


At 0mmHg

277px
view width

100.0%
of baseline



'Middle finger' method at 0mmHg
with M. nitrile glove, Trial #2
Filename: 20220423_143310.jpg
Size: 4032x2268px (reduced 10x to display here)

At 40mmHg

210px
view width

75.8%
of baseline



'Middle finger' method at 0mmHg
with M. nitrile glove, Trial #2
Filename: 20220423_143324.jpg
Size: 4032x2268px (reduced 10x to display here)

At 80mmHg

146px
view width

52.7%
of baseline



'Middle finger' method at 0mmHg
with M. nitrile glove, Trial #2
Filename: 20220423_143336.jpg
Size: 4032x2268px (reduced 10x to display here)

At 120mmHg

116px
view width

41.9%
of baseline




'Middle finger' method at 0mmHg
with M. nitrile glove, Trial #2
Filename: 20220423_143346.jpg
Size: 4032x2268px (reduced 10x to display here)

At 160mmHg

84px
view width

30.3%
of baseline



'Middle finger' method at 0mmHg
with M. nitrile glove, Trial #2
Filename: 20220423_143355.jpg
Size: 4032x2268px (reduced 10x to display here)

At 200mmHg

67px
view width

24.2%
of baseline



'Middle finger' method at 0mmHg
with M. nitrile glove, Trial #2
Filename: 20220423_143401.jpg
Size: 4032x2268px (reduced 10x to display here)