


At 0mmHg

257px
view width

100.0%
of baseline




'Middle finger' method at 0mmHg
with M. nitrile glove, Trial #1
Filename: 20220423_142023.jpg
Size: 4032x2268px (reduced 10x to display here)

At 40mmHg

190px
view width

73.9%
of baseline




'Middle finger' method at 40mmHg
with M. nitrile glove, Trial #1
Filename: 20220423_142031.jpg
Size: 4032x2268px (reduced 10x to display here)

At 80mmHg

120px
view width

46.7%
of baseline



'Middle finger' method at 80mmHg
with M. nitrile glove, Trial #1
Filename: 20220423_142043.jpg
Size: 4032x2268px (reduced 10x to display here)

At 120mmHg

90px
view width

35.0%
of baseline



'Middle finger' method at 120mmHg
with M. nitrile glove, Trial #1
Filename: 20220423_142054.jpg
Size: 4032x2268px (reduced 10x to display here)

At 160mmHg

65px
view width

25.3%
of baseline



'Middle finger' method at 160mmHg
with M. nitrile glove, Trial #1
Filename: 20220423_142100.jpg
Size: 4032x2268px (reduced 10x to display here)

At 200mmHg

44px
view width

17.1%
of baseline



'Middle finger' method at 200mmHg
with M. nitrile glove, Trial #1
Filename: 20220423_142108.jpg
Size: 4032x2268px (reduced 10x to display here)