

Palindromic, Polynomially-Growing Free-by-Cyclic Groups are CAT(0)

Rylee Lyman – Tufts University

Graduate Student Conference in Algebra, Geometry and Topology,
Temple University, June 1, 2019

A Tale of Two Free-by-Cyclic Groups

Write $F_3\langle a, b, c \rangle$. Let φ and ψ in $\text{Aut}(F_3)$ be the automorphisms

$$\varphi \begin{cases} a \mapsto a \\ b \mapsto ba \\ c \mapsto ca^2 \end{cases} \quad \psi \begin{cases} a \mapsto a \\ b \mapsto aba \\ c \mapsto a^2ca^2 \end{cases}$$

A Tale of Two Free-by-Cyclic Groups


Write $F_3\langle a, b, c \rangle$. Let φ and ψ in $\text{Aut}(F_3)$ be the automorphisms

$$\varphi \begin{cases} a \mapsto a \\ b \mapsto ba \\ c \mapsto ca^2 \end{cases} \quad \psi \begin{cases} a \mapsto a \\ b \mapsto aba \\ c \mapsto a^2ca^2 \end{cases}$$

$$G_\varphi = F_3 \rtimes_\varphi \mathbb{Z} = \langle F_3, t \mid txt^{-1} = \varphi(x), x \in F_3 \rangle$$


$$G_\psi = F_3 \rtimes_\psi \mathbb{Z} = \langle F_3, t \mid txt^{-1} = \psi(x), x \in F_3 \rangle$$

Free-by-Cyclic Groups as Mapping Tori



$$G_\varphi = \pi_1 \left((\mathbb{S}^1 \times \{0\}) \bigvee_{(\rho, 0) \sim (\varphi(\rho), 1)} \right)$$

Free-by-Cyclic Groups as Mapping Tori



$$G_\varphi = \pi_1 \left((\mathcal{G} \times \{0\}) \middle/_{(p,0) \sim (\varphi(p),1)} \right)$$

Although φ and ψ appear similar, G_φ and G_ψ turn out to have very different geometry!

Motivation

Theorem (Gersten, '94)

$\text{Aut}(F_n)$, $\text{Out}(F_{n+1})$ are not CAT(0) groups for $n \geq 3$

Key observation: G_φ cannot be a subgroup of a CAT(0) group, but $G_\varphi \leq \text{Aut}(F_n)$ and $\text{Out}(F_{n+1})$ for $n \geq 3$.

Motivation

Theorem (Gersten, '94)

$\text{Aut}(F_n)$, $\text{Out}(F_{n+1})$ are not CAT(0) groups for $n \geq 3$

Key observation: G_φ cannot be a subgroup of a CAT(0) group, but $G_\varphi \leq \text{Aut}(F_n)$ and $\text{Out}(F_{n+1})$ for $n \geq 3$.

Original Goal Theorem

$\text{Aut}(W_n)$, $\text{Out}(W_{n+1})$ are not CAT(0) groups for $n \geq 4$.

Motivation

Theorem (Gersten, '94)

$\text{Aut}(F_n)$, $\text{Out}(F_{n+1})$ are not CAT(0) groups for $n \geq 3$

Key observation: G_φ cannot be a subgroup of a CAT(0) group, but $G_\varphi \leq \text{Aut}(F_n)$ and $\text{Out}(F_{n+1})$ for $n \geq 3$.

Original Goal Theorem

$\text{Aut}(W_n)$, $\text{Out}(W_{n+1})$ are not CAT(0) groups for $n \geq 4$.

Hopeful tool: Use G_ψ or a similar group to mimic Gersten's proof.

Motivation

Theorem (Gersten, '94)

$\text{Aut}(F_n), \text{Out}(F_{n+1})$ are not CAT(0) groups for $n \geq 3$

Key observation: G_φ cannot be a subgroup of a CAT(0) group, but $G_\varphi \leq \text{Aut}(F_n)$ and $\text{Out}(F_{n+1})$ for $n \geq 3$.

Original Goal Theorem

$\text{Aut}(W_n), \text{Out}(W_{n+1})$ are not CAT(0) groups for $n \geq 4$.

Hopeful tool: Use G_ψ or a similar group to mimic Gersten's proof.

Actually, this is still wide open!

Results

Theorem (L, '19)

We fail! G_ψ is a CAT(0) group. More generally, any polynomially-growing $\theta \in \text{Aut}(W_n)$ has a power $k < n!$ such that G_{θ^k} is a CAT(0) group.

Results

Theorem (L, '19)

We fail! G_ψ is a CAT(0) group. More generally, any polynomially-growing $\theta \in \text{Aut}(W_n)$ has a power $k < n!$ such that G_{θ^k} is a CAT(0) group.

Our proof is constructive: We build a 2-dimensional CW complex X_{θ^k} with a CAT(0) metric that G_{θ^k} acts on.

Results

Theorem (L, '19)

We fail! G_ψ is a CAT(0) group. More generally, any polynomially-growing $\theta \in \text{Aut}(W_n)$ has a power $k < n!$ such that G_{θ^k} is a CAT(0) group.

Our proof is constructive: We build a 2-dimensional CW complex X_{θ^k} with a CAT(0) metric that G_{θ^k} acts on.

The k is needed to ensure that θ^k is in the kernel of the map $\text{Aut}(W_n) \rightarrow S_n$. This restriction isn't good enough for Gersten's theorem.

Background

What is a W_n , anyway?

Let $W_n = \underbrace{\mathbb{Z}/2\mathbb{Z} * \cdots * \mathbb{Z}/2\mathbb{Z}}_{n \text{ times}} = \langle a_1, \dots, a_n \mid a_i^2 \rangle$.

Background

What is a W_n , anyway?

Let $W_n = \underbrace{\mathbb{Z}/2\mathbb{Z} * \cdots * \mathbb{Z}/2\mathbb{Z}}_{n \text{ times}} = \langle a_1, \dots, a_n \mid a_i^2 \rangle$.

$W_n \cong F_{n-1} \rtimes \mathbb{Z}/2\mathbb{Z}$. We get $\text{Aut}(W_n) \hookrightarrow \text{Aut}(F_{n-1})$, and $\text{Out}(W_n) \hookrightarrow \text{Out}(F_{n-1})$.

$$W_4 = \pi_1 \left(\begin{array}{c} \mathbb{Z}/2\mathbb{Z} \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ \mathbb{Z}/2\mathbb{Z} \end{array} \right)$$

$$F_3 = \pi_1 \left(\begin{array}{c} \text{flower shape} \\ \bullet \end{array} \right)$$

Background

Facts about $\text{Out}(F_n)$

Given $\Phi \in \text{Out}(F_n)$ and $[g]$ a conjugacy class in F_n , the word length of the shortest representative of $\Phi^k([g])$ grows either polynomially or exponentially.

Background

Facts about $\text{Out}(F_n)$

Given $\Phi \in \text{Out}(F_n)$ and $[g]$ a conjugacy class in F_n , the word length of the shortest representative of $\Phi^k([g])$ grows either polynomially or exponentially.

Φ is *exponentially-growing* if at least one conjugacy class grows exponentially, and is *polynomially-growing* otherwise.

Examples: φ and ψ are polynomially (in fact linearly) growing.

Background

Facts about $\text{Out}(F_n)$

Given $\Phi \in \text{Out}(F_n)$ and $[g]$ a conjugacy class in F_n , the word length of the shortest representative of $\Phi^k([g])$ grows either polynomially or exponentially.

Φ is *exponentially-growing* if at least one conjugacy class grows exponentially, and is *polynomially-growing* otherwise.

Examples: φ and ψ are polynomially (in fact linearly) growing.

Pseudo-Anosov homeomorphisms of surfaces with nonempty boundary induce exponentially-growing $\Phi \in \text{Out}(F_n)$.

Bakground

Facts about $\text{Out}(F_n)$

(Up to finite index) elements in the image $\text{Out}(W_n) \hookrightarrow \text{Out}(F_{n-1})$ are *palindromic*—they send generators to palindromes.

Bakground

Facts about $\text{Out}(F_n)$

(Up to finite index) elements in the image $\text{Out}(W_n) \hookrightarrow \text{Out}(F_{n-1})$ are *palindromic*—they send generators to palindromes.

Example: ψ is palindromic!

Bakground

Facts about $\text{Out}(F_n)$

(Up to finite index) elements in the image $\text{Out}(W_n) \hookrightarrow \text{Out}(F_{n-1})$ are *palindromic*—they send generators to palindromes.

Example: ψ is palindromic!

Our theorem is one of the first signs of a difference in structure between $\Phi \in \text{Out}(W_n)$ and those in $\text{Out}(F_n)$.

Background

Facts about $\text{Out}(F_n)$

Exponentially-growing Φ feel a little more like mapping classes, while polynomially-growing Φ are more like unipotent matrices.

Background

Facts about $\text{Out}(F_n)$

Exponentially-growing Φ feel a little more like mapping classes, while polynomially-growing Φ are more like unipotent matrices.

Polynomially-growing $F_n \rtimes \mathbb{Z}$ admit a *hierarchy*, with all edge groups \mathbb{Z} and vertex groups either $F_m \rtimes \mathbb{Z}$ with $m < n$, or (at the base) \mathbb{Z}^2 .

Background

Facts about $\text{Out}(F_n)$


Exponentially-growing Φ feel a little more like mapping classes, while polynomially-growing Φ are more like unipotent matrices.

Polynomially-growing $F_n \rtimes \mathbb{Z}$ admit a *hierarchy*, with all edge groups \mathbb{Z} and vertex groups either $F_m \rtimes \mathbb{Z}$ with $m < n$, or (at the base) \mathbb{Z}^2 .


$$G_\varphi = \pi_1 \left(\begin{array}{c} \overset{\alpha^b}{\curvearrowright} \\ \overset{\beta^b}{\curvearrowright} \\ F_2 \rtimes \mathbb{Z} \\ \langle a, b, t \rangle \end{array} \right)$$
$$\langle a, b, t \rangle = \pi_1 \left(\begin{array}{c} \overset{\alpha^t}{\curvearrowright} \\ \overset{\beta^t}{\curvearrowright} \\ \mathbb{Z}^2 \\ \langle a, b, t \rangle \end{array} \right)$$

$$G_\varphi = \pi_1 \left(\begin{array}{c} \overset{\alpha^t}{\curvearrowright} \\ \overset{\beta^t}{\curvearrowright} \\ F_2 \rtimes \mathbb{Z} \\ \langle a, b, t \rangle \end{array} \right)$$
$$\langle a, b, t \rangle = \pi_1 \left(\begin{array}{c} \overset{\alpha^t}{\curvearrowright} \\ \overset{\beta^t}{\curvearrowright} \\ \mathbb{Z}^2 \\ \langle a, b, t \rangle \end{array} \right)$$

CAT(0) Geometry



\mathbb{R}^2



$$d_X(p, q) \leq d_{\mathbb{R}^2}(\tilde{p}, \tilde{q})$$

CAT(0) Geometry

The Flat Torus Theorem

Theorem (Flat Torus Theorem; Bridson–Haefliger '99)

Given a “nice” action of G on a CAT(0) space X and $\mathbb{Z}^k \cong A \leq G$, there is an isometric copy of $\mathbb{R}^k \subset X$ that A acts on by translation, and the quotient \mathbb{R}^k/A is a k -torus.

CAT(0) Geometry


The Flat Torus Theorem

$$G \curvearrowright X$$

$$A \cong \mathbb{Z}^k \quad A \leq G$$



$$\mathbb{R}^k \subseteq X$$




$$\mathbb{R}^k / A$$




$$T^k$$

Gersten: G_φ is not CAT(0)


$$G_\varphi: b^{-1}tb = at, \quad c^{-1}tc = a^2t$$




G_ψ fails to fail to be CAT(0)



Blowing up the Bass–Serre Tree



Blowing up the Bass–Serre Tree



Blowing up the Bass–Serre Tree

