Exact stress-tensor flow in four-dimensional single-scalar Dirac–Born–Infeld theory

Ryosuke Suda

Graduate School of Science and Engineering, Saitama University,
255 Shimo-Okubo, Sakura-ku, Saitama 338-8570, Japan
r.suda.813@ms.saitama-u.ac.jp

Abstract. We consider four-dimensional Dirac–Born–Infeld theory containing one Abelian gauge field and one real scalar field, and derive a flow equation that expresses the derivative of the Lagrangian with respect to the deformation parameter solely in terms of local invariants of the stress tensor. For a non-null scalar gradient, the flow admits a closed rational expression. The same expression follows both from a direct calculation in the four-dimensional DBI theory and from dimensional reduction of the known five-dimensional Born–Infeld flow. When the scalar gradient is null, the rational expression degenerates and is replaced by a finite stress-tensor formula; in the limit where the scalar is removed, this formula reduces to the known four-dimensional Born–Infeld flow. We also derive an algebraic relation among the stress-tensor invariants realized by the DBI theory. The closed form was first suggested by a neural-network-assisted numerical search and is established analytically in the paper.

Keywords: Dirac–Born–Infeld theory; non-linear electrodynamics; stress-tensor deformation; D-branes

Draft — August 2026

1 Introduction

Non-linear electrodynamics provides a broad setting in which to study electromagnetic duality, causality, and wave propagation beyond Maxwell theory. Born–Infeld (BI) theory is the canonical example. More recently, the structure of four-dimensional non-linear electrodynamics (NLED) has received renewed attention through the ModMax theory, which preserves both conformal invariance and electromagnetic duality, its Born–Infeld-type extensions, and classifications of theories without birefringence [1, 4, 5, 9, 10].

In parallel, deformations generated by local composite operators built from the stress tensor have been studied extensively. Their prototype is the two-dimensional $T\overline{T}$ deformation, an irrelevant deformation that nevertheless retains substantial analytic control and, in suitable cases, preserves integrability and allows exact access to finite-volume spectra [2]. There is no unique higher-dimensional analogue of $T\overline{T}$, but flow equations in which the variation of a Lagrangian with respect to a deformation parameter is expressed in terms of stress-tensor invariants have been identified for BI theory, ModMax–BI theory, and brane actions, among other examples [3, 7, 8, 10, 13].

The situation is more involved for Dirac–Born–Infeld (DBI) theories containing both a gauge field and worldvolume scalars. In generic dimension, the derivative of the DBI Lagrangian with respect to the deformation parameter contains invariants built explicitly from the field strength and scalar gradients, and these quantities need not be reconstructible from the stress tensor alone. Blair, Lahnsteiner, Obers and Yan studied flows generating Dp-brane DBI actions and obtained complete flow equations including the gauge field for $p = 1, 2$, while for the four-dimensional worldvolume case $p = 3$ they gave the result only to first order in the deformation parameter [12]. Brizio, Kade, Sfondrini and Sorokin derived stress-tensor flows for BI theory in general dimension and analyzed DBI theories in two and three dimensions in detail [13]. They identified the extension to $d \geq 4$ DBI theories as an open problem and pointed out that the four-dimensional theory with a single real scalar is a particularly promising case in which a formulation entirely in terms of stress-tensor invariants may exist. Relatedly, Ferko, Hu, Huang, Koutrolikos and Tartaglino-Mazzucchelli constructed stress-tensor-like flows for three-dimensional Born–Infeld theory and for the scalar sector of DBI theory, and identified the construction of a flow for the full gauge-plus-scalar DBI action as a natural open direction [11].

In this paper we address this four-dimensional single-scalar DBI theory. Before carrying out the analytic calculation, we numerically regressed the derivative of the DBI Lagrangian with respect to the deformation parameter against four invariants constructed from the stress tensor on the same field configurations. A neural network reproduced this quantity with a small error on independent test data, indicating that the derivative may be a function of the stress-tensor invariants alone. A simple rational ansatz applied to independent numerical data then produced a candidate closed form.

Guided by this numerical observation, we derive the flow analytically in four dimensions. Intermediate expressions contain invariants built from the field strength and the scalar gradient, but identities specific to four-dimensional antisymmetric matrices allow these quantities to be eliminated in favor of stress-tensor invariants. We then derive the same four-dimensional flow by reducing the known five-dimensional Born–Infeld flow. The formula obtained for a non-null scalar gradient becomes indeterminate when the scalar gradient is null; we therefore treat this locus separately and recover the pure Born–Infeld flow as a limiting case. The details of the numerical search are collected in appendix A; none of the analytic derivations below assumes the numerically inferred formula.

All flow equations considered below are classical Lagrangian flow equations; no quantum definition of the four-dimensional deformation is assumed.

The paper is organized as follows. Section 2 fixes the conventions for the four-dimensional DBI theory and its stress tensor. In section 3 we derive the four-dimensional flow equation directly and compare its weak-deformation limit with the perturbative $p = 3$ result of ref. [12]. Section 4

derives the same relation by dimensional reduction from five-dimensional Born–Infeld theory. Section 5 treats null scalar gradients and the Born–Infeld limit. In section 6 we derive an algebraic relation among the stress-tensor invariants realized by the DBI theory. Section 7 summarizes the results and discusses their interpretation and possible extensions.

2 Four-dimensional DBI theory

We first fix conventions for four-dimensional DBI theory with a single real scalar. This section introduces the action and the Hilbert stress tensor. Since the characteristic polynomial of the stress tensor will be used below, we distinguish throughout between the covariant tensor $T_{\mu\nu}$ and the mixed tensor $T^{\mu}{}_{\nu}$.

2.1 Conventions and action

We use metric signature $(-,+,+,+)$. In canonical four-dimensional normalization the deformation parameter has mass dimension $[\lambda] = -4$, and we take $\lambda > 0$ in order to describe the real DBI branch. The algebraic formulas below may be analytically continued wherever the relevant square roots remain defined. The Abelian gauge potential is denoted by A_μ, with field strength

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu. \tag{2.1}$$

We denote the real scalar field by ϕ. The four-dimensional DBI action is

$$I_\lambda = \int \mathrm{d}^4x\sqrt{-g}\,\mathcal{L}_\lambda, \qquad \mathcal{L}_\lambda = \frac{1-\Delta}{\lambda}, \tag{2.2}$$

where

$$\mathcal{G}_{\mu\nu} = g_{\mu\nu} + \lambda\,\partial_\mu\phi\,\partial_\nu\phi + \sqrt{\lambda}F_{\mu\nu}, \qquad \Delta = \sqrt{\frac{\det\mathcal{G}}{\det g}}. \tag{2.3}$$

The branch is chosen so that $\Delta \to 1$ as $\lambda \to 0$.

We write the Lorentz norm of the scalar gradient as

$$X = \partial_\mu\phi\,\partial^\mu\phi. \tag{2.4}$$

For later use, we also introduce the standard NLED invariant

$$S = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu}. \tag{2.5}$$

Expanding eq. (2.3) around $\lambda = 0$ gives

$$\mathcal{L}_\lambda = S - \frac{1}{2}X + O(\lambda). \tag{2.6}$$

Thus the seed theory at $\lambda = 0$ consists of a Maxwell field and a free massless scalar.

2.2 Stress tensor

To formulate the flow in terms of stress-tensor invariants, we first determine the Hilbert stress tensor of the DBI action. We define

$$T_{\mu\nu} = -\frac{2}{\sqrt{-g}}\frac{\delta I_\lambda}{\delta g^{\mu\nu}}. \tag{2.7}$$

Metric variation of eq. (2.2) gives

$$T^{\mu\nu} = \frac{1}{\lambda}\left[g^{\mu\nu} - \Delta(\mathcal{G}^{-1})^{(\mu\nu)}\right], \qquad (\mathcal{G}^{-1})^{(\mu\nu)} = \frac{1}{2}\left[(\mathcal{G}^{-1})^{\mu\nu} + (\mathcal{G}^{-1})^{\nu\mu}\right]. \tag{2.8}$$

All matrix invariants of the stress tensor below are taken for the mixed tensor

$$T^{\mu}{}_{\nu} = g^{\mu\rho}T_{\rho\nu}. \tag{2.9}$$

Equation (2.8) will be the starting point for the calculation below. In the next section we compute the derivative of the DBI Lagrangian on the same field configuration and eliminate the explicit field variables in favor of invariants of $T^{\mu}{}_{\nu}$.

3 Four-dimensional flow equation

In this section we differentiate the four-dimensional DBI Lagrangian with respect to the deformation parameter while holding the metric, field strength, and scalar gradient fixed, and then express the result in terms of stress-tensor invariants. We first introduce the relevant invariants and briefly describe the numerical search that motivated the analytic calculation. We then derive the flow directly from the DBI Lagrangian without assuming the numerically inferred expression.

We define

$$Q_\lambda \equiv \left.\frac{\partial \mathcal{L}_\lambda}{\partial \lambda}\right|_{g_{\mu\nu},\, F_{\mu\nu},\, \partial_\mu\phi}. \tag{3.1}$$

The goal is to express Q_λ solely through local invariants of the stress tensor. Introduce

$$M^{\mu}{}_{\nu} = \delta^{\mu}{}_{\nu} - \lambda T^{\mu}{}_{\nu}. \tag{3.2}$$

For a four-dimensional matrix N, define the elementary symmetric polynomials $e_k(N)$ by

$$\det(z\mathbf{1} - N) = z^4 - e_1(N)z^3 + e_2(N)z^2 - e_3(N)z + e_4(N), \tag{3.3}$$

and write $E_k = e_k(M)$.

The numerical search was carried out in the region where $X \neq 0$ and the scalar gradient is timelike. Random DBI field configurations were used to compute (E_1, E_2, E_3, E_4) and $\lambda^2 Q_\lambda$, and a neural network was trained to regress the map

$$(E_1, E_2, E_3, E_4) \longmapsto \lambda^2 Q_\lambda. \tag{3.4}$$

Its small error on independent test data suggested that Q_λ could be written as a function of these invariants alone. The network architecture, loss function, and rational readout are described in appendix A. We now proceed analytically and do not assume this numerical result.

For an antisymmetric 4×4 matrix $F^{\mu}{}_{\nu}$, the Cayley–Hamilton identity reduces to a form involving the two standard four-dimensional electromagnetic invariants. In addition to S, define

$$P = -\frac{1}{4}F_{\mu\nu}\widetilde{F}^{\mu\nu}, \qquad \widetilde{F}^{\mu\nu} = \frac{1}{2\sqrt{-g}}\varepsilon^{\mu\nu\rho\sigma}F_{\rho\sigma}, \qquad \varepsilon^{0123} = +1. \tag{3.5}$$

Then

$$(\sqrt{\lambda}F)^4 - 2\lambda S(\sqrt{\lambda}F)^2 - \lambda^2 P^2 \mathbf{1} = 0. \tag{3.6}$$

We will also use the matrix determinant lemma

$$\det(A + uv^{\mathsf{T}}) = \det A \left(1 + v^{\mathsf{T}}A^{-1}u\right) \tag{3.7}$$

for an invertible matrix A and column vectors u, v.

Applying these relations to eq. (2.8) gives first

$$E_4 = \det M = 1 + \lambda X. \tag{3.8}$$

The intermediate expressions for E_1 and E_3 involve the mixed gauge–scalar invariant

$$Z = \partial_\mu\phi\, F^{\mu}{}_{\rho}F^{\rho}{}_{\nu}\, \partial^\nu\phi, \tag{3.9}$$

which does not appear in the final flow equation. A direct application of eqs. (3.6) and (3.7) to the DBI matrix yields

$$\Delta^2 = 1 + \lambda X - 2\lambda S + \lambda^2 \left(Z - 2SX - P^2\right), \tag{3.10}$$

$$\Delta E_1 = 4 + 3\lambda X - 4\lambda S + \lambda^2 (Z - 2SX), \tag{3.11}$$

$$\Delta E_3 = (1 + \lambda X)(4 + \lambda X - 4\lambda S) + \lambda^2 Z. \tag{3.12}$$

The matrix algebra leading to these formulas is given in appendix B.

On the other hand, eq. (2.2) implies

$$\lambda^2 Q_\lambda = \Delta - 1 - \lambda \frac{\partial \Delta}{\partial \lambda}. \tag{3.13}$$

Differentiating eq. (3.10) gives

$$2\Delta \left(1 + \lambda^2 Q_\lambda\right) = 2 + \lambda X - 2\lambda S. \tag{3.14}$$

Taking the difference of eqs. (3.11) and (3.12) and using eq. (3.8), we obtain

$$E_1 - E_3 = 2(1 - E_4) \left(1 + \lambda^2 Q_\lambda\right). \tag{3.15}$$

This relation remains valid at $X = 0$ and will also be used in section 5.

For $X \neq 0$, eq. (3.8) implies $E_4 \neq 1$, so eq. (3.15) can be solved for Q_λ:

$$\lambda^2 Q_\lambda = \frac{E_1 - E_3 + 2E_4 - 2}{2(1 - E_4)}. \tag{3.16}$$

To rewrite this entirely in terms of $T^\mu{}_\nu$, use $M = \mathbf{1} - \lambda T$. With $e_1(T) = \operatorname{tr} T$ and $e_4(T) = \det T$,

$$\begin{aligned} E_1 &= 4 - \lambda \operatorname{tr} T, \\ E_3 &= 4 - 3\lambda \operatorname{tr} T + 2\lambda^2 e_2(T) - \lambda^3 e_3(T), \\ E_4 &= 1 - \lambda \operatorname{tr} T + \lambda^2 e_2(T) - \lambda^3 e_3(T) + \lambda^4 \det T. \end{aligned} \tag{3.17}$$

It follows that

$$Q_\lambda = \frac{2\lambda \det T - e_3(T)}{2\left[\operatorname{tr} T - \lambda e_2(T) + \lambda^2 e_3(T) - \lambda^3 \det T\right]}. \tag{3.18}$$

Moreover, comparing eqs. (3.8) and (3.17) gives

$$\operatorname{tr} T - \lambda e_2(T) + \lambda^2 e_3(T) - \lambda^3 \det T = -X. \tag{3.19}$$

Thus eq. (3.18) applies for $X \neq 0$, while the locus $X = 0$ must be treated separately.

As a first consistency check, consider the weak-deformation limit. For a seed configuration with $X \neq 0$, eq. (3.18) gives

$$Q_0 = -\frac{e_3(T_0)}{2 \operatorname{tr} T_0}, \tag{3.20}$$

where T_0 is the stress tensor of the Maxwell–scalar theory in eq. (2.6).

Blair, Lahnsteiner, Obers and Yan denote their deformation parameter by t [12]. Restricting their flat-worldvolume DBI Lagrangian to $p = 3$ and one real scalar, and identifying $t = \lambda$ and $X^{A'} = \phi$, gives the normalization in eq. (2.2) after extracting the overall factor t^{-1} from the four-dimensional determinant. In these conventions, the t^0 term in their $p = 3$ flow can be written as

$$Q_0 = \frac{1}{4} \operatorname{tr}(T_0^2) - \frac{1}{8} (\operatorname{tr} T_0)^2 - \frac{1}{8} \left[\operatorname{tr}(F^4) - \frac{1}{4} \left(\operatorname{tr} F^2\right)^2\right], \tag{3.21}$$

where F denotes the mixed matrix $F^{\mu}{}_{\nu}$. Substituting the Maxwell–scalar stress tensor shows that eqs. (3.20) and (3.21) agree. At $X = 0$, the weak-deformation limit should instead be taken from the expression derived in section 5.

Equation (3.18) therefore resums the deformation parameter to all orders, while reproducing the known perturbative $p = 3$ result at weak deformation. The derivation above uses only the four-dimensional DBI Lagrangian. Since the DBI action itself arises by dimensional reduction of higher-dimensional Born–Infeld theory, it is natural to ask whether the flow equation behaves in the same way. We address this next.

4 Five-dimensional Born–Infeld reduction

The previous section derived the flow directly in four dimensions. We now obtain the same expression from the known five-dimensional Born–Infeld flow. A gauge-potential component along a compact direction becomes a scalar field after reduction, so reducing five-dimensional Born–Infeld theory on one direction produces precisely the four-dimensional DBI system considered above. We first verify the reduction at the level of the action and then apply it to the flow equation.

Five-dimensional quantities will carry hats. Let the compact coordinate be $y \sim y + L_y$. We do not consider a general Kaluza–Klein reduction: throughout this section we take a product background with $\widehat{g}_{\mu y} = 0$ and $\widehat{g}_{yy} = 1$, omit fluctuations of the Kaluza–Klein vector and radion, and retain only zero modes, so all fields are independent of y.

It is useful to distinguish the canonically normalized deformation parameters before and after reduction. Denote the five-dimensional parameter by $\widehat{\lambda}$ and define the four-dimensional zero modes by

$$\lambda = \frac{\widehat{\lambda}}{L_y}, \qquad A_\mu = \sqrt{L_y}\,\widehat{A}_\mu, \qquad \phi = \sqrt{L_y}\,\widehat{A}_y. \tag{4.1}$$

Thus $[\widehat{\lambda}] = -5$ and $[\lambda] = -4$ in canonical normalization. The field strengths obey $\widehat{F}_{\mu\nu} = F_{\mu\nu}/\sqrt{L_y}$ and $\widehat{F}_{\mu y} = \partial_\mu\phi/\sqrt{L_y}$.

We take

$$\widehat{g}_{MN} = \begin{pmatrix} g_{\mu\nu} & 0 \\ 0 & 1 \end{pmatrix}, \qquad \widehat{\mathcal{G}}_{MN} = \widehat{g}_{MN} + \sqrt{\widehat{\lambda}}\,\widehat{F}_{MN}. \tag{4.2}$$

The five-dimensional action and determinant factor are

$$\widehat{I}_{\widehat{\lambda}} = \int \mathrm{d}^5x\sqrt{-\widehat{g}}\,\widehat{\mathcal{L}}_{\widehat{\lambda}}, \qquad \widehat{\mathcal{L}}_{\widehat{\lambda}} = \frac{1-\widehat{\Delta}}{\widehat{\lambda}}, \qquad \widehat{\Delta} = \sqrt{\frac{\det\widehat{\mathcal{G}}}{\det\widehat{g}}}, \tag{4.3}$$

and the Hilbert stress tensor is

$$\widehat{T}_{MN} = -\frac{2}{\sqrt{-\widehat{g}}}\frac{\delta\widehat{I}_{\widehat{\lambda}}}{\delta\widehat{g}^{MN}}, \qquad \widehat{T}^{M}{}_{N} = \widehat{g}^{MP}\widehat{T}_{PN}. \tag{4.4}$$

Using eq. (4.1), the zero-mode matrix takes the form

$$\widehat{\mathcal{G}} = \begin{pmatrix} g + \sqrt{\lambda}F & \sqrt{\lambda}\,\partial\phi \\ -\sqrt{\lambda}\,(\partial\phi)^{\mathsf{T}} & 1 \end{pmatrix}. \tag{4.5}$$

Since the lower-right block is unity, we may use

$$\det\begin{pmatrix} A & b \\ c^{\mathsf{T}} & 1 \end{pmatrix} = \det\left(A - bc^{\mathsf{T}}\right). \tag{4.6}$$

Applied to eq. (4.5), this gives

$$\det\widehat{\mathcal{G}} = \det\left(g + \sqrt{\lambda}F + \lambda\,\partial\phi\,(\partial\phi)^{\mathsf{T}}\right) = \det\mathcal{G}. \tag{4.7}$$

After integrating over y, the factor L_y combines with $\widehat{\lambda} = L_y\lambda$, so that $\widehat{I}_{\widehat{\lambda}} = I_\lambda$ with the four-dimensional action in eq. (2.2). Metric variation similarly gives $T^\mu{}_\nu = L_y \widehat{T}^\mu{}_\nu$ for the four-dimensional block, and hence $\widehat{\lambda}\,\widehat{T}^\mu{}_\nu = \lambda T^\mu{}_\nu$.

The inverse matrix needed for reducing the flow follows from the same Schur complement:

$$\widehat{\mathcal{G}}^{-1} = \begin{pmatrix} \mathcal{G}^{-1} & -\sqrt{\lambda}\,\mathcal{G}^{-1}\partial\phi \\ \sqrt{\lambda}\,(\partial\phi)^{\mathsf{T}}\mathcal{G}^{-1} & 1-\lambda(\partial\phi)^{\mathsf{T}}\mathcal{G}^{-1}\partial\phi \end{pmatrix}. \tag{4.8}$$

As in four dimensions, define

$$\widehat{M}^M{}_N \equiv \delta^M{}_N - \widehat{\lambda}\,\widehat{T}^M{}_N = \widehat{\Delta}\,(\widehat{\mathcal{G}}^{-1})^{(MP)}\widehat{g}_{PN}. \tag{4.9}$$

Equation (4.7) implies $\widehat{\Delta} = \Delta$, so the trace and determinant of $\widehat{M}$ can be expressed directly through four-dimensional quantities.

The known five-dimensional Born–Infeld flow is [13]

$$\frac{\partial\widehat{\mathcal{L}}_{\widehat{\lambda}}}{\partial\widehat{\lambda}} = \frac{1}{4\widehat{\lambda}^2} - \frac{1}{4\widehat{\lambda}}\operatorname{tr}\widehat{T} - \frac{1}{4\widehat{\lambda}^2}\det(\mathbf{1}_5 - \widehat{\lambda}\widehat{T}). \tag{4.10}$$

Since $\mathcal{L}_\lambda = L_y\widehat{\mathcal{L}}_{\widehat{\lambda}}$ and $\widehat{\lambda} = L_y\lambda$ at fixed L_y and fixed four-dimensional fields,

$$\lambda^2 Q_\lambda = \widehat{\lambda}^2\frac{\partial\widehat{\mathcal{L}}_{\widehat{\lambda}}}{\partial\widehat{\lambda}}. \tag{4.11}$$

Using eqs. (4.8), (4.9), and the four-dimensional stress tensor in eq. (2.8), one finds

$$\det(\mathbf{1}_5 - \widehat{\lambda}\widehat{T}) = \Delta, \tag{4.12}$$

$$\operatorname{tr}(\mathbf{1}_5 - \widehat{\lambda}\widehat{T}) = E_1 + \Delta\left[1 - \lambda\,\partial_\mu\phi(\mathcal{G}^{-1})^{(\mu\nu)}\partial_\nu\phi\right]. \tag{4.13}$$

Equations (4.10)–(4.13) then reduce to

$$\lambda^2 Q_\lambda = \frac{1}{4}\left[E_1 - 4 - \Delta\lambda\,\partial_\mu\phi(\mathcal{G}^{-1})^{(\mu\nu)}\partial_\nu\phi\right]. \tag{4.14}$$

The remaining scalar contraction can be eliminated in favor of four-dimensional stress-tensor invariants. Combining eqs. (3.11), (3.12), and (3.8) gives

$$(E_4 - 1)\Delta\lambda\,\partial_\mu\phi(\mathcal{G}^{-1})^{(\mu\nu)}\partial_\nu\phi = E_1(1 + E_4) - 2E_3. \tag{4.15}$$

For $X \neq 0$, substituting eq. (4.15) into eq. (4.14) reproduces eq. (3.16). This derivation does not differentiate the four-dimensional DBI Lagrangian with respect to λ and therefore provides a second route to the same four-dimensional flow from the higher-dimensional Born–Infeld flow.

Both derivations divide by $1 - E_4$ in the final step. According to eq. (3.8), this factor vanishes precisely at $X = 0$. We therefore analyze this locus separately.

5 Null scalar gradient

The expression in eq. (3.16) was derived under the assumption $X \neq 0$. Equation (3.8) shows that $X = 0$ is equivalent to $E_4 = 1$, in which case both numerator and denominator of eq. (3.16) vanish. To obtain a finite expression at $X = 0$, we return to the undivided relation in eq. (3.15) and use the characteristic polynomial of M.

For four-dimensional DBI theory with a single real scalar, the characteristic polynomial of M factorizes into two quadratics, as shown in appendix B. On the locus $X = 0$, this factorization gives

$$\lambda^2 Q_\lambda = \frac{1}{4}\left[E_1 + \sqrt{E_1^2 - 4E_2 + 8}\right] - 1. \tag{5.1}$$

The sign is fixed by continuity from the DBI branch with $X \neq 0$. Rewriting E_1 and E_2 in terms of stress-tensor traces gives

$$Q_\lambda = \frac{-\lambda \operatorname{tr} T + \sqrt{4\lambda \operatorname{tr} T + \lambda^2 \left[2 \operatorname{tr}(T^2) - (\operatorname{tr} T)^2\right]}}{4\lambda^2}. \tag{5.2}$$

If the scalar gradient is further set to zero, the theory reduces to four-dimensional Born–Infeld electrodynamics. Equation (5.1) then simplifies to

$$Q_\lambda = -\frac{1}{4\lambda} \operatorname{tr} T, \tag{5.3}$$

in agreement with the known four-dimensional Born–Infeld flow [7, 13]. Thus a stress-tensor-only flow also exists at $X = 0$, and the pure Born–Infeld result is recovered continuously.

Having treated both $X \neq 0$ and $X = 0$, we now examine the algebraic relations among the invariants E_1, E_2, E_3, E_4 realized by DBI configurations.

6 Algebraic relation among stress-tensor invariants

The flow has been written in terms of $E_k = e_k(\mathbf{1} - \lambda T)$. It is useful to determine whether these four quantities are independent when they arise from DBI field configurations. The factorization of the characteristic polynomial derived in appendix B implies

$$\begin{aligned}(E_4 - 1)^2 E_2 = & -E_1^2 E_4 + E_1 E_3 (E_4 + 1) - E_3^2 \\ & + E_4^3 - E_4^2 - E_4 + 1.\end{aligned} \tag{6.1}$$

Hence the four E_k generated by DBI field configurations are not independent.

Equation (6.1) shows directly that (E_1, E_2, E_3, E_4) cannot vary independently on DBI field configurations. The existence of the flow in eq. (3.18) therefore does not follow from a simple counting argument based on the number of available invariants. The same relation also makes the algebraic representation of the flow operator non-unique: any multiple of the difference between the two sides of eq. (6.1) may be added to a stress-tensor expression without changing its value on DBI configurations.

7 Summary and discussion

We have derived a stress-tensor flow for four-dimensional DBI theory containing one real scalar and one Abelian gauge field. For $X \neq 0$, the derivative $\partial \mathcal{L}_\lambda / \partial \lambda$ is given by eq. (3.18), which depends only on local invariants of the mixed stress tensor. The same expression follows from a direct four-dimensional calculation and from dimensional reduction of the five-dimensional Born–Infeld flow. At $X = 0$, eq. (5.2) provides a finite stress-tensor expression, and the pure Born–Infeld limit reduces to the known trace flow.

The immediate significance of the result is structural. Stress-tensor flows provide a way of describing trajectories in the space of classical Lagrangians without referring explicitly to all microscopic field invariants. In four-dimensional NLED this viewpoint has been useful for organizing families of duality-invariant theories and their deformations [10, 9]. The present calculation shows that such a description can persist when a gauge field and a worldvolume scalar are present simultaneously, at least in the special four-dimensional single-scalar DBI theory. In this sense it provides a four-dimensional single-scalar realization of the gauge-plus-scalar problem highlighted in ref. [11]. Since the undeformed limit is Maxwell theory plus a free massless scalar, one natural question is whether eq. (3.18), together with the seed Lagrangian in eq. (2.6) and the $X = 0$ prescription, uniquely reconstructs the DBI family within an appropriate class of local

derivative-free Lagrangians. Establishing such a uniqueness statement would give an alternative characterization of this DBI theory directly in terms of its stress tensor.

The dimensional-reduction derivation provides a second direction. Brizio *et al.* found stress-tensor flows for Born–Infeld theory in general dimension and for DBI theories in two and three dimensions [13]. The present five-to-four-dimensional reduction gives another explicit case in which a higher-dimensional Born–Infeld flow descends to a DBI flow, but only after the explicit field-strength dependence can be eliminated. This suggests studying dimensional reduction as a way to classify when a DBI deformation can be written solely in terms of stress-tensor invariants. The algebraic constraint in eq. (6.1) is relevant to such a classification: stress-tensor expressions that differ by a function vanishing on DBI configurations represent the same flow on this theory, so these algebraic relations must be taken into account when comparing candidate flow operators.

Several extensions are then concrete. With multiple worldvolume scalars, the rank-one structure used here is lost, while in other spacetime dimensions the algebra of a two-form is different; both changes can obstruct the elimination of explicit $F_{\mu\nu}$ invariants. It would therefore be useful to determine systematically which dimensions and field contents admit a stress-tensor-only description. Another question is whether the bosonic relation found here admits a supersymmetric completion or a formulation compatible with electromagnetic duality in the gauge sector, paralleling developments for four-dimensional NLED stress-tensor flows [9, 10]. Finally, the numerical search used in this work can serve as a diagnostic in these more complicated theories: it can test whether a candidate deformation is approximately single-valued as a function of stress-tensor invariants before one attempts an analytic elimination. The four-dimensional single-scalar theory provides a controlled test case in which this numerical strategy is known to lead to an exact relation.

All statements in this paper concern classical Lagrangian flows. Whether the relation admits a useful quantum interpretation in four dimensions is a separate question.

Acknowledgments

Use of generative AI. OpenAI ChatGPT was used in this project to assist with exploration of research directions, preparation of numerical and symbolic-computation code, literature searches, and organization and editing of the manuscript. The analytic and numerical calculations and the references supporting the scientific claims were checked by the author, who takes responsibility for the content of the paper.

Data availability

No external experimental or observational data were used in this work. Numerical validation data generated for the study are maintained in the accompanying repository.

Code availability

Code for the numerical search, checks of the analytic formulas, and the five-to-four-dimensional reduction, together with numerical outputs, is maintained in the accompanying repository:

`https://github.com/ryos-physrockme/dbi-stress-tensor-flow`.

Code under the repository's `code/` directory is distributed under the MIT License.

A Neural-network-assisted numerical search

Before the analytic derivation, we investigated numerically whether Q_λ could be predicted from stress-tensor invariants alone. On the flat metric $\eta = \mathrm{diag}(-1, 1, 1, 1)$, define dimensionless fields

$$a_\mu = \sqrt{\lambda}\,\partial_\mu \phi, \qquad b_{\mu\nu} = \sqrt{\lambda}\,F_{\mu\nu}. \tag{A.1}$$

For a timelike scalar gradient, a Lorentz transformation brings a_μ to $(a, 0, 0, 0)$, and the remaining spatial rotations reduce the independent nonzero components of $b_{\mu\nu}$ to three. We used

$$b_{01} = e, \qquad b_{23} = p, \qquad b_{31} = b \tag{A.2}$$

and drew samples uniformly from

$$a \in [0.18, 0.72], \quad e \in [-0.42, 0.42], \quad p, b \in [-0.52, 0.52]. \tag{A.3}$$

We retained only configurations on the real DBI branch and excluded points with $\Delta < 0.30$, $|1 - E_4| < 5 \times 10^{-3}$, or $|\lambda^2 Q_\lambda| > 2$ in order to stabilize the initial regression.

For each field configuration, the DBI Lagrangian and stress tensor were evaluated directly and used to form

$$\boldsymbol{x} = (E_1, E_2, E_3, E_4) \in \mathbb{R}^4, \qquad y = \lambda^2 Q_\lambda \in \mathbb{R}. \tag{A.4}$$

The neural network approximated the map

$$f_\theta : \mathbb{R}^4 \longrightarrow \mathbb{R}, \qquad (E_1, E_2, E_3, E_4) \longmapsto f_\theta(E_1, E_2, E_3, E_4) \simeq \lambda^2 Q_\lambda. \tag{A.5}$$

Inputs and targets were standardized using the mean and standard deviation of the training set. If $\widetilde{\boldsymbol{x}}_n$ and $\widetilde{y}_n$ denote the standardized input and target of sample n, the loss was the mean-squared error

$$\mathcal{J}(\theta) = \frac{1}{N_{\mathrm{train}}} \sum_{n \in \mathcal{D}_{\mathrm{train}}} \left[f_\theta(\widetilde{\boldsymbol{x}}_n) - \widetilde{y}_n \right]^2 . \tag{A.6}$$

No known flow equation or analytic candidate was included in the loss.

The network architecture was 4–64–64–64–1, with tanh activations in the hidden layers. We generated 18,000 field configurations, using 80% for training and 20% for testing. The network was trained with Adam, learning rate 2×10^{-3}, batch size 512, and 260 epochs. On the test set, the root-mean-square error was 7.31×10^{-4} and the relative L^2 error was 1.37×10^{-2}.

To extract an analytic candidate, we generated an independent set of 12,000 canonical-frame configurations and considered the simplest rational ansatz whose numerator and denominator are affine functions of the E_k,

$$y = \frac{a_0 + a_1 E_1 + a_2 E_2 + a_3 E_3 + a_4 E_4}{b_0 + b_1 E_1 + b_2 E_2 + b_3 E_3 + b_4 E_4}. \tag{A.7}$$

After clearing the denominator, the coefficients a_i, b_i obey a homogeneous linear system. The singular vector associated with the smallest singular value reproduced, up to an overall normalization, the coefficients in eq. (3.16).

After obtaining this candidate, we tested eq. (3.16) on 30,000 field configurations generated without using the canonical frame, with all four components of a_μ and all six independent components of $b_{\mu\nu}$ sampled independently. The final claims of the paper are based on the analytic derivation in the main text and appendix B, rather than on these numerical tests. The code and numerical outputs are available in the repository listed in the code-availability statement.

B Auxiliary matrix identities

B.1 Determinant identities in four dimensions

For compactness, define within this appendix

$$C_{\pm}{}^{\mu}{}_{\nu} = \delta^{\mu}{}_{\nu} \pm \sqrt{\lambda} F^{\mu}{}_{\nu} + \lambda\, \partial^{\mu}\phi\, \partial_{\nu}\phi. \tag{B.1}$$

Equation (2.3) gives $\det C_+ = \det C_- = \Delta^2$. The stress tensor in eq. (2.8) may be written as

$$M = \frac{\Delta}{2}(C_+^{-1} + C_-^{-1}) = \Delta C_-^{-1}\left(\mathbf{1} + \lambda\, \partial\phi \otimes \partial\phi\right) C_+^{-1}. \tag{B.2}$$

Using the matrix determinant lemma

$$\det(A + uv^{\mathsf{T}}) = \det A \left(1 + v^{\mathsf{T}} A^{-1} u\right), \tag{B.3}$$

one immediately finds

$$\det M = 1 + \lambda X. \tag{B.4}$$

To obtain E_1, consider

$$D(a) = \det\Big(a\mathbf{1} + \sqrt{\lambda} F + \lambda\, \partial\phi \otimes \partial\phi\Big) \tag{B.5}$$

for real a. The four-dimensional antisymmetric matrix $F^{\mu}{}_{\nu}$ obeys

$$(\sqrt{\lambda}F)^4 - 2\lambda S(\sqrt{\lambda}F)^2 - \lambda^2 P^2 \mathbf{1} = 0. \tag{B.6}$$

Combining this with the matrix determinant lemma gives

$$D(a) = a^4 + \lambda X a^3 - 2\lambda S a^2 + \lambda^2(Z - 2SX)a - \lambda^2 P^2. \tag{B.7}$$

The Jacobi formula,

$$\frac{\mathrm{d}}{\mathrm{d}a} \det B(a) = \det B(a)\, \operatorname{tr}\left(B(a)^{-1} \frac{\mathrm{d}B(a)}{\mathrm{d}a}\right), \tag{B.8}$$

implies $D'(1) = \det C_+ \operatorname{tr} C_+^{-1}$ and hence eq. (3.11). To obtain E_3, we use the rank-one inverse formula

$$(A + uv^{\mathsf{T}})^{-1} = A^{-1} - \frac{A^{-1} u v^{\mathsf{T}} A^{-1}}{1 + v^{\mathsf{T}} A^{-1} u}. \tag{B.9}$$

Substituting eq. (B.2) into $E_3 = \det M \operatorname{tr} M^{-1}$ then yields eq. (3.12).

B.2 Factorization of the characteristic polynomial

We now derive the factorization used in sections 5 and 6. In this subsection only, write

$$v_\mu \equiv \partial_\mu \phi \tag{B.10}$$

and define

$$h_{\mu\nu} = g_{\mu\nu} + \lambda v_\mu v_\nu, \qquad r \equiv \frac{\det h}{\det g} = 1 + \lambda X. \tag{B.11}$$

We first assume $r \neq 0$ so that $h_{\mu\nu}$ is invertible. The final characteristic-polynomial relation is polynomial in r, and therefore extends continuously to $r = 0$ wherever the DBI Lagrangian itself remains defined. Introduce

$$J^{\mu}{}_{\nu} = \sqrt{\lambda}\, h^{\mu\rho} F_{\rho\nu}. \tag{B.12}$$

Then J is antisymmetric with respect to $h_{\mu\nu}$ and $\mathcal{G} = h(\mathbf{1} + J)$, so

$$\Delta^2 = r \det(\mathbf{1} + J). \tag{B.13}$$

The h-antisymmetry of J implies

$$(\mathcal{G}^{-1})^{(\mu\rho)} h_{\rho\nu} = \frac{1}{2}\left[(\mathbf{1}+J)^{-1} + (\mathbf{1}-J)^{-1}\right]^{\mu}{}_{\nu} = (\mathbf{1}-J^2)^{-1\,\mu}{}_{\nu}. \tag{B.14}$$

The rank-one inverse formula also gives

$$(h^{-1}g)^{\mu}{}_{\nu} = \delta^{\mu}{}_{\nu} - \frac{\lambda}{r} v^{\mu} v_{\nu}. \tag{B.15}$$

Substituting these relations into eq. (2.8) yields

$$M = U\left(\mathbf{1} - \frac{R}{r}\right), \qquad U \equiv \Delta(\mathbf{1}-J^2)^{-1}, \qquad R^{\mu}{}_{\nu} \equiv \lambda v^{\mu} v_{\nu}. \tag{B.16}$$

Here $\operatorname{tr} R = \lambda X = r - 1$.

The eigenvalues of a four-dimensional antisymmetric matrix J occur in opposite-sign pairs, so J^2 has two eigenvalues, each with multiplicity two. The same is therefore true of U. Moreover, $\det(\mathbf{1} - J) = \det(\mathbf{1} + J)$ together with eq. (B.13) gives

$$\det U = r^2. \tag{B.17}$$

On the branch connected continuously to $\lambda = 0$, the product of the two distinct eigenvalues of U is r. Hence there exists a coefficient α such that

$$U^2 - \alpha U + r\mathbf{1} = 0. \tag{B.18}$$

Defining $q(z) = z^2 - \alpha z + r$, we have

$$\det(z\mathbf{1} - U) = q(z)^2, \qquad (z\mathbf{1} - U)^{-1} U = \frac{zU - r\mathbf{1}}{q(z)}. \tag{B.19}$$

Equation (B.16) is a rank-one update of U. Applying the matrix determinant lemma gives

$$\begin{aligned} \det(z\mathbf{1} - M) &= q(z)^2 \left[1 + \frac{1}{r}\operatorname{tr}\big(R(z\mathbf{1}-U)^{-1}U\big)\right] \\ &= q(z)\left[q(z) + \frac{\operatorname{tr}(RU)}{r} z - \operatorname{tr} R\right]. \end{aligned} \tag{B.20}$$

Using $\operatorname{tr} R = r - 1$ and defining

$$\beta \equiv \alpha - \frac{\operatorname{tr}(RU)}{r}, \tag{B.21}$$

we obtain

$$\det(z\mathbf{1} - M) = (z^2 - \alpha z + r)(z^2 - \beta z + 1). \tag{B.22}$$

Since $r = E_4$ by eq. (3.8), comparison of coefficients yields

$$E_1 = \alpha + \beta, \qquad E_2 = 1 + E_4 + \alpha\beta, \qquad E_3 = \alpha + E_4\beta. \tag{B.23}$$

For $E_4 \neq 1$, eq. (3.15) implies

$$\beta = \frac{E_1 - E_3}{1 - E_4} = 2(1 + \lambda^2 Q_\lambda). \tag{B.24}$$

Taking the continuous DBI branch as $E_4 \to 1$, eq. (B.23) gives

$$\beta = \frac{1}{2}\left[E_1 + \sqrt{E_1^2 - 4E_2 + 8}\right], \tag{B.25}$$

which yields eq. (5.1). The discriminant is $(\alpha - \beta)^2$ on this factorized characteristic polynomial and is therefore non-negative on the real DBI branch. Eliminating α and β from eq. (B.23) gives eq. (6.1).

References

[1] M. Born and L. Infeld, "Foundations of the New Field Theory," Proc. Roy. Soc. Lond. A **144** (1934) 425–451.

[2] F. A. Smirnov and A. B. Zamolodchikov, "On space of integrable quantum field theories," Nucl. Phys. B **915** (2017) 363–383, arXiv:1608.05499 [hep-th].

[3] M. Taylor, "$T\overline{T}$ deformations in general dimensions," Adv. Theor. Math. Phys. **24** (2020) 37–63, arXiv:1805.10287 [hep-th].

[4] I. Bandos, K. Lechner, D. Sorokin and P. K. Townsend, "A non-linear duality-invariant conformal extension of Maxwell's equations," Phys. Rev. D **102** (2020) 121703, arXiv:2007.09092 [hep-th].

[5] D. P. Sorokin, "Introductory Notes on Non-linear Electrodynamics and its Applications," Fortsch. Phys. **70** (2022) 2200092, arXiv:2112.12118 [hep-th].

[6] H. Babaei-Aghbolagh, K. Babaei Velni, D. M. Yekta and H. Mohammadzadeh, "Emergence of non-linear electrodynamic theories from $T\overline{T}$-like deformations," Phys. Lett. B **829** (2022) 137079, arXiv:2202.11156 [hep-th].

[7] C. Ferko, L. Smith and G. Tartaglino-Mazzucchelli, "On Current-Squared Flows and ModMax Theories," SciPost Phys. **13** (2022) 012, arXiv:2203.01085 [hep-th].

[8] R. Conti, J. Romano and R. Tateo, "Metric approach to a $T\overline{T}$-like deformation in arbitrary dimensions," JHEP **09** (2022) 085, arXiv:2206.03415 [hep-th].

[9] C. Ferko, L. Smith and G. Tartaglino-Mazzucchelli, "Stress Tensor Flows, Birefringence in Non-Linear Electrodynamics, and Supersymmetry," SciPost Phys. **15** (2023) 198, arXiv:2301.10411 [hep-th].

[10] C. Ferko, S. M. Kuzenko, L. Smith and G. Tartaglino-Mazzucchelli, "Duality-Invariant Non-linear Electrodynamics and Stress Tensor Flows," Phys. Rev. D **108** (2023) 106021, arXiv:2309.04253 [hep-th].

[11] C. Ferko, Y. Hu, Z. Huang, K. Koutrolikos and G. Tartaglino-Mazzucchelli, "$T\overline{T}$-like flows and $3d$ nonlinear supersymmetry," SciPost Phys. **16** (2024) 038, arXiv:2302.10410 [hep-th].

[12] C. D. A. Blair, J. Lahnsteiner, N. A. Obers and Z. Yan, "Matrix Theory Reloaded: A BPS Road to Holography," JHEP **02** (2025) 024, arXiv:2410.03591 [hep-th].

[13] N. Brizio, M. Kade, A. Sfondrini and D. P. Sorokin, "More on $T\overline{T}$-like deformations in higher dimensions," JHEP **06** (2026) 262, arXiv:2602.24058 [hep-th].