


# Lecture 1: Mathematical Preliminaries

Shi Pu

School of Data Science (SDS)  
The Chinese University of Hong Kong, Shenzhen



- 1 Spaces, inner products and norms
- 2 Eigenvalues and eigenvectors
- 3 Basic topological concepts
- 4 Differentiability

1 Spaces, inner products and norms

2 Eigenvalues and eigenvectors

3 Basic topological concepts

4 Differentiability

# The Space $\mathbb{R}^n$

- $\mathbb{R}^n$  - the set of  $n$ -dimensional column vectors with real components endowed with the component-wise addition operator:

$$\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} + \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} = \begin{pmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_n \end{pmatrix},$$

and the scalar-vector product

$$\lambda \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} \lambda x_1 \\ \lambda x_2 \\ \vdots \\ \lambda x_n \end{pmatrix}.$$

- $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$  - standard/canonical basis.
- $\mathbf{e}$  and  $\mathbf{0}$  - all ones and all zeros column vectors.

- Nonnegative orthant:

$$\mathbb{R}_+^n = \{(x_1, x_2, \dots, x_n)^\top : x_1, x_2, \dots, x_n \geq 0\}.$$

- Positive orthant:

$$\mathbb{R}_{++}^n = \{(x_1, x_2, \dots, x_n)^\top : x_1, x_2, \dots, x_n > 0\}.$$

- If  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ , the closed line segment between  $\mathbf{x}$  and  $\mathbf{y}$  is given by

$$[\mathbf{x}, \mathbf{y}] = \{\mathbf{x} + \alpha(\mathbf{y} - \mathbf{x}) : \alpha \in [0, 1]\}.$$

- The open line segment  $(\mathbf{x}, \mathbf{y})$  is similarly defined as

$$(\mathbf{x}, \mathbf{y}) = \{\mathbf{x} + \alpha(\mathbf{y} - \mathbf{x}) : \alpha \in (0, 1)\}$$

for  $\mathbf{x} \neq \mathbf{y}$  and  $(\mathbf{x}, \mathbf{x}) = \emptyset$

- Unit-simplex:

$$\Delta_n = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{x} \geq 0, \mathbf{e}^\top \mathbf{x} = 1\}.$$

- The set of all real valued matrices is denoted by  $\mathbb{R}^{m \times n}$ .
- $\mathbf{I}_n$  -  $n \times n$  identity matrix.
- $\mathbf{0}_{m \times n}$  -  $m \times n$  zeros matrix.

## Definition

An inner product on  $\mathbb{R}^n$  is a map  $\langle \cdot, \cdot \rangle : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  with the following properties:

1. (symmetry)  $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle$  for any  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ .
2. (additivity)  $\langle \mathbf{x}, \mathbf{y} + \mathbf{z} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle + \langle \mathbf{x}, \mathbf{z} \rangle$  for any  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}^n$ .
3. (homogeneity)  $\langle \lambda \mathbf{x}, \mathbf{y} \rangle = \lambda \langle \mathbf{x}, \mathbf{y} \rangle$  for any  $\lambda \in \mathbb{R}$  and  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ .
4. (positive definiteness)  $\langle \mathbf{x}, \mathbf{x} \rangle \geq 0$  for any  $\mathbf{x} \in \mathbb{R}^n$  and  $\langle \mathbf{x}, \mathbf{x} \rangle = 0$  if and only if  $\mathbf{x} = \mathbf{0}$ .

## Definition

An inner product on  $\mathbb{R}^n$  is a map  $\langle \cdot, \cdot \rangle : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  with the following properties:

1. (symmetry)  $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle$  for any  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ .
2. (additivity)  $\langle \mathbf{x}, \mathbf{y} + \mathbf{z} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle + \langle \mathbf{x}, \mathbf{z} \rangle$  for any  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}^n$ .
3. (homogeneity)  $\langle \lambda \mathbf{x}, \mathbf{y} \rangle = \lambda \langle \mathbf{x}, \mathbf{y} \rangle$  for any  $\lambda \in \mathbb{R}$  and  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ .
4. (positive definiteness)  $\langle \mathbf{x}, \mathbf{x} \rangle \geq 0$  for any  $\mathbf{x} \in \mathbb{R}^n$  and  $\langle \mathbf{x}, \mathbf{x} \rangle = 0$  if and only if  $\mathbf{x} = \mathbf{0}$ .

- The “dot product”

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^\top \mathbf{y} = \sum_{i=1}^n x_i y_i \text{ for any } \mathbf{x}, \mathbf{y} \in \mathbb{R}^n.$$

- The “weighted dot product”

$$\langle \mathbf{x}, \mathbf{y} \rangle_{\mathbf{w}} = \sum_{i=1}^n w_i x_i y_i, \text{ where } \mathbf{w} \in \mathbb{R}_{++}^n.$$



## Definition

An norm  $\|\cdot\|$  on  $\mathbb{R}^n$  is a function  $\|\cdot\| : \mathbb{R}^n \rightarrow \mathbb{R}$  satisfying

1. (nonnegativity)  $\|\mathbf{x}\| \geq 0$  for any  $\mathbf{x} \in \mathbb{R}^n$  and  $\|\mathbf{x}\| = 0$  if and only if  $\mathbf{x} = \mathbf{0}$ .
2. (positive homogeneity)  $\|\lambda\mathbf{x}\| = |\lambda|\|\mathbf{x}\|$  for any  $\mathbf{x} \in \mathbb{R}^n$  and  $\lambda \in \mathbb{R}$ .
3. (triangle inequality)  $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$  for any  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ .

- One natural way to generate a norm on  $\mathbb{R}^n$  is to take any inner product  $\langle \cdot, \cdot \rangle$  defined on  $\mathbb{R}^n$ , and define the associated norm

$$\|\mathbf{x}\| \equiv \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle}, \text{ for all } \mathbf{x} \in \mathbb{R}^n,$$

- The norm associated with the dot-product is the so-called Euclidean norm or  $l_2$ -norm:

$$\|\mathbf{x}\|_2 \equiv \sqrt{\sum_{i=1}^n x_i^2} \text{ for all } \mathbf{x} \in \mathbb{R}^n.$$

- The  $l_p$ -norm ( $p \geq 1$ ) is defined by  $\|x\|_p \equiv \sqrt[p]{\sum_{i=1}^n |x_i|^p}$ .
- The  $l_\infty$ -norm is

$$\|\mathbf{x}\|_\infty \equiv \max_{i=1,2,\dots,n} |x_i|.$$

- It can be shown that

$$\|\mathbf{x}\|_\infty = \lim_{p \rightarrow \infty} \|\mathbf{x}\|_p.$$

Example:  $l_{1/2}$  is not a norm. why?

# The Cauchy-Schwartz Inequality

## Lemma

For any  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ :

$$|\mathbf{x}^\top \mathbf{y}| \leq \|\mathbf{x}\| \cdot \|\mathbf{y}\|.$$

## Proof.

For any  $\lambda \in \mathbb{R}$ :

$$\|\mathbf{x} + \lambda \mathbf{y}\|^2 = \|\mathbf{x}\|^2 + 2\lambda \langle \mathbf{x}, \mathbf{y} \rangle + \lambda^2 \|\mathbf{y}\|^2$$

Therefore (why?),

$$4\langle \mathbf{x}, \mathbf{y} \rangle^2 - 4\|\mathbf{x}\|^2 \|\mathbf{y}\|^2 \leq 0,$$

establishing the desired result. □

## Definition

A norm  $\|\cdot\|$  on  $\mathbb{R}^{m \times n}$  is a function  $\|\cdot\| : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$  satisfying

1. (nonnegativity)  $\|\mathbf{A}\| \geq 0$  for any  $\mathbf{A} \in \mathbb{R}^{m \times n}$  and  $\|\mathbf{A}\| = 0$  if and only if  $\mathbf{A} = \mathbf{0}$ .
2. (positive homogeneity)  $\|\lambda \mathbf{A}\| = |\lambda| \|\mathbf{A}\|$  for any  $\mathbf{A} \in \mathbb{R}^{m \times n}$  and  $\lambda \in \mathbb{R}$ .
3. (triangle inequality)  $\|\mathbf{A} + \mathbf{B}\| \leq \|\mathbf{A}\| + \|\mathbf{B}\|$  for any  $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{m \times n}$ .
4. (submultiplicativity)  $\|\mathbf{AB}\| \leq \|\mathbf{A}\| \|\mathbf{B}\|$  for any compatible  $\mathbf{A}, \mathbf{B}$ .

- Given a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  and two norms  $\|\cdot\|_a$  and  $\|\cdot\|_b$  on  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, the induced matrix norm  $\|\mathbf{A}\|_{a,b}$  (called  $(a, b)$ -norm) is defined by

$$\|\mathbf{A}\|_{a,b} = \max_{\mathbf{x}} \{\|\mathbf{Ax}\|_b : \|\mathbf{x}\|_a \leq 1\}.$$

Conclusion:

$$\|\mathbf{Ax}\|_b \leq \|\mathbf{A}\|_{a,b} \|\mathbf{x}\|_a$$

- An induced norm is a norm (satisfies nonnegativity, positive homogeneity and triangle inequality, and submultiplicativity).
- We refer to the matrix-norm  $\|\cdot\|_{a,b}$  as the  $(a, b)$ -norm. When  $a = b$ , we will simply refer to it as an  $a$ -norm.

- Spectral norm: If  $\|\cdot\|_a = \|\cdot\|_b = \|\cdot\|_2$ , the induced  $(2,2)$ -norm of a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  is the maximum singular value of  $\mathbf{A}$

$$\|\mathbf{A}\|_2 = \|\mathbf{A}\|_{2,2} = \sqrt{\lambda_{\max}(\mathbf{A}^\top \mathbf{A})} \equiv \sigma_{\max}(\mathbf{A}).$$

- $l_1$ -norm: when  $\|\cdot\|_a = \|\cdot\|_b = \|\cdot\|_1$ , the induced  $(1,1)$ -matrix norm of a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  is given by

$$\|\mathbf{A}\|_1 = \max_{j=1,2,\dots,n} \sum_{i=1}^m |A_{i,j}|.$$

- $l_\infty$ -norm: when  $\|\cdot\|_a = \|\cdot\|_b = \|\cdot\|_\infty$ , the induced  $(\infty, \infty)$ -matrix norm of a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  is given by

$$\|\mathbf{A}\|_\infty = \max_{i=1,2,\dots,m} \sum_{j=1}^n |A_{i,j}|.$$

# The Frobenius norm

$$\|\mathbf{A}\|_F = \sqrt{\sum_{i=1}^m \sum_{j=1}^n A_{ij}^2}, \mathbf{A} \in \mathbb{R}^{m \times n}$$

The Frobenius norm is not an induced norm.  
Why is it a norm?

- 1 Spaces, inner products and norms
- 2 Eigenvalues and eigenvectors
- 3 Basic topological concepts
- 4 Differentiability



- Let  $\mathbf{A} \in \mathbb{R}^{n \times n}$ . Then a nonzero vector  $\mathbf{v} \in \mathbb{R}^n$  is called an eigenvector of  $\mathbf{A}$  if there exists a  $\lambda \in \mathbb{C}$  for which

$$\mathbf{A}\mathbf{v} = \lambda\mathbf{v}.$$

The scalar  $\lambda$  is the eigenvalue corresponding to the eigenvector  $\mathbf{v}$ .

- In general, real-valued matrices can have complex eigenvalues, but when the matrix is *symmetric* the eigenvalues are necessarily *real*.
- The eigenvalues of a symmetric  $n \times n$  matrix  $\mathbf{A}$  are denoted by

$$\lambda_1(\mathbf{A}) \geq \lambda_2(\mathbf{A}) \geq \dots \geq \lambda_n(\mathbf{A}).$$

- The maximum eigenvalue is also denoted by  $\lambda_{\max}(\mathbf{A})(= \lambda_1(\mathbf{A}))$ , and the minimum eigenvalue is also denoted by  $\lambda_{\min}(\mathbf{A})(= \lambda_n(\mathbf{A}))$ .

## Theorem

Let  $\mathbf{A} \in \mathbb{R}^{n \times n}$  be an  $n \times n$  symmetric matrix. Then there exists an orthogonal matrix  $\mathbf{U} \in \mathbb{R}^{n \times n}$  ( $\mathbf{U}^\top \mathbf{U} = \mathbf{U} \mathbf{U}^\top = \mathbf{I}$ ) and a diagonal matrix  $\mathbf{D} = \text{diag}(d_1, d_2, \dots, d_n)$  for which

$$\mathbf{U}^\top \mathbf{A} \mathbf{U} = \mathbf{D}.$$

- The columns of the matrix  $\mathbf{U}$  constitute an orthogonal basis comprising eigenvectors of  $\mathbf{A}$  and the diagonal elements of  $\mathbf{D}$  are the corresponding eigenvalues.
- A direct result is that  $\text{Tr}(\mathbf{A}) = \sum_{i=1}^n \lambda_i(\mathbf{A})$  and  $\det(\mathbf{A}) = \prod_{i=1}^n \lambda_i(\mathbf{A})$ .

- 1 Spaces, inner products and norms
- 2 Eigenvalues and eigenvectors
- 3 Basic topological concepts**
- 4 Differentiability

# Basic Topological Concepts

- The open ball with center  $\mathbf{c} \in \mathbb{R}^n$  and radius  $r$ :

$$B(\mathbf{c}, r) = \{\mathbf{x} : \|\mathbf{x} - \mathbf{c}\| < r\}.$$

- The closed ball with center  $\mathbf{c} \in \mathbb{R}^n$  and radius  $r$ :

$$B[\mathbf{c}, r] = \{\mathbf{x} : \|\mathbf{x} - \mathbf{c}\| \leq r\}.$$

## Definition

Given a set  $U \subseteq \mathbb{R}^n$ , a point  $\mathbf{c} \in U$  is called an interior point of  $U$  if there exists  $r > 0$  for which  $B(\mathbf{c}, r) \subseteq U$ .

The set of all interior points of a given set  $U$  is called the interior of the set and is denoted by  $\text{int}(U)$ :

$$\text{int}(U) = \{\mathbf{x} \in U : B(\mathbf{x}, r) \subseteq U \text{ for some } r > 0.\}$$

Examples:

$$\text{int}(\mathbb{R}_+^n) = \mathbb{R}_{++}^n, \quad \text{int}(B[\mathbf{c}, r]) = B(\mathbf{c}, r), \quad \text{int}([\mathbf{x}, \mathbf{y}]) = ?$$

- An open set is a set that contains only interior points, meaning that  $U = \text{int}(U)$ .
- Examples of open sets are open balls (hence the name...) and the positive orthant  $\mathbb{R}_{++}^n$ .

Result: a union of any number of open sets is an open set and the intersection of a finite number of open sets is open.

- A set  $U \subseteq \mathbb{R}^n$  is closed if it contains all the limits of convergent sequences of vectors in  $U$ , that is, if  $\{\mathbf{x}_i\}_{i=1}^{\infty} \subseteq U$  satisfies  $\mathbf{x}_i \rightarrow \mathbf{x}^*$  as  $i \rightarrow \infty$ , then  $\mathbf{x}^* \in U$ .
- A known result states that  $U$  is closed iff its complement  $U^c$  is open.
- Examples of closed sets are the closed ball  $B[\mathbf{c}, r]$ , closed line segments, the nonnegative orthant  $\mathbb{R}_+^n$  and the unit simplex  $\Delta_n$ .

What about  $\mathbb{R}^n$ ?  $\emptyset$ ?

## Definition

Given a set  $U \subseteq \mathbb{R}^n$ , a boundary point of  $U$  is a vector  $\mathbf{x} \in \mathbb{R}^n$  satisfying the following: any neighborhood of  $\mathbf{x}$  contains at least one point in  $U$  and at least one point in its complement  $U^c$ .

- The set of all boundary points of a set  $U$  is denoted by  $\text{bd}(U)$ .

Examples:

$$\text{bd}(B(\mathbf{c}, r)) =$$

$$\text{bd}(B[\mathbf{c}, r]) =$$

$$\text{bd}(\mathbb{R}_{++}^n) =$$

$$\text{bd}(\mathbb{R}_+^n) =$$

$$\text{bd}(\mathbb{R}^n) =$$

$$\text{bd}(\Delta_n) =$$

- The closure of a set  $U \subseteq \mathbb{R}^n$  is denoted by  $\text{cl}(U)$  and is defined to be the smallest closed set containing  $U$ :

$$\text{cl}(U) = \bigcap \{ T : U \subseteq T, T \text{ is closed} \}$$

- Another equivalent definition of  $\text{cl}(U)$  is:

$$\text{cl}(U) = U \cup \text{bd}(U).$$

Examples:

$$\begin{aligned}\text{cl}(\mathbb{R}_{++}^n) &= \\ \text{cl}(B(\mathbf{c}, r)) &= \\ (\mathbf{x} \neq \mathbf{y}), \text{cl}((\mathbf{x}, \mathbf{y})) &= \end{aligned}$$



- A set  $U \subseteq \mathbb{R}^n$  is called bounded if there exists  $M > 0$  for which  $U \subseteq B(\mathbf{0}, M)$ .
- A set  $U \subseteq \mathbb{R}^n$  is called compact if it is closed and bounded.
- Examples of compact sets: closed balls, unit simplex, closed line segments.

- 1 Spaces, inner products and norms
- 2 Eigenvalues and eigenvectors
- 3 Basic topological concepts
- 4 Differentiability**

## Definition

Let  $f$  be a function defined on a set  $S \subseteq \mathbb{R}^n$ . Let  $\mathbf{x} \in \text{int}(S)$  and let  $\mathbf{d} \in \mathbb{R}^n$ . If the limit

$$\lim_{t \rightarrow 0^+} \frac{f(\mathbf{x} + t\mathbf{d}) - f(\mathbf{x})}{t}$$

exists, then it is called the directional derivative of  $f$  at  $\mathbf{x}$  along the direction  $\mathbf{d}$  and is denoted by  $f'(\mathbf{x}; \mathbf{d})$ .

- For any  $i = 1, 2, \dots, n$ , if the limit

$$\lim_{t \rightarrow 0} \frac{f(\mathbf{x} + t\mathbf{e}_i) - f(\mathbf{x})}{t}$$

exists, then its value is called the  $i$ -th partial derivative and is denoted by  $\frac{\partial f}{\partial x_i}(\mathbf{x})$ .

- If all the partial derivatives of a function  $f$  exist at a point  $\mathbf{x} \in \mathbb{R}^n$ , then the gradient of  $f$  at  $\mathbf{x}$  is  $\nabla f(\mathbf{x}) = (\frac{\partial f}{\partial x_1}(\mathbf{x}), \frac{\partial f}{\partial x_2}(\mathbf{x}), \dots, \frac{\partial f}{\partial x_n}(\mathbf{x}))^\top$ .

## Continuous Differentiability

A function  $f$  defined on an open set  $U \subseteq \mathbb{R}^n$  is called continuously differentiable over  $U$  if all the partial derivatives exist and are continuous on  $U$ . In that case,

$$f'(\mathbf{x}; \mathbf{d}) = \nabla f(\mathbf{x})^\top \mathbf{d}, \quad \mathbf{x} \in U, \mathbf{d} \in \mathbb{R}^n$$

### Proposition

*Let  $f : U \rightarrow \mathbb{R}$  be defined on an open set  $U \subseteq \mathbb{R}^n$ . Suppose that  $f$  is continuously differentiable over  $U$ . Then*

$$\lim_{\mathbf{d} \rightarrow 0} \frac{f(\mathbf{x} + \mathbf{d}) - f(\mathbf{x}) - \nabla f(\mathbf{x})^\top \mathbf{d}}{\|\mathbf{d}\|} = 0, \quad \forall \mathbf{x} \in U.$$

Another way to write the above result is as follows:

$$f(\mathbf{y}) = f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + o(\|\mathbf{y} - \mathbf{x}\|),$$

where  $o(\cdot) : \mathbb{R}_+^n \rightarrow \mathbb{R}$  is a one-dimensional function satisfying  $\frac{o(t)}{t} \rightarrow 0$  as  $t \rightarrow 0^+$ .

- The partial derivatives  $\partial f$  are themselves real-valued functions that can be partially differentiated. The  $(i, j)$ -partial derivatives of  $f$  at  $\mathbf{x} \in U$  (if exists) is defined by

$$\frac{\partial^2 f}{\partial x_i \partial x_j}(\mathbf{x}) = \frac{\partial(\frac{\partial f}{\partial x_j})}{\partial x_i}(\mathbf{x}).$$

- A function  $f$  defined on an open set  $U \subseteq \mathbb{R}^n$  is called twice continuously differentiable over  $U$  if all the second order partial derivatives exist and are continuous over  $U$ . In that case, for any  $i \neq j$  and any  $\mathbf{x} \in U$ :

$$\frac{\partial^2 f}{\partial x_i \partial x_j}(\mathbf{x}) = \frac{\partial^2 f}{\partial x_j \partial x_i}(\mathbf{x}).$$

The Hessian of  $f$  at a point  $\mathbf{x} \in U$  is the  $n \times n$  matrix:

$$\nabla^2 f(\mathbf{x}) = \begin{pmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & & \vdots \\ \vdots & \vdots & & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_n^2} \end{pmatrix}$$

- For twice continuously differentiable functions, the Hessian is a symmetric matrix.

## Theorem

Let  $f : U \rightarrow \mathbb{R}$  be defined on an open set  $U \subseteq \mathbb{R}^n$ . Suppose that  $f$  is twice continuously differentiable over  $U$ . Let  $\mathbf{x} \in U$  and  $r > 0$  satisfy  $B(\mathbf{x}, r) \subseteq U$ . Then for any  $\mathbf{y} \in B(\mathbf{x}, r)$  there exists  $\xi \in [\mathbf{x}, \mathbf{y}]$  such that:

$$f(\mathbf{y}) = f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + \frac{1}{2}(\mathbf{y} - \mathbf{x})^\top \nabla^2 f(\xi)(\mathbf{y} - \mathbf{x}).$$

## Theorem

Let  $f : U \rightarrow \mathbb{R}$  be defined on an open set  $U \subseteq \mathbb{R}^n$ . Suppose that  $f$  is twice continuously differentiable over  $U$ . Let  $\mathbf{x} \in U$  and  $r > 0$  satisfy  $B(\mathbf{x}, r) \subseteq U$ . Then for any  $\mathbf{y} \in B(\mathbf{x}, r)$ :

$$f(\mathbf{y}) = f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + \frac{1}{2}(\mathbf{y} - \mathbf{x})^\top \nabla^2 f(\mathbf{x})(\mathbf{y} - \mathbf{x}) + o(\|\mathbf{y} - \mathbf{x}\|^2).$$