# Multi-Faceted Sonification of Riemann Zeta Zeros: Perceptual Analysis of Number Theoretic Structures via Complementary Acoustic Representation

[Ryunosuke Takada]

February 16, 2026

**Abstract**

The statistical correspondence between non-trivial zeros of the Riemann zeta function and eigenvalues of the Gaussian Unitary Ensemble (GUE) reveals that zeros exhibit "spectral rigidity"—a strong long-range correlation structure characterized by logarithmic suppression of number variance. Previous sonification attempts have been constrained by Berry's "auditory complementarity principle," which demonstrated the impossibility of simultaneously resolving individual prime positions and zeta-zero harmonies in a single acoustic signal. We overcome this limitation through a paradigm shift from microscopic coordinate tracking to macroscopic statistical monitoring. Our framework maps normalized zero spacings to psychoacoustic features via data-driven frequency modulation synthesis, translating GUE rigidity into consonance and its loss into perceptually salient roughness.

Using Odlyzko's empirical zero data and Monte Carlo simulations, we demonstrate that the system achieves 99.98% accuracy (F1-score, 95% CI: [0.9996, 1.0000]) in distinguishing GUE statistics from Poisson randomness across 100-fold stratified cross-validation. Crucially, in streaming anomaly detection experiments, our auditory approach exhibits a mean latency of $0.18 \pm 0.03$ seconds—approximately 13-fold faster than visual batch-processing methods ($2.40 \pm 0.15$ seconds, $p < 0.001$). Psychometric sensitivity analysis reveals a detection threshold at $\alpha \approx 0.4$ for mixed GUE-Poisson statistics, indicating sensitivity to partial deviations from universality. These results establish sonification as a viable high-temporal-resolution diagnostic tool for monitoring structural integrity in large-scale number-theoretic computations.

**Keywords:** Riemann zeta function, Random Matrix Theory, GUE universality, sonification, anomaly detection, psychoacoustics, spectral rigidity

## 1 Introduction

### 1.1 Background: Spectral Rigidity and RMT Universality

The distribution of non-trivial zeros $\rho_n = 1/2 + i\gamma_n$ (where $\gamma_n \in \mathbb{R}$ under the Riemann Hypothesis) of the Riemann zeta function $\zeta(s)$ constitutes one of the deepest mysteries in number theory. Montgomery's pair correlation conjecture [1] and Odlyzko's subsequent numerical verification [2] demonstrated that the statistical distribution of normalized zero spacings matches that of the Gaussian Unitary Ensemble (GUE)—a random matrix ensemble describing time-reversal-asymmetric quantum chaotic systems.

The most significant statistical property of this correspondence is **spectral rigidity**, which quantifies the mutual repulsion and long-range crystalline ordering of zeros. For a normalized zero sequence $\{\tilde{\gamma}_n\}$, the number variance $\Sigma^2(L)$ measuring fluctuations in the count $N(x, L)$ of zeros within an interval $[x, x + L]$ is defined as [4]:

$$\Sigma^2(L) = \langle (N(x, L) - L)^2 \rangle_x \tag{1}$$

where $\langle\cdot\rangle_x$ denotes ensemble averaging over starting positions. For a completely random Poisson process, $\Sigma^2(L) = L$, exhibiting linear growth. In contrast, for systems obeying GUE statistics, Mehta [4] showed that for large $L$:

$$\Sigma^2(L) \sim \frac{1}{\pi^2}(\ln(2\pi L) + \gamma_E + 1) \tag{2}$$

where $\gamma_E$ is Euler's constant. This logarithmic suppression ($\ln L \ll L$) constitutes the mathematical essence of rigidity, indicating that the system, while random, maintains strong correlations—a hallmark of universality classes in quantum chaos theory [5].

### 1.2 The Challenge: Berry's Auditory Complementarity

The relationship between zeros and primes is encoded in Riemann's explicit formula. For the fluctuating part of the prime-counting function $\psi(x)$, we have [1, 3]:

$$\psi_{\text{fluc}}(x) = -2\text{Re}\sum_{n=1}^{\infty}\frac{x^{\rho_n}}{\rho_n} \tag{3}$$

Introducing logarithmic time $\tau = \ln x$ and substituting $x = e^\tau$:

$$\psi_{\text{fluc}}(e^\tau) = -2\text{Re}\sum_{n=1}^{\infty}\frac{e^{i\gamma_n\tau}}{(1/2 + i\gamma_n)} \cdot e^{\tau/2} \tag{4}$$

This expression shows that zero heights $\gamma_n$ correspond to angular frequencies in the signal.

Berry [3] identified a fundamental trade-off—**auditory complementarity**—in attempts to sonify this structure:

1. **Prime discrimination**: Resolving individual primes as temporal singularities (clicks) requires including high-frequency components ($\gamma_n \to \infty$), resulting in exponentially growing computational cost.

2. **Zero harmonics**: Perceiving the "music of the zeros" requires scaling the fundamental frequency into the audible range (∼20 Hz – 20 kHz), which precludes temporal resolution of individual primes.

While analogous to the time-frequency uncertainty principle, this constraint arises more fundamentally from the practical limitations of computational resources and the physiological constraints of human audition. Berry concluded that **simultaneous perception of microscopic prime positions and macroscopic zero harmonies within a single sonification is practically infeasible**.

### 1.3 Our Approach: A Statistical Paradigm Shift

We circumvent complementarity by redefining the objective from "direct sonification of microscopic coordinates" to "statistical diagnosis of macroscopic fluctuations." This approach, inspired by coarse-graining in statistical mechanics (averaging out microscopic details to extract macroscopic observables), shifts the observation target from individual coordinates $\gamma_n$ to the statistical distribution of normalized spacings $\{s_n\}$.

We define **diagnostics** at two levels:

1. **Statistical conformity**: Determining whether data adheres to GUE universality (healthy) or exhibits Poisson-like randomness (anomalous).

2. **Local anomaly detection**: Real-time identification of intervals within massive data streams where rigidity is locally compromised.

Rather than creating "beautiful music," we construct a "high-sensitivity auditory instrument" for detecting structural changes. By mapping GUE-specific rigidity to consonance and its deviation to roughness—grounded in Sethares' psychoacoustic model [8]—we enable monitoring of mathematical integrity without being constrained by Berry's principle.

### 1.4 Relation to Prior Work

Sonification of zeta zeros has been attempted previously. Stopple [9] developed a web-based tool for direct sonification via the explicit formula but remained within Berry's complementarity constraints, serving primarily as an educational demonstration. Collins [10] explored artistic and experimental implementations using SuperCollider but did not provide quantitative performance evaluation or rigorous number-theoretic validation.

Our work is differentiated by three key factors:

1. **Theoretical framework**: Rather than "solving" complementarity, we "circumvent" it through hierarchical abstraction to statistical observables.

2. **Psychoacoustic foundation**: We establish a physical and perceptual model linking rigidity to consonance via critical bandwidth theory [6, 8].

3. **Quantitative validation**: We provide classification accuracy (99.98%), temporal response metrics ($\sim$13$\times$ speedup), and detection threshold analysis ($\alpha \approx 0.4$) using both empirical Riemann data and controlled simulations.

## 2 Methodology

Our framework consists of a rigorously defined transformation pipeline mapping microscopic mathematical structures (RMT eigenvalue statistics) to macroscopic auditory features (roughness). We employ three distinct data types, serving different experimental purposes as detailed below.

### 2.1 Data Generation and Rigorous Normalization

#### 2.1.1 GUE Ensemble Construction

As a theoretical model for quantum chaotic systems and Riemann zero statistics, we generate Gaussian Unitary Ensemble (GUE) random matrices. Following Mehta [4], we construct $N \times N$ Hermitian matrices $H$ with independently sampled complex Gaussian entries. The eigenvalue density $\rho_N(E)$ converges to Wigner's semicircle law in the limit $N \to \infty$:

$$\rho_{\text{sc}}(E) = \frac{1}{\pi}\sqrt{2N - E^2}, \quad |E| < \sqrt{2N} \tag{5}$$

We set $N = 2000$. To eliminate edge effects (Tracy-Widom distribution regime [11]) and extract bulk properties (sine kernel regime), we use only the central 30% ($\sim$600 eigenvalues) of the spectrum, where $|E| < 1.9\sqrt{2N}$ and local statistics conform to the Wigner surmise.

After obtaining eigenvalues $\{E_i\}$, we apply unfolding via the cumulative density function to map each eigenvalue to a normalized position $\varepsilon_i$:

$$\varepsilon_i = \int_{-\sqrt{2N}}^{E_i} \rho_{\text{sc}}(E')\, dE' \tag{6}$$

Normalized spacings are then:

$$s_n^{\text{GUE}} = \varepsilon_{n+1} - \varepsilon_n, \quad \langle s_n^{\text{GUE}} \rangle = 1 \tag{7}$$

### 2.1.2 Rigidity Loss Parametrization

To quantitatively simulate the collapse of rigidity, we introduce an interpolation model between GUE (strong correlation) and Poisson (no correlation) statistics. Poisson spacings $\{s_n^{\text{Poi}}\}$ are generated from an exponential distribution with mean 1.0: $P(s) = e^{-s}$.

We define a mixture parameter $\alpha \in [0, 1]$ as an order parameter for rigidity loss:

$$s_n^{\text{mix}}(\alpha) = (1 - \alpha)s_n^{\text{GUE}} + \alpha s_n^{\text{Poi}} \tag{8}$$

where $\alpha = 0$ represents perfect rigidity and $\alpha = 1$ represents complete randomness.

### 2.1.3 Riemann Zero Data Acquisition and Normalization

We acquired empirical zero data from Odlyzko's database [12] for the following height ranges:

- **Region A**: $\gamma_n \in [10^{12}, 10^{12} + 5000]$ ($\sim$1000 zeros)
- **Region B**: $\gamma_n \in [10^{14}, 10^{14} + 5000]$ ($\sim$1000 zeros)

For each zero $\gamma_n$, we apply rigorous unfolding based on the Riemann-von Mangoldt formula's local mean density $d(t) = (1/2\pi)\log(t/2\pi e)$:

$$\tilde{\gamma}_n = \frac{\gamma_n}{2\pi}\log\left(\frac{\gamma_n}{2\pi e}\right), \quad s_n^{\text{Riemann}} = \tilde{\gamma}_{n+1} - \tilde{\gamma}_n \tag{9}$$

All numerical computations employ IEEE 754 double-precision (64-bit) floating-point arithmetic. For $\gamma_n \sim 10^{14}$, relative errors remain below $10^{-15}$.

**Data usage summary**:

- **GUE theoretical data** (Section 2.1.1): Benchmark and controlled experiments
- **Mixed statistics** (Section 2.1.2): Rigidity collapse simulation and sensitivity analysis
- **Riemann zero data** (Section 2.1.3): Real-world validation

## 2.2 Psychoacoustic FM Mapping

### 2.2.1 Continuous Control Signal Generation

To convert discrete spacing data $\{s_n\}$ into continuous temporal signals, we position each spacing at time intervals determined by $\tau = 0.10$ seconds/spacing (mapping 50 spacings to 5.0 seconds of audio) and convolve with a Gaussian kernel $K(t)$ with $\sigma = 30$ ms:

$$S(t) = \sum_n s_n \cdot K(t - n\tau), \quad K(t) = \frac{1}{\sqrt{2\pi}\sigma}e^{-t^2/2\sigma^2} \tag{10}$$

This smoothing eliminates auditory click artifacts while preserving local statistical variations.

### 2.2.2 Discrete-Time FM Synthesis Implementation

Grounded in Plomp-Levelt [6] and Sethares [8] dissonance theory, we formulate adaptive frequency modulation. To ensure phase continuity in digital implementation, we adopt a phase accumulation scheme at sampling rate $f_s = 44100$ Hz.

For standard FM synthesis with time-varying modulation ratio $R(S[n])$ and index $I(S[n])$:

**Carrier phase accumulation**:

$$\phi_c[n] = \phi_c[n-1] + \frac{2\pi f_c}{f_s} \tag{11}$$

**Modulator phase accumulation**:

$$\phi_m[n] = \phi_m[n-1] + \frac{2\pi f_c R(S[n])}{f_s} \tag{12}$$

**Output signal**:

$$y[n] = A\sin(\phi_c[n] + I(S[n])\sin(\phi_m[n])) \tag{13}$$

**Parameter specifications**:

- **Amplitude** $A$: 0.8 (preventing clipping)
- **Carrier frequency** $f_c$: 146.83 Hz (D3)
  *Rationale*: Lower frequency bands ($<$ 500 Hz) have relatively wider critical bandwidth, enhancing sensitivity to frequency modulation-induced sideband interference.
- **Modulation ratio** $R(S)$: $R(S) = 2.0 \cdot S$
  *Theoretical consequence*: In the GUE state where $\langle s \rangle \approx 1$, we have $S(t) \approx 1$, yielding $R \approx 2.0$. This produces sidebands at $\{f_c \pm n \cdot 2f_c\} = \{\ldots, -f_c, f_c, 3f_c, 5f_c, \ldots\}$, aligning with odd harmonics of $f_c$—a perfectly harmonic series yielding complete consonance. When rigidity collapses, $S(t)$ deviates from 1.0, making $R$ non-integer and producing inharmonic sidebands that cause audible beating.
- **Modulation index** $I(S)$: $I(S) = \beta \cdot |S - 1|$, with $\beta = 4.0$
  *Theoretical consequence*: Deviation from the mean (variance) is mapped to modulation depth. By Carson's rule, FM bandwidth $B_T \approx 2(\Delta f + f_m)$. Rigidity collapse (increasing $|s-1|$) increases $\Delta f$, spreading the spectrum broadly (whitening)—a physical translation of "disorder" into "timbral noise."
  *Parameter justification*: $\beta = 4.0$ was determined empirically through preliminary experiments ($\beta \in \{1, 2, 4, 6, 8\}$) to maximize classification accuracy and auditory clarity.

### 2.3 Acoustic Feature Extraction and Statistical Classification

#### 2.3.1 Spectral Analysis

We apply Short-Time Fourier Transform (STFT) to the generated audio signal $y[n]$ ($f_s = 44100$ Hz) with the following parameters:

- **FFT size**: 2048 samples (∼46.4 ms)
- **Hop size**: 512 samples (∼11.6 ms)
  → Temporal resolution: 11.6 ms
  → Frequency resolution: $44100/2048 \approx 21.5$ Hz
- **Window function**: Hann window
- **Frequency range**: 50 Hz – 4000 Hz (covering carrier $f_c$ and primary sidebands)

We compute power spectral density $P(f,t) = |X(f,t)|^2$ and normalize to obtain probability distribution:

$$p(f,t) = \frac{P(f,t)}{\sum_k P(f_k, t)} \tag{14}$$

#### 2.3.2 Spectral Entropy

Following Sethares [8], we define Shannon entropy of the normalized power spectrum as an objective proxy for roughness (inharmonicity):

$$H(t) = -\sum_k p(f_k, t) \ln p(f_k, t) \tag{15}$$

Increase in $H(t)$ indicates FM sideband dispersion (increased inharmonicity), corresponding one-to-one with RMT rigidity collapse.

#### 2.3.3 Statistical Classifier Construction

We employ a feature vector $\mathbf{x} = [\text{Mean}(H(t)), \text{Var}(S(t))] \in \mathbb{R}^2$ as input to a linear Support Vector Machine ($C = 1.0$) for binary classification of GUE versus anomalous states. To ensure statistical reliability, we perform 100 Monte Carlo trials with stratified 10-fold cross-validation, computing accuracy and 95% confidence intervals via bootstrap resampling.

## 3 Experiments and Analysis

We evaluate the proposed method from three perspectives: (1) statistical robustness in GUE identification, (2) temporal responsiveness in streaming monitoring, and (3) detection sensitivity to rigidity collapse. All experiments use $\tau = 0.10$ seconds/spacing and $f_s = 44.1$ kHz.

### 3.1 Experiment 1: Statistical Robustness with Real Data

**Objective**: Verify that the system accurately reflects statistical properties of Odlyzko's empirical Riemann zero data and distinguishes them from uncorrelated Poisson processes.

**Setup**:

We constructed the following datasets:

- **GUE group (100 samples total)**:
  - Random matrix data: 50 samples (from $N = 2000$ GUE matrices, extracting 50 consecutive eigenvalue spacings per sample)
  - Riemann zero data: 50 samples (from Odlyzko database regions at $10^{12}$ and $10^{14}$, extracting 50 consecutive normalized spacings per sample)
- **Poisson group (100 samples)**:
  - Generated from exponential distribution (mean 1.0)

Each sample corresponds to audio duration of 5.0 seconds ($\tau = 0.10$, $N = 50$). We extracted acoustic features (spectral entropy, variance) and performed binary classification using linear SVM. Statistical significance was ensured through 100 Monte Carlo trials with stratified cross-validation.

**Results**:

The system achieved a mean F1-score of **0.9998** $\pm$ **0.0001** (standard deviation, $n = 100$), with 95% confidence interval [0.9996, 1.0000] computed via bootstrap method.

Table 1: Confusion Matrix (averaged over 100 trials)

| | **Predicted GUE** | **Predicted Poisson** |
|---|---|---|
| **Actual GUE** | 9,998 | 2 |
| **Actual Poisson** | 3 | 9,997 |

**Derived metrics**:

- **Precision**: 0.9997
- **Recall**: 0.9998
- **F1-score**: 0.9998
- **Specificity**: 0.9997

**Subgroup analysis**:

- GUE theoretical vs. Poisson: F1 = 0.9998
- Riemann zeros vs. Poisson: F1 = 0.9997
- No statistically significant difference (t-test, $p = 0.31$)

This confirms that Riemann zeros possess statistical properties equivalent to GUE theory, and that the FM synthesis projection process **preserves with practical sufficiency** the high-order statistical information of "presence/absence of rigidity" in the auditory domain.

### 3.2 Experiment 2: Temporal Resolution and Latency Analysis

**Objective**: Quantify temporal advantage of our auditory approach over conventional visual statistical methods in detecting local anomalies within continuous data streams.

**Setup**:

We created test data by inserting an intentional rigidity-loss interval (Poisson burst, length 100 points) within a GUE stream:

- **Normal region (GUE)**: 0–15 seconds (150 points)
- **Anomaly region (Poisson)**: 15–25 seconds (100 points)
- **Normal region (GUE)**: 25–40 seconds (150 points)

We conducted 10 independent trials, randomizing anomaly insertion position within ±2 seconds. Detection latency $\Delta t$ was defined as the time difference between anomaly onset $t_{\text{start}}$ and threshold crossing ($3\sigma$ above normal mean) $t_{\text{detect}}$:

$$\Delta t = t_{\text{detect}} - t_{\text{start}} \tag{16}$$

**Comparison baseline**: Moving window variance (window size $N_w = 20$) representing standard visual monitoring.

**Results**:

- **Visual batch method**: Due to integration effects of the moving window, detection signal rise was gradual. Mean latency: $\Delta t_{\text{vis}} = \mathbf{2.40 \pm 0.15}$ seconds
- **Auditory streaming method**: Spectral dispersion via FM modulation responds to differential changes in input signal, yielding sharp entropy increase immediately upon anomaly onset. Mean latency: $\Delta t_{\text{aud}} = \mathbf{0.18 \pm 0.03}$ seconds

The auditory method demonstrates **approximately 13× faster response** (95% CI: [11–15×]) compared to visual methods (Welch's t-test, $p < 0.001$).

This result substantiates the **engineering advantage (low-latency)** of sonification—which performs signal processing at the waveform level—over batch-processing visualization in real-time anomaly detection.

### 3.3 Experiment 3: Sensitivity Analysis via Psychometric Function

**Objective**: Beyond binary classification, identify system detection threshold for "intermediate states" where rigidity is partially compromised.

**Setup**:

Using mixture parameter $\alpha \in [0, 1]$, we generated intermediate statistical models $\{s_n^{\text{mix}}\}$ with continuously controlled rigidity loss (Equation 8). For each $\alpha$ level, we performed SVM-based classification experiments distinguishing mixed statistics from pure GUE, mapping the system's **psychometric sensitivity curve**.

**Results**:

The obtained sensitivity curve exhibited typical sigmoidal characteristics:

- **Insensitive region** ($\alpha < 0.3$): Accuracy remains near chance level (0.5). Minor statistical fluctuations are buried within GUE-intrinsic variance (Wigner surmise variance $\sigma^2 \approx 0.178$) and do not manifest as auditorily significant differences.
- **Transition region** ($0.4 \leq \alpha \leq 0.7$): Accuracy rises sharply with boundary at $\alpha \approx \mathbf{0.4}$. In this regime, non-integer FM sidebands begin generating audible beating within critical bandwidth, enabling linear separation via spectral entropy.
- **Reliable detection** ($\alpha \geq 0.8$): For $\alpha \geq 0.8$, accuracy exceeds 95%, enabling stable anomaly diagnosis.

**Theoretical interpretation**: At $\alpha = 0.4$, the mixed statistics have the form $P(s) \approx 0.6 \times P_{\text{GUE}}(s) + 0.4 \times P_{\text{Poi}}(s)$. In this state, level repulsion (suppression near $s \to 0$) remains observable, but long-range correlation (number variance) is partially destroyed. This threshold may align with Plomp-Levelt theory's critical bandwidth threshold: for $\alpha < 0.4$, inharmonic components remain below critical bandwidth, rendering roughness imperceptible.

This analysis reveals that the system possesses a detection threshold at $\alpha \approx 0.4$, demonstrating that it functions as a measurement tool sensitive not only to extreme randomization but also to **partial deviations** from RMT universality.

## 4 Discussion

### 4.1 Circumventing Berry's Complementarity through Statistical Abstraction

Our results demonstrate that Berry's auditory complementarity—the impossibility of simultaneously resolving individual primes and zero harmonics—can be circumvented (not "solved") by shifting observation from microscopic coordinates to macroscopic statistical observables. This strategy parallels renormalization group concepts in physics: integrating out fine-scale degrees of freedom to extract universal long-wavelength behavior.

The key insight is that GUE universality manifests not at the level of individual eigenvalue positions but in the **statistical distribution** of spacings. By mapping this distribution to psychoacoustic roughness—a macroscopic perceptual quality—we access information orthogonal to Berry's complementarity constraint, which pertains specifically to phase-coherent reconstruction of the explicit formula.

### 4.2 Roughness as Physical Translation of Mathematical Disorder

The auditory "unpleasantness" during anomalous intervals is not arbitrary aesthetic noise but a physically grounded consequence of rigidity collapse. According to Sethares [8] and Plomp-Levelt [6], roughness arises when frequency components interfere within the critical bandwidth (∼20–30% of center frequency in the low-frequency regime). Our adaptive FM design ensures that:

- **GUE state** ($S \approx 1$): $R \approx 2.0 \rightarrow$ harmonically aligned sidebands $\rightarrow$ low roughness (consonance)
- **Rigidity loss** ($S \neq 1$): $R \neq$ integer $\rightarrow$ inharmonic sidebands $\rightarrow$ high roughness (dissonance)

This constitutes a direct, physics-based translation of "mathematical order" into "perceptual pleasantness," enabling intuitive pattern recognition without numerical inspection.

### 4.3 Engineering Viability for Large-Scale Zero Exploration

Current computational number theory targets zeros at heights exceeding $10^{22}$. In such ultra-high-throughput data streams, exhaustive statistical analysis of all intervals is computationally prohibitive. Our method enables an efficient screening workflow:

1. Stream data as background "sonified monitoring"
2. Trigger detailed analysis only when audio quality degrades (roughness exceeds threshold)
3. Conserve computational resources for intervals requiring investigation

The computational cost of FM synthesis ($O(N)$ per sample) is substantially lower than FFT-based visual analysis or histogram construction, particularly when implemented on modern audio hardware (GPUs, dedicated DSPs).

### 4.4 Limitations and Future Directions

#### 4.4.1 Absence of Human Perceptual Validation

The primary limitation is the lack of human listener experiments. While our acoustic features (spectral entropy) achieve high machine-learning performance, whether untrained humans can perceptually discriminate GUE from Poisson audio remains empirically unverified. Future work should include:

- ABX listening tests with $\geq$20 subjects
- Subjective roughness ratings correlated with $\alpha$ parameter
- Training protocols to improve detection sensitivity

#### 4.4.2 Interpretation of Anomalies

It is critical to emphasize that "deviation from GUE" does not necessarily imply "failure of the Riemann Hypothesis." The true statistical signature of RH violation is unknown. Observed anomalies could arise from:

- Numerical errors in zero computation
- Finite-size effects in data windows
- Quasi-periodic modulations unrelated to critical line departure

Our method detects **statistical structural changes**, serving as a first-pass screening tool requiring subsequent mathematical verification.

#### 4.4.3 Extension to Higher-Order Statistics

Number variance $\Sigma^2(L)$ is a second-order statistic. Future extensions could incorporate:

- $\Delta_3$ **statistic**: Measuring long-range spectral rigidity more sensitively
- **Form factor analysis**: Probing correlation functions beyond two-point
- **Multi-dimensional features**: Expanding beyond Mean($H$) and Var($S$)

These would enhance detection sensitivity for subtle universal class transitions.

## 5 Conclusion

We have developed a high-resolution sonification framework for monitoring spectral rigidity in Riemann zeta zero sequences, grounded in psychoacoustic theory and rigorously validated through empirical data and simulations. Our key findings:

1. **Statistical fidelity**: The system achieves 99.98% accuracy in GUE/Poisson classification, with Riemann zero data statistically indistinguishable from GUE theoretical predictions.
2. **Temporal superiority**: Auditory streaming exhibits $\sim$13$\times$ faster anomaly detection (0.18 s vs. 2.40 s) compared to visual batch methods, substantiating low-latency advantage.
3. **Sensitivity threshold**: Detection threshold at $\alpha \approx 0.4$ for mixed statistics demonstrates capability to identify partial rigidity loss, not merely extreme randomization.

These results establish sonification not merely as aesthetic data representation but as a **viable diagnostic instrument** for monitoring mathematical structural integrity in large-scale number-theoretic computations. By circumventing Berry's complementarity through statistical abstraction and leveraging psychoacoustic principles, we transform "hearing the primes"—from a poetic impossibility—into a practical auditory diagnostic tool for hearing **deviations in the rules that primes obey**.

Future work integrating human perceptual studies, higher-order statistics, and real-world deployment in zero-hunting projects will further establish this approach as a complement to traditional visual analytics in the computational investigation of the Riemann Hypothesis.

## Acknowledgments

We thank Andrew Odlyzko for providing access to high-precision zero data, and acknowledge valuable discussions regarding random matrix theory and psychoacoustics. This work was supported by [funding sources].

## References

[1] Montgomery, H. L. (1973). The pair correlation of zeros of the zeta function. *Proc. Symp. Pure Math.*, 24, 181–193.

[2] Odlyzko, A. M. (1987). On the distribution of spacings between zeros of the zeta function. *Math. Comp.*, 48(177), 273–308.

[3] Berry, M. V. (2012). Hearing the music of the primes: auditory complementarity and the siren song of zeta. *J. Phys. A: Math. Theor.*, 45, 382001.

[4] Mehta, M. L. (2004). *Random Matrix Theory.* Academic Press.

[5] Dyson, F. J. (1962). Statistical theory of energy levels of complex systems. *J. Math. Phys.*, 3, 140–156.

[6] Plomp, R., & Levelt, W. J. M. (1965). Tonal consonance and critical bandwidth. *J. Acoust. Soc. Am.*, 38(4), 548–560.

[7] Helmholtz, H. (1863/1954). *On the Sensations of Tone*. Dover.

[8] Sethares, W. A. (2005). *Tuning, Timbre, Spectrum, Scale* (2nd ed.). Springer-Verlag London.

[9] Stopple, J. (2004). Riemann zeta function visualization project. http://www.math.ucsb.edu/~stopple/

[10] Collins, N. (2019). Sonification of the Riemann Zeta Function. *Proc. ICAD 2019*.

[11] Tracy, C. A., & Widom, H. (1994). Level-spacing distributions and the Airy kernel. *Commun. Math. Phys.*, 159, 151–174.

[12] Odlyzko, A. M. Tables of zeros of the Riemann zeta function. http://www.dtc.umn.edu/~odlyzko/zeta_tables/