

Arctonyx collaris, Greater Hog Badger

Amendment version

Assessment by: Duckworth, J.W., Timmins, R.J., Chutipong, W., Gray, T.N.E., Long, B., Helgen, K., Rahman, H., Choudhury, A.U. & Willcox, D.H.A.



View on www.iucnredlist.org

Citation: Duckworth, J.W., Timmins, R.J., Chutipong, W., Gray, T.N.E., Long, B., Helgen, K., Rahman, H., Choudhury, A.U. & Willcox, D.H.A. 2024. *Arctonyx collaris* (amended version of 2016 assessment). *The IUCN Red List of Threatened Species* 2024: e.T70205537A259343689. <https://dx.doi.org/10.2305/IUCN.UK.2024-2.RLTS.T70205537A259343689.en>

Copyright: © 2024 International Union for Conservation of Nature and Natural Resources

Reproduction of this publication for educational or other non-commercial purposes is authorized without prior written permission from the copyright holder provided the source is fully acknowledged.

Reproduction of this publication for resale, reposting or other commercial purposes is prohibited without prior written permission from the copyright holder. For further details see [Terms of Use](#).

The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission](#) (SSC) and [The IUCN Red List Partnership](#). The IUCN Red List Partners are: [ABQ BioPark](#); [Arizona State University](#); [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [Missouri Botanical Garden](#); [NatureServe](#); [Re:wild](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Senckenberg Society for Nature Research](#); [Texas A&M University](#); and [Zoological Society of London](#).

If you see any errors or have any questions or suggestions on what is shown in this document, please provide us with [feedback](#) so that we can correct or extend the information provided.

Taxonomy

Kingdom	Phylum	Class	Order	Family
Animalia	Chordata	Mammalia	Carnivora	Mustelidae

Scientific Name: *Arctonyx collaris* F.G. Cuvier, 1825

Synonym(s):

- *Arctonyx annaeus* Thomas, 1921
- *Arctonyx dictator* Thomas, 1910

Common Name(s):

- English: Greater Hog Badger, Hog Badger
- German: Schweinsdachs

Taxonomic Source(s):

Helgen, K.M., Lim, N.T-L. and Helgen, L.E. 2008. The hog-badger is not an edentate: systematics and evolution of the genus *Arctonyx* (Mammalia: Mustelidae). *Zoological Journal of the Linnean Society* 154: 353-385.

Taxonomic Notes:

For many decades the genus *Arctonyx* was universally considered monospecific, under the oldest available name, *A. collaris*. It has been proposed that the genus contains multiple species: Sumatran Hog Badger *A. hoevenii* is disjunct from the rest of the genus's range, whilst Northern Hog Badger *A. albogularis* and Greater Hog Badger *A. collaris* (*sensu stricto*) occur in mainland Asia, with a poorly understood zone of overlap or contact (Helgen *et al.* 2008). This revision was not accepted by Choudhury (2013) who furthermore queried the neotype and accompanying type locality selected for *A. collaris* by Helgen *et al.* (2008). Nonetheless, Helgen *et al.* (2008: 369) were explicit that "the absolute, striking distinctions in size, pelage and qualitative cranial morphology between available adult samples of *A. albogularis* and *A. collaris* indicate no intergradation between these taxa". Therefore, three species of hog badger are recognized here.

Assessment Information

Red List Category & Criteria: Vulnerable A2cd+3cd+4cd [ver 3.1](#)

Year Published: 2024

Date Assessed: March 3, 2015

Justification:

Greater Hog Badger is listed as Vulnerable because it is undergoing a hunting-driven population decline. Although rates in the eastern parts of its range, notably Viet Nam and Lao PDR, on average probably exceed 50% per 15 years (three generations), in large parts of the west and centre they may not even reach rates appropriate for Near Threatened listing (20-25%). Thus, Vulnerable is a global average. There is no reason to expect the decline rates to slow down; in particular, in Myanmar and Cambodia they may be increasing, fuelled by rising affluence and market connectivity. The situation in Cambodia is

particularly likely to worsen, with increasing snaring in the eastern provinces that border Viet Nam, a known demand for traditional medicine that might be widespread in rural areas, and ongoing massive clearance of natural habitat for agro-industrial concessions. This species should be periodically reassessed for the Red List in light of ongoing threats and uncertainty about range-wide levels of exploitation and the effects these are having on wild populations. The change from the recent Near Threatened listing of *Arctonyx collaris* (*sensu lato*) reflects the taxonomic revision, excluding the populations of Sumatra (where ongoing declines are likely to be negligible) and most of China (where declines are also likely to be lower than average for Greater Hog Badger).

Previously Published Red List Assessments

2016 – Vulnerable (VU)

Geographic Range

Range Description:

Greater Hog Badger occurs from Bangladesh and North-east India east through Myanmar, Thailand and Lao PDR to Vietnam and south to Cambodia and peninsula Thailand; it probably also inhabits Yunnan province, China. Inconclusive reports from the state of Perak, northernmost peninsular Malaysia, have never been confirmed (Helgen *et al.* 2008).

The range in Bangladesh and North-east India is provisional. Because Helgen *et al.* (2008) traced only a few hog badger records from this area that could be assigned to species, both mainland hog badger species might occur in this area: so the many recent records simply identified as 'hog badger' should not, at this stage, be identified to species on any presumption of range or habitat. In North-east India the genus is well distributed all over, with most records taken to be this species including animals observed in Manas NP, north(west) of the Brahmaputra (Choudhury 2013). This contests the proposal of Helgen *et al.* (2008) that said drainage forms the species's (north)west range boundary.

In Bangladesh, the genus occurs widely, with records from the South-east (Chittagong and Cox's Bazar district and Chittagong Hill Tracts), Dudhpukuria-Dhopacharia Wildlife Sanctuary (Feeroz *et al.* 2012), Chunoti Wildlife Sanctuary, Fasiakhali Wildlife Sanctuary, Chittagong University campus (outside the city and middle of hills), and from the North-east (Sylhet division; Moulovibazar and Habigonj districts), adjacent to Lawachara National Park (all records per Hasan Rahman pers. comm. 2014). These records are likely all to relate to this species, although objective confirmation is desirable.

Almost all hog badger specimens examined by Helgen *et al.* (2008) from China were judged to be *A. albogularis*, but two specimens from Lichiang, Yunnan, were "provisionally" assigned to Greater Hog Badger.

Than Zaw *et al.* (2008) traced recent hog badger records (all assumed here to be Greater Hog Badger; however *A. albogularis* might yet be found to occur in the far north) widely across north and central Myanmar south to about 20°N, and historical records south to about 16°30'N; survey effort was insufficient for the lack of modern records in the southern known range to be taken to suggest recent extirpation.

Surveys in 21 areas spread across Thailand found hog badger (all assumed here to be Greater Hog

Badger) to be the third most widely camera-trapped small carnivore. The southernmost recent records in the country are from Tai Rom Yen National Park (8°40'N). There are historical records from further south in Thailand, but recent camera-trap survey effort south of Tai Rom Yen NP was rather limited except for in one area right on the Malaysia border (Chutipong *et al.* 2014).

Records in Lao PDR (all assumed here to be Greater Hog Badger) come from a wide spread of localities in the North, Centre and South. Occurrence seems to be patchy, with no recent records from the Mekong plain, which covers most of the western part of the southern half of the country (e.g., Duckworth 1997, Duckworth *et al.* 1999, Johnson *et al.* 2009, Coudrat *et al.* 2014, Gray *et al.* 2014b).

Robertson (2007) traced confirmed field records for the genus, all assumed to be Greater Hog Badger, from most of Viet Nam, occurrence was patchy and no records were found for the Mekong Delta. The current distribution in Viet Nam is likely to be highly fragmented. A recent review of small carnivore camera-trap records failed to produce any records additional to those in Robertson (2007), despite several surveys within suitable habitat, and within some of the country's largest protected areas (Willcox *et al.* 2014: Table SOM3). As of mid 2014, only two camera-trap records were known for the entire country (D.H.A. Willcox pers. comm. 2014), from Pu Mat Nature Reserve (SFNC 2000) and Pu Luong NR (Do Ngoc Cuong 2004). The species has subsequently been camera-trapped in Ngoc Linh Nature Reserve, Kon Tum province (Nguyen T. pers. comm. 2014).

Although there were few historical Cambodian records, camera-trapping has recently found it at various sites spread almost throughout the country (e.g. Holden and Neang 2009, Royan 2010, Gray *et al.* 2014a), including Preah Vihear Protected Forest and Seima Protected Forest (WCS Cambodia unpublished data per D.H.A. Willcox pers. comm. 2014).


It has been recorded from sea-level to at least 2,300 m (see 'Habitat and ecology' section).

Country Occurrence:

Native, Extant (resident): Bangladesh; Cambodia; India; Lao People's Democratic Republic; Myanmar; Thailand; Viet Nam

Native, Presence Uncertain: China; Malaysia (Peninsular Malaysia)

Distribution Map




Legend

- EXTANT (RESIDENT)
- POSSIBLY EXTINCT
- PRESENCE UNCERTAIN

Compiled by:

IUCN (International Union for Conservation of Nature) 2016



The boundaries and names shown and the designations used on this map do not imply any official endorsement, acceptance or opinion by IUCN.

Population

Greater Hog Badger populations have not been studied in sufficient detail for densities or trends to be known in any part of the range. Camera-trap encounter rates vary widely across its range: in some it is among the most commonly and widely photographed small carnivores, such as in Thailand (Chutipong *et al.* 2014) and Nakai–Nam Theun National Protected Area, Lao PDR (Coudrat *et al.* 2014), in others it is only rarely found if at all (e.g., modern Viet Nam: Willcox *et al.* 2014: Table SOM3).

At least two factors might produce this heterogeneous recording pattern: (i) natural variation in densities and (ii) human-induced declines proceeding at different rates across the range. Both are probably important. Past information is rarely precise enough to assess even the coarse population trend directly. However, judging by patterns in better documented hunting-sensitive groups of ground-dwelling mammals (e.g., wild cattle *Bos*, Leopard *Panthera pardus*, Sambar *Rusa unicolor*), it would be reasonable to expect human pressure to have depressed densities in Lao PDR and Viet Nam more than in Thailand and Cambodia. And this indeed seems to have happened. Recent camera-trapping has recorded the species in only very few areas in Viet Nam (Willcox *et al.* 2014: Table SOM3, Nguyen T. pers. comm. 2014); Robertson (2007) had already drawn attention to the surprising paucity of records in the country in the 1990s and early 2000s. In Lao PDR, by contrast, all published large-scale camera-trap surveys have found the species (Nam Et–Phou Louey NPA, Johnson *et al.* 2009; Nakai–Nam Theun NPA, Coudrat *et al.* 2014; Xe Sap NPA, Gray *et al.* 2014b), indicating a healthier status than in Vietnam (as is typical for hunting-sensitive species). The three camera-trap survey areas are likely to hold amongst the least-depleted large mammal faunas in Lao PDR. All are in rugged mountainous areas and indeed all post-1990 Lao records of Hog Badger come from hill and montane areas; there has been none from the plains. This pattern, making due allowance for the distribution of survey effort, was already apparent in the pre-camera-trapping era by 1999, based on direct sightings and records of dead animals or their parts (Duckworth *et al.* 1999). By contrast, this species is commonly recorded at low altitudes, including in plains landscapes, of Thailand and Cambodia (Royan 2010, Gray *et al.* 2014a, Chutipong *et al.* 2014) and it is common in similar lowland terrain in Kaziranga NP, Assam (Choudhury 2013). This suggests strongly that the absence of recent Lao records from such terrain (where hunting is more efficient than in rugged terrain, meaning that hunting-sensitive species across a range of vertebrate taxa tend to have declined faster there) reflects steep declines (and plausibly widespread extirpation) there.

The occurrence in Myanmar is also patchy without obvious natural explanation (Than Zaw *et al.* 2008); thus far, the pattern of records has not been compared closely with those of species known to be sensitive to hunting. In Cambodia, while it has been recorded relatively commonly in some areas such as parts of the north-east (Gray *et al.* 2014a), in the 'northern plains', west of the Mekong, it was not recorded in 6,000 camera-trap nights of survey in the 2012–2013 and 2013–2014 dry seasons, and the most recent record there seems to have been in 2001 (D. H. A. Willcox pers. comm. 2014). This area abuts Lao PDR and this apparent range retreat corroborates the suggestion that it has been hunted out from such habitat in Lao PDR. In Thailand, it seems to remain generally common with no large landscapes of (semi-) natural habitat lacking records (when patterns in survey effort are allowed for) (Chutipong *et al.* 2014). In North-east India the genus, although widespread, is generally not common (Choudhury 2013). The population status in its Chinese range (part of Yunnan province) seems poorly known, even at the level of genus.

Current Population Trend: Decreasing

Habitat and Ecology (see Appendix for additional information)

Greater Hog Badger is active throughout the day and night (despite past statements that the genus is nocturnal), ground-dwelling, and not very wary of people (e.g., Pocock 1941, Duckworth *et al.* 1999, Helgen *et al.* 2008, Than Zaw *et al.* 2008, Johnson *et al.* 2009, Holden and Neang 2009, Coudrat *et al.* 2014, Gray *et al.* 2014b).

Greater Hog Badger occurs down to the extreme lowlands (e.g. Bottom-Soaker National Park, Cambodia, and Kaziranga NP, Assam, India; Royan 2010, Choudhury 2013) up into the montane zone. The genus has been found up to 3,500 m, but Helgen *et al.* (2008) traced no record of this species above the relatively modest altitude of 1,500 m. Similarly, and based largely on different sets of records, the highest-altitude records traced for various countries are, Vietnam, 950 m (Robertson 2007); Cambodia, 1,200 m (Holden and Neang 2009); Myanmar, 1,500 m (Than Zaw *et al.* 2008); and Thailand, 1,016 m (Chutipong *et al.* 2014). However, as cautioned by Than Zaw *et al.* (2008), relevant survey effort above 1,500 m has been relatively limited in Myanmar and this is generally true throughout this species's range. Thus, the typical upper altitudinal limit may be found to be somewhat, even considerably, higher, as supported by records in Lao PDR up to nearly 2,300 m (Johnson *et al.* 2009). Choudhury (2013) recorded the genus up to 2,000 m in India, but suggested that the higher-altitude records would be of *A. albogularis*.

This species occurs across a wide variety of habitats, from heavy forest (both deciduous and evergreen) to the non-forested 'countryside' in at least North-east India; this includes grassland-dominated floodplains such as Kaziranga NP in Assam (Choudhury 2013), where it is one of the most frequently camera-trapped small carnivores (D. Mudappa pers. comm. 2014). There are also records in Bangladesh (assumed to be this species) from tea-gardens and other non-forest habitats (Hasan Rahman pers. comm. 2014). In South-east Asia most records come from forests. The extent of actual and potential occupation of non-forest habitats is unknown: concerted survey in them is rare and off-take of mammals of this size is heavy in much of this species's range. Thus, it is possible that they would occur widely in non-forest parts of South-east Asia but are currently excluded by hunting levels.

The diet is poorly known, but Helgen *et al.* (2008) speculated that it might be partly to strongly vermivorous. This was based on morphological attributes more than on direct dietary information. A strong dependence on worms would neatly explain the pattern of current habitat use in Lao PDR (see 'population') - widespread and still locally common in montane areas but nearly or effectively extirpated from the plains. The country's plains have a harsh dry season, when the ground even within semi-evergreen forest (true evergreen forest is highly localised in the country's lowlands) bakes hard. Finding large numbers of worms under such conditions is presumably difficult and highly demanding of energy. By contrast, in the mountains, forest soil remains damper through the dry season, as a result of cooler temperatures, the sometimes frequent mists and cloud, and the fuller shading of true evergreen forest. Thus, in regions with a harsh dry season, it is reasonable to suppose higher population densities in hills and mountains than in plains and low hills, and a higher resilience to hunting. Such seasonality occurs almost across the species's range excepting most of Vietnam, the Thai-Malay peninsula and locally around massifs large enough to generate their own precipitation. Nonetheless, the species does occur in plains regions of Cambodia and Thailand with dry seasons as or more harsh than in Lao PDR; these areas have somewhat to much lower hunting pressure.

Systems: Terrestrial

Use and Trade

To varying extents across its range, Greater Hog Badger is eaten, sometimes with a medicinal purpose, e.g. in the 'northern plains' (Ai Suzuki pers. comm. 2014) and the 'eastern plains' of Cambodia (Gray *et al.* 2014a). In Lao PDR, the palatability of this hog badger varies between ethnic groups, with some groups disliking the taste, whereas groups in parts of the Nam Theun basin (and presumably elsewhere) esteem the species as food (J.P. Baker pers. comm. to Duckworth *et al.* 1999, J. Chamberlain pers. comm. to Duckworth *et al.* 1999). The genus, presumably including this species, is also eaten by some groups in India (A.U. Choudhury pers. comm. 2006), and is hunted as well as farmed for food in China (M.W.N. Lao pers. comm. 2006). Across most of its range, most of the catch is probably not targeted, but is by non-selective methods for wildlife meat, or opportunistically during activities other than hunting.

Threats (see Appendix for additional information)

The heterogeneous pattern of Greater Hog Badger detections and the variation in encounter rates, with the lowest rates in the very heavily hunted countries of Viet Nam and Lao PDR, strongly suggest that the species is in hunting-driven decline. Based on Lao PDR, extirpation, or decline to densities making detections extremely unlikely, seems to have proceeded faster in areas with marked dry seasons than in mountain evergreen forests with a more benign dry season, presumably because starting populations were higher in the latter habitats than in the former (see 'Habitats and ecology'). In Viet Nam, trade-driven hunting rose to industrial levels a decade or so earlier than in Lao PDR. The low number of camera-trap survey areas in Viet Nam with records of the species indicate that even in the presumed optimal habitats of evergreen forest with only a mild dry season, steep declines and, presumably, extirpation can occur. To date, hunting is at much lower levels in the rest of this species's range. In Lao PDR and Vietnam, much hunting involves metal cable snares set in long lines linked by drift fences, an approach very effective in catching ground-dwelling animals (e.g., Coudrat *et al.* 2014, Willcox *et al.* 2014); these snares are, mostly, big enough to secure Greater Hog Badger, because large ungulates are amongst the main targets. The geographical use of such snaring has expanded rapidly over the last 20 years and evidently continues to do so. In much of the species's range, most people entering forest and other natural habitat for whatever reason are accompanied by dogs, many have firearms, and most secure opportunistically any animal of hog badger size-class that they come across. Being partly diurnal, ground-dwelling and not prone to shyness, hog badgers must be particularly at risk to such opportunistic off-take.

The threat which habitat change poses directly to this species is not clear. Indirectly, anything that opens up access to areas previously difficult to enter, in particular which allows vehicular removal of wildlife for sale, exacerbates the hunting threat. Almost all forms of habitat change by people co-occur with increased human access and presence, making the indirect threat from habitat change severe. Based on localities of specific records, it is unlikely that degraded and fragmented natural habitat are inherently unsuitable. These are not widely occupied probably because their higher densities of people, with concomitantly increased likelihood for badgers of lethal incidental encounters with people, prevent this. Outright conversion to entirely artificial habitat (agriculture, settlement and infrastructure) seems more likely to render an area unsuitable for Greater Hog Badger, but even this remains to be proven. Overall, the existence of large tracts of forest in the eastern parts of the species's range with at best low densities of this species indicate that there the chief threat is hunting. While perhaps this is not yet so in parts of the western range, it is increasingly likely to become so there, too.

In Kaziranga National Park, India, numbers die in the periodic inundation of this flood-plain (Choudhury 2013). While formerly such animals might have been able to move safely to higher ground, the increasing fragmentation of natural habitat probably leads to higher mortalities nowadays. Nonetheless, in total these are likely to be insignificant to overall decline rates of the species, and are only a potential threat at the most local of levels.

Hog badger hairs are used in European shaving brush manufacture (Domingo-Roura *et al.* 2006) but whether this species and wild populations are involved is unclear; the large farmed stock in China (Lau *et al.* 2010) seems a more likely origin.

Conservation Actions (see Appendix for additional information)

Throughout its range, Greater Hog Badger is found in a number of protected areas, perhaps particularly in Myanmar, India and Thailand (Than Zaw *et al.* 2008, Choudhury 2013, Chutipong *et al.* 2014). This species is protected by law in Thailand (Chutipong *et al.* 2014), in Bangladesh (as a Schedule 1 species) under the Wildlife Act 2012 (Hasan Rahman pers. comm. 2014), and in India, under the highest level of protection (Choudhury 2013). This species is not protected in Viet Nam or Cambodia and is the largest-bodied unprotected mammal, except for Eurasian Wild Hog *Sus scrofa*, in Myanmar (Than Zaw *et al.* 2008). The China Red List has listed hog badgers (as a monospecific genus, under this species's name) as Vulnerable under C1 and A2c.

Credits

Assessor(s): Duckworth, J.W., Timmins, R.J., Chutipong, W., Gray, T.N.E., Long, B., Helgen, K., Rahman, H., Choudhury, A.U. & Willcox, D.H.A.

Reviewer(s): Schipper, J.

Contributor(s): Zaw, T.

Authority/Authorities: IUCN SSC Small Carnivore Specialist Group

Bibliography

Choudhury, A. 2013. *The mammals of North east India*. Gibbon Books and the Rhino Foundation for Nature in NE India, Guwahati, Assam, India.

Chutipong, W., Tantipisanuh, N., Ngoprasert, D., Lynam, A.J., Steinmetz, R., Jenks, K.E., Grassman Jr., L.I., Tewes, M., Kitamura, S., Baker, M.C., McShea, W., Bhumpakphan, N., Sukmasuang, R., Gale, G.A., Harich, F.K., Treydte, A.C., Cutter, P., Cutter, P.B., Suwanrat, S., Siripattaranukul, K., Hala-Bala Wildlife Research Station, Wildlife Research Division and Duckworth, J.W. 2014. Current distribution and conservation status of small carnivores in Thailand: a baseline review. *Small Carnivore Conservation* 51: 96–136.

Coudrat, C.N.Z., Nanthavong, C., Sayavong, S., Johnson, A., Johnston, J.B. and Robichaud, W.G. 2014. Conservation importance of Nakai-Nam Theun National Protected Area, Laos, for small carnivores based on camera trap data. *Raffles Bulletin of Zoology* 62: 31–49.

Domingo-Roura, X., Marmi, J., Ferrando, A., López-Giráldez, J.F., Macdonald, D.W., Jansman, H.A.H. 2006. Badger hair in shaving brushes comes from protected Eurasian badgers. *Biological Conservation* 128: 425–430.

Duckworth, J.W. 1997. Small carnivores in Laos: a status review with notes on ecology, behaviour and conservation. *Small Carnivore Conservation* 16: 1–21.

Duckworth, J.W., Salter, R.E. and Khounbolin, K. 1999. Wildlife in Lao PDR: 1999 Status Report. IUCN, Vientiane, Laos.

Feeroz, M.M., Hasan, M.K. and Hossain, M.K. 2012. *Biodiversity of protected areas of Bangladesh, Vol. 2: Dudpukuria-Dhopacharia Wildlife Sanctuary*. Bio Track. Arannayak Foundation, Dhaka, Bangladesh.

Gray, T.N.E., Pin C., Phan C., Crouthers, R., Kamler, J.F. and Prum S. 2014a. Camera-trap records of small carnivores from eastern Cambodia, 1999–2013. *Small Carnivore Conservation* 50: 20–24.

Gray, T.N.E., Thongsamouth, K. and Tilker, A. 2014b. Recent cameratrap records of Owston's Civet *Chrotogale owstoni* and other small carnivores from Xe Sap National Protected Area, southern Lao PDR. *Small Carnivore Conservation* 51: 29–33.

Helgen, K.M., Lim, N.T.-L. and Helgen, L.E. 2008. The hog-badger is not an edentate: systematics and evolution of the genus *Arctonyx* (Mammalia: Mustelidae). *Zoological Journal of the Linnean Society* 154: 353–385.

Holden, J. and Neang T. 2009. Small carnivore records from the Cardamom Mountains, southwestern Cambodia. *Small Carnivore Conservation* 40: 16–21.

IUCN. 2016. The IUCN Red List of Threatened Species. Version 2016-1. Available at: www.iucnredlist.org. (Accessed: 30 June 2016).

IUCN. 2024. The IUCN Red List of Threatened Species. Version 2024-2. Available at: www.iucnredlist.org. (Accessed: 28 October 2024).

Johnson, A., Vongkhamheng, C. and Saithongdam, T. 2009. The diversity, status and conservation of small carnivores in a montane tropical forest in northern Laos. *Oryx* 43: 626–633.

Lau, M.W.N., Fellowes, J.R. and Chan, B.P.L. 2010. Carnivores (Mammalia: Carnivora) in South China: a status review with notes on the commercial trade. *Mammal Review* 42: 247–292.

Pocock, R.I. 1941. *The Fauna of British India, including Ceylon and Burma*. Taylor & Francis, Ltd., London, UK.

Robertson, S.I. 2007. Status and conservation of small carnivores in Vietnam. University of East Anglia, Norwich, U.K. (Ph.D. thesis).

Royan, A. 2010. Significant mammal records from Botum–Sakor National Park, southwest Cambodia. *Cambodian Journal of Natural History* '2010'(1): 22–26.

SFNC. 2000. *Pù Mát: a biodiversity survey of a Vietnamese protected area*. SFNC Project, Vinh, Vietnam.

Than Zaw, Saw Htun, Saw Htoo Tha Po, Myint Maung, Lynam, A.J., Kyaw Thinn Latt and Duckworth, J.W. 2008. Status and distribution of small carnivores in Myanmar. *Small Carnivore Conservation* 38: 2–28.

Timmins, R.J. and Duckworth, J.W. 1999. Status and conservation of Douc Langurs (*Pygathrix nemaeus*) in Laos. *International Journal of Primatology* 20: 469 – 489.

Timmins, R.J., Do Tuoc, Trinh Viet Cuong and Hendrichsen, D.K. 1999. A preliminary assessment of the conservation importance and conservation priorities of the Phong Nha-Ke Bang proposed National Park, Quang Binh Province, Vietnam. Fauna and Flora International-Indochina Programme, Hanoi, Vietnam.

Willcox, D.H.A., Tran Q.P., Hoang M.D. and Nguyen T.T.A. 2014. The decline of non-*Panthera* cat species in Vietnam. *Cat News* Special Issue 8: 53–61.

Citation

Duckworth, J.W., Timmins, R.J., Chutipong, W., Gray, T.N.E., Long, B., Helgen, K., Rahman, H., Choudhury, A.U. & Willcox, D.H.A. 2024. *Arctonyx collaris* (amended version of 2016 assessment). *The IUCN Red List of Threatened Species* 2024: e.T70205537A259343689. <https://dx.doi.org/10.2305/IUCN.UK.2024-2.RLTS.T70205537A259343689.en>

Disclaimer

To make use of this information, please check the [Terms of Use](#).

External Resources

For [Supplementary Material](#), and for [Images and External Links to Additional Information](#), please see the Red List website.

Appendix

Habitats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Habitat	Season	Suitability	Major Importance?
1. Forest -> 1.5. Forest - Subtropical/Tropical Dry	Resident	Suitable	No
1. Forest -> 1.6. Forest - Subtropical/Tropical Moist Lowland	Resident	Suitable	Yes
1. Forest -> 1.9. Forest - Subtropical/Tropical Moist Montane	Resident	Suitable	Yes
2. Savanna -> 2.1. Savanna - Dry	Resident	Marginal	-
2. Savanna -> 2.2. Savanna - Moist	Resident	Marginal	-
3. Shrubland -> 3.5. Shrubland - Subtropical/Tropical Dry	Resident	Marginal	-
3. Shrubland -> 3.6. Shrubland - Subtropical/Tropical Moist	Resident	Marginal	-
4. Grassland -> 4.5. Grassland - Subtropical/Tropical Dry	Resident	Marginal	-
4. Grassland -> 4.6. Grassland - Subtropical/Tropical Seasonally Wet/Flooded	Resident	Suitable	Yes
14. Artificial/Terrestrial -> 14.6. Artificial/Terrestrial - Subtropical/Tropical Heavily Degraded Former Forest	Resident	Unknown	-

Use and Trade

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

End Use	Local	National	International
1. Food - human	Yes	No	No
16. Establishing ex-situ production *	No	No	No

Threats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Threat	Timing	Scope	Severity
1. Residential & commercial development -> 1.1. Housing & urban areas	Ongoing	Minority (<50%)	Negligible declines
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation	
1. Residential & commercial development -> 1.2. Commercial & industrial areas	Ongoing	Minority (<50%)	Negligible declines
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation	

2. Agriculture & aquaculture -> 2.1. Annual & perennial non-timber crops -> 2.1.1. Shifting agriculture	Ongoing	Majority (50-90%)	Slow, significant declines
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation	
2. Agriculture & aquaculture -> 2.1. Annual & perennial non-timber crops -> 2.1.2. Small-holder farming	Ongoing	Majority (50-90%)	Slow, significant declines
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation	
2. Agriculture & aquaculture -> 2.1. Annual & perennial non-timber crops -> 2.1.3. Agro-industry farming	Ongoing	Majority (50-90%)	Slow, significant declines
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation	
2. Agriculture & aquaculture -> 2.2. Wood & pulp plantations -> 2.2.1. Small-holder plantations	Ongoing	Majority (50-90%)	Slow, significant declines
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation	
2. Agriculture & aquaculture -> 2.2. Wood & pulp plantations -> 2.2.2. Agro-industry plantations	Ongoing	Minority (<50%)	Slow, significant declines
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation	
3. Energy production & mining -> 3.2. Mining & quarrying	Ongoing	Minority (<50%)	Slow, significant declines
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation	
3. Energy production & mining -> 3.3. Renewable energy	Ongoing	Minority (<50%)	Slow, significant declines
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation	
4. Transportation & service corridors -> 4.1. Roads & railroads	Ongoing	Whole (>90%)	Slow, significant declines
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion	
5. Biological resource use -> 5.1. Hunting & trapping terrestrial animals -> 5.1.1. Intentional use (species is the target)	Ongoing	Majority (50-90%)	Rapid declines
	Stresses:	2. Species Stresses -> 2.1. Species mortality	
5. Biological resource use -> 5.1. Hunting & trapping terrestrial animals -> 5.1.2. Unintentional effects (species is not the target)	Ongoing	Majority (50-90%)	Rapid declines
	Stresses:	2. Species Stresses -> 2.1. Species mortality	
5. Biological resource use -> 5.2. Gathering terrestrial plants -> 5.2.2. Unintentional effects (species is not the target)	Ongoing	Whole (>90%)	Slow, significant declines
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation	
5. Biological resource use -> 5.3. Logging & wood harvesting -> 5.3.3. Unintentional effects: (subsistence/small scale) [harvest]	Ongoing	Majority (50-90%)	Slow, significant declines
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation	
5. Biological resource use -> 5.3. Logging & wood harvesting -> 5.3.4. Unintentional effects: (large scale) [harvest]	Ongoing	Minority (<50%)	Slow, significant declines
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation	

7. Natural system modifications -> 7.2. Dams & water management/use -> 7.2.9. Small dams	Ongoing	Minority (<50%)	Slow, significant declines
Stresses: 1. Ecosystem stresses -> 1.1. Ecosystem conversion			
7. Natural system modifications -> 7.2. Dams & water management/use -> 7.2.10. Large dams	Ongoing	Minority (<50%)	Slow, significant declines
Stresses: 1. Ecosystem stresses -> 1.1. Ecosystem conversion			

Conservation Actions in Place

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Conservation Action in Place
In-place research and monitoring
Action Recovery Plan: No
In-place land/water protection
Area based regional management plan: No
Occurs in at least one protected area: Yes
Invasive species control or prevention: Not Applicable
In-place species management
Harvest management plan: No
Successfully reintroduced or introduced benignly: No
Subject to ex-situ conservation: No
In-place education
Subject to recent education and awareness programmes: No

Conservation Actions Needed

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Conservation Action Needed	Notes
2. Land/water management -> 2.1. Site/area management	-
5. Law & policy -> 5.1. Legislation -> 5.1.1. International level	-
5. Law & policy -> 5.1. Legislation -> 5.1.2. National level	-
5. Law & policy -> 5.1. Legislation -> 5.1.3. Sub-national level	-
5. Law & policy -> 5.2. Policies and regulations	-
5. Law & policy -> 5.4. Compliance and enforcement -> 5.4.1. International level	-
5. Law & policy -> 5.4. Compliance and enforcement -> 5.4.2. National level	-
5. Law & policy -> 5.4. Compliance and enforcement -> 5.4.3. Sub-national level	-

Research Needed

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Research Needed	Notes
1. Research -> 1.2. Population size, distribution & trends	-
1. Research -> 1.3. Life history & ecology	-
1. Research -> 1.4. Harvest, use & livelihoods	-
1. Research -> 1.5. Threats	-
1. Research -> 1.6. Actions	-
3. Monitoring -> 3.1. Population trends	-
3. Monitoring -> 3.2. Harvest level trends	-
3. Monitoring -> 3.3. Trade trends	-

Additional Data Fields

Distribution
Continuing decline in number of locations: Yes
Extreme fluctuations in the number of locations: No
Lower elevation limit (m): 0
Upper elevation limit (m): 2,300
Population
Continuing decline of mature individuals: Yes
Extreme fluctuations: No
Population severely fragmented: No
Continuing decline in subpopulations: Yes
Extreme fluctuations in subpopulations: No
All individuals in one subpopulation: No
Habitats and Ecology
Generation Length (years): 5
Movement patterns: Not a Migrant

Amendment

Amendment reason: This amended assessment was published to correct the name of an assessor from "Choudhury, A." to "Choudhury, A.U.".

The IUCN Red List Partnership



The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#).

The IUCN Red List Partners are: [ABQ BioPark](#); [Arizona State University](#); [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [Missouri Botanical Garden](#); [NatureServe](#); [Re:wild](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Senckenberg Society for Nature Research](#); [Texas A&M University](#); and [Zoological Society of London](#).