


Libellula angelina

Assessment by: Inoue, K.



View on www.iucnredlist.org

Citation: Inoue, K. 2006. *Libellula angelina*. The IUCN Red List of Threatened Species 2006: e.T11928A3315484. <http://dx.doi.org/10.2305/IUCN.UK.2006.RLTS.T11928A3315484.en>

Copyright: © 2015 International Union for Conservation of Nature and Natural Resources

Reproduction of this publication for educational or other non-commercial purposes is authorized without prior written permission from the copyright holder provided the source is fully acknowledged.

Reproduction of this publication for resale, reposting or other commercial purposes is prohibited without prior written permission from the copyright holder. For further details see [Terms of Use](#).

The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#). The IUCN Red List Partners are: [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [Microsoft](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); [Wildscreen](#); and [Zoological Society of London](#).

If you see any errors or have any questions or suggestions on what is shown in this document, please provide us with [feedback](#) so that we can correct or extend the information provided.

Taxonomy

Kingdom	Phylum	Class	Order	Family
Animalia	Arthropoda	Insecta	Odonata	Libellulidae

Taxon Name: *Libellula angelina* Selys, 1883

Assessment Information

Red List Category & Criteria: Critically Endangered A3ce [ver 3.1](#)

Year Published: 2006

Date Assessed: January 31, 2006

Justification:

Libellula angelina was recorded at 86 localities in 29 prefectures until the 1950s. It declined in the 1960s, with drastic declines in the 1990s. In 2000 it was found at only 18 localities in six prefectures. Developments have destroyed and degraded habitat and introduced predators threaten the species directly. These declines are expected to continue.

Previously Published Red List Assessments

1996 – Critically Endangered (CR)

1994 – Endangered (E)

1990 – Endangered (E)

1988 – Endangered (E)

1986 – Endangered (E)

Geographic Range

Range Description:

China (central and north), Japan (Honshu, Shikoku, Kyushu and offshore islands) and Korea.

Country Occurrence:

Native: China; Japan; Korea, Democratic People's Republic of; Korea, Republic of

Population

Current population size is estimated at < 5,000.

Current Population Trend: Decreasing

Habitat and Ecology (see Appendix for additional information)

Old and stable ponds with a moderate growth of emergent vegetation in lowland hill areas. An area of open water is also necessary.

Systems: Terrestrial, Freshwater

Threats (see Appendix for additional information)

Filling up of ponds to build houses and factories, and reforming to artificial ponds with concrete banks have much diminished the habitats. Destruction of grassy fields caused decrease of this species because the females and immature males need such fields as refuges. Introducing alien animals like *Procambarus clarkii* (Crustacea) and *Micropterus salmoides* (carnivorous fish) is a serious problem because of direct predation and habitat destruction.

Conservation Actions (see Appendix for additional information)

The Ministry of Environment started to protect this species in 1993, and collection is prohibited by law. But this is not enough for the protection of habitats from destruction. Some NGOs have begun to conserve this insect and its habitats, for example at Okegaya Pond, Shizuoka Pref., Imuta Pond, Kagoshima Pref. and Noyori-shin Pond, Oita Pref.

Credits

Assessor(s): Inoue, K.

Reviewer(s): Clausnitzer, V. & Suhling, F. (Odonata Red List Authority)

Bibliography

- Baillie, J. and Groombridge, B. (eds). 1996. *1996 IUCN Red List of Threatened Animals*. pp. 378. International Union for Conservation of Nature, Gland, Switzerland and Cambridge, UK.
- Fukui, M. 1987. Records of interspecific hybrid between *Libellula quadrimaculata asahinai* and *L. angelina*. *Tombo* 30(1–4): 36–43.
- Fukui, M. 1993. Observation on the larval growth of *Libellula angelina*. *Tombo* 36(1–4): 18
- Groombridge, B. (ed.). 1994. *1994 IUCN Red List of Threatened Animals*. IUCN, Gland, Switzerland and Cambridge, UK.
- Hamada, K and Inoue, K. 1983. *The dragonflies of Japan in colour*. Kodansha, Tokyo
- Inoue, K. 2004. Critical species of Odonata in Japan. *Journal of Odonatology* 7(2): 311–324.
- IUCN. 1990. *IUCN Red List of Threatened Animals*. IUCN, Gland, Switzerland and Cambridge, UK.
- IUCN. 2006. *2006 IUCN Red List of Threatened Species*. www.iucnredlist.org. Downloaded on 04 May 2006.
- Okudaira, M., Sugimura, M., Ishida, S., Kojima, K., Ishida, K. and Aoki, T. 2001. *Dragonflies of the Japanese Archipelago in Colour — The Optimum Pictorial Guidebook to the Odonata of Japan*. Hokkaido Uni. Press., Sapporo, Japan.

Citation

Inoue, K. 2006. *Libellula angelina*. *The IUCN Red List of Threatened Species 2006*: e.T11928A3315484. <http://dx.doi.org/10.2305/IUCN.UK.2006.RLTS.T11928A3315484.en>

Disclaimer

To make use of this information, please check the [Terms of Use](#).

External Resources

For [Images and External Links to Additional Information](#), please see the Red List website.

Appendix

Habitats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Habitat	Season	Suitability	Major Importance?
5. Wetlands (inland) -> 5.6. Wetlands (inland) - Seasonal/Intermittent Freshwater Lakes (over 8ha)	-	Suitable	-
5. Wetlands (inland) -> 5.8. Wetlands (inland) - Seasonal/Intermittent Freshwater Marshes/Pools (under 8ha)	-	Suitable	-

Threats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Threat	Timing	Scope	Severity	Impact Score
1. Residential & commercial development -> 1.1. Housing & urban areas	Ongoing	-	-	-
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation		
1. Residential & commercial development -> 1.2. Commercial & industrial areas	Ongoing	-	-	-
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation		
8. Invasive & other problematic species & genes -> 8.1. Invasive non-native/alien species -> 8.1.1. Unspecified species	Ongoing	-	-	-
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality		
8. Invasive & other problematic species & genes -> 8.1. Invasive non-native/alien species -> 8.1.2. Named species (<i>Procambarus clarkii</i>)	Ongoing	-	-	-
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality		
8. Invasive & other problematic species & genes -> 8.1. Invasive non-native/alien species -> 8.1.2. Named species (<i>Micropterus salmoides</i>)	Ongoing	-	-	-
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality		

Conservation Actions Needed

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Conservation Actions Needed
2. Land/water management -> 2.1. Site/area management

Conservation Actions Needed
2. Land/water management -> 2.3. Habitat & natural process restoration

The IUCN Red List Partnership



The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission](#) (SSC) and [The IUCN Red List Partnership](#). The IUCN Red List Partners are: [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [Microsoft](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); [Wildscreen](#); and [Zoological Society of London](#).