INS-001

Screening versus diagnosis?

INS-002

PHQ-9: structure and severity bands?

INS-003

PHQ-9 item 9 is positive (thoughts of death or self-harm). Next step?

INS-004

PHQ-2?

INS-005

GAD-7: structure and bands?

INS-006

Geriatric Depression Scale (short form)?

INS-007

Edinburgh Postnatal Depression Scale (EPDS)?

INS-008

EPDS item 10 is positive (thoughts of self-harm)?

INS-002

9 items over the past 2 weeks, scored 0-27. Scores of 5, 10, 15 and 20 mark mild, moderate, moderately severe and severe depression.

Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2001). The PHQ-9. Journal...

INS-001

A positive screen means 'look closer,' not a diagnosis. Follow it with a clinical assessment, and consider language, culture and literacy.

DSM-5-TR (APA, 2022) Section III: assessment measures

INS-004

The first two PHQ items (low mood and loss of interest), scored 0-6. A score of 3 or more calls for the full PHQ-9 or further assessment.

Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2003). The Patient Health...

INS-003

A suicide risk assessment right away, whatever the total score.

Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2001). The PHQ-9. Journal...

INS-006

15 yes/no items, easy for older adults to answer. A score above 5 suggests depression that warrants further evaluation.

Sheikh, J. I., & Yesavage, J. A. (1986). Geriatric Depression Scale (GDS): R...

INS-005

7 items over 2 weeks, scored 0-21. Scores of 5, 10 and 15 mark mild, moderate and severe anxiety; 10 is a common cut point.

Spitzer, R. L., Kroenke, K., Williams, J. B. W., & Lowe, B. (2006). A brief m...

INS-008

Assess safety right away, regardless of the total score.

Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal de...

INS-007

10 items, scored 0-30, for perinatal depression. Cox and colleagues used 12/13 for probable depression; lower cut points are used for screening.

Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal de...

INS-009

Beck Hopelessness Scale: why it matters?

INS-010

SCARED (child anxiety)?

INS-011

Columbia Suicide Severity Rating Scale (C-SSRS)?

INS-012

Ask Suicide-Screening Questions (ASQ)?

INS-013

SAD PERSONS scale: use it alone to predict suicide?

INS-014

Safety planning intervention (Stanley and Brown)?

INS-015

AUDIT: structure and key cut point?

INS-016

AUDIT risk zones?

INS-010

41 items with child and parent forms. A total of 25 or more may indicate an anxiety disorder; subscales cover panic, generalized, separation, social and school avoidance.

Birmaher, B., et al. (1999). Psychometric properties of the Screen for Chil...

INS-009

Hopelessness is a stronger predictor of suicide than depression severity alone.

Beck, A. T., Weissman, A., Lester, D., & Trexler, L. (1974). The measurem...

INS-012

Four yes/no questions for youth and adults in medical settings, plus a question about current thoughts. Any yes is positive; yes to current thoughts is acute and needs immediate evaluation.

Horowitz, L. M., et al. (2012). Ask Suicide-Screening Questions (ASQ). Arc...

INS-011

Structured questions about the wish to be dead, suicidal thoughts, method, intent, plan and recent suicidal behavior. Intent, a plan or recent behavior signal higher risk.

Posner, K., et al. (2011). The Columbia-Suicide Severity Rating Scale. Am...

INS-014

A written, collaborative plan: warning signs, coping strategies, people and places for distraction, people to ask for help, professional contacts, and making the environment safer.

Stanley, B., & Brown, G. K. (2012). Safety planning intervention. Cognitiv...

INS-013

No. It is an old mnemonic of risk factors with poor predictive accuracy; a full clinical risk assessment is needed.

Patterson, W. M., Dohn, H. H., Bird, J., & Patterson, G. A. (1983). Evaluati...

INS-016

0-7 low risk (education), 8-15 (simple advice), 16-19 (brief counseling and monitoring), 20-40 (refer for diagnostic evaluation and treatment).

Babor, T. F., Higgins-Biddle, J. C., Saunders, J. B., & Monteiro, M. G. (2001...

INS-015

WHO's 10-item alcohol screen, scored 0-40. A score of 8 or more indicates hazardous or harmful drinking.

Babor, T. F., Higgins-Biddle, J. C., Saunders, J. B., & Monteiro, M. G. (2001...

INS-017

AUDIT-C?

INS-018

CAGE?

INS-019

CRAFFT?

INS-020

DAST-10?

INS-021

SBIRT?

INS-022

CIWA-Ar?

INS-023

COWS?

INS-024

ASAM Criteria: what are they for?

INS-018

Four yes/no questions: Cut down, Annoyed by criticism, Guilty, Eye-opener. Two or more yes answers is clinically significant. Less sensitive for at-risk drinking.

Ewing, J. A. (1984). Detecting alcoholism: The CAGE questionnaire. JAMA,...

INS-017

The 3 consumption items of the AUDIT, scored 0-12. Commonly positive at 4 or more for men and 3 or more for women.

Bradley, K. A., et al. (2007). AUDIT-C as a brief screen for alcohol misuse...

INS-020

10 yes/no items screening drug use other than alcohol over the past year; higher scores mean more severe problems.

Skinner, H. A. (1982). The Drug Abuse Screening Test. Addictive Behavio...

INS-019

Adolescent substance screen (Car, Relax, Alone, Forget, Friends, Trouble). A score of 2 or more indicates high risk and need for further assessment.

Knight, J. R., et al. (2002). Validity of the CRAFFT substance abuse screen...

INS-022

A 10-item clinician scale (maximum 67) that rates alcohol withdrawal severity and guides symptom-triggered medical treatment.

Sullivan, J. T., et al. (1989). Assessment of alcohol withdrawal: The revise...

INS-021

Screening, Brief Intervention and Referral to Treatment: universal screening, a short motivational conversation, and referral when risk is high.

Babor, T. F., et al. (2007). Screening, Brief Intervention, and Referral to T...

INS-024

A multidimensional framework for matching substance use treatment to the right level of care; the least intensive safe and effective level is preferred.

ASAM Criteria (4th ed., 2023) dimensions and levels of care

INS-023

An 11-item clinician scale for opioid withdrawal: 5-12 mild, 13-24 moderate, 25-36 moderately severe, above 36 severe.

Wesson, D. R., & Ling, W. (2003). The Clinical Opiate Withdrawal Scale (C...

INS-025

PCL-5?

INS-026

PC-PTSD-5?

INS-027

ACE questionnaire?

INS-028

MoCA?

INS-029

MMSE versus MoCA?

INS-030

Mini-Cog?

INS-031

Confusion Assessment Method (CAM)?

INS-032

Katz ADLs versus Lawton IADLs?

INS-026

5 yes/no items for primary care, asked after a trauma exposure question. A cut point of 4 is recommended; lower may be considered for women.

VA National Center for PTSD. Primary Care PTSD Screen for DSM-5 (PC-P...

INS-025

20-item self-report of DSM-5 PTSD symptoms, scored 0-80. Scores around 31-33 suggest probable PTSD; a 10-point drop suggests meaningful response.

VA National Center for PTSD. PTSD Checklist for DSM-5 (PCL-5) scoring

INS-028

30-point cognitive screen sensitive to mild impairment; covers executive function, memory, attention and more. 26 or above is considered normal; add 1 point for 12 or fewer years of education.

Nasreddine, Z. S., et al. (2005). The Montreal Cognitive Assessment, MoC...

INS-027

10 categories of childhood abuse, neglect and household dysfunction. Higher scores show population-level health risk, not an individual diagnosis.

Felitti, V. J., et al. (1998). Relationship of childhood abuse and household...

INS-030

A quick cognitive screen: three-word recall plus clock drawing.

Borson, S., et al. (2000). The Mini-Cog. International Journal of Geriatric P...

INS-029

Both 30-point screens. The MMSE is less sensitive to mild impairment and executive problems, and scores are affected by education and language.

Nasreddine, Z. S., et al. (2005). The Montreal Cognitive Assessment, MoC...

INS-032

ADLs: bathing, dressing, toileting, transferring, continence, feeding. IADLs: phone, shopping, meals, housekeeping, laundry, transport, medications, finances.

Lawton, M. P., & Brody, E. M. (1969). Assessment of older people: Self-m...

INS-031

Delirium screen: acute onset and fluctuating course, plus inattention, plus either disorganized thinking or altered level of consciousness.

Inouye, S. K., et al. (1990). Clarifying confusion: The Confusion Assessme...

INS-033

Which tasks usually decline first in neurocognitive disorder: ADLs or IADLs?

INS-034

Zarit Burden Interview?

INS-035

M-CHAT-R/F: age and scoring?

INS-036

Vanderbilt and Conners scales: why both parent and teacher forms?

INS-037

Strengths and Difficulties Questionnaire (SDQ)?

INS-038

HITS?

INS-039

Danger Assessment (Campbell)?

INS-040

Genogram versus ecomap?

INS-034

Measures caregiver burden, for example in family members caring for someone with dementia.

Zarit, S. H., Reever, K. E., & Bach-Peterson, J. (1980). Relatives of the im...

INS-033

IADLs, such as managing money and medications; basic ADLs decline later.

DSM-5-TR (APA, 2022) Neurocognitive Disorders: Major and Mild Neuroco...

INS-036

ADHD requires symptoms in two or more settings, so multi-informant ratings are needed.

Wolraich, M. L., et al. (2003). Psychometric properties of the Vanderbilt A...

INS-035

Parent-report autism screen for 16-30 months. 0-2 low risk; 3-7 medium (give the follow-up interview); 8-20 high (refer for evaluation and early intervention).

Robins, D. L., et al. (2014). Validation of the Modified Checklist for Autis...

INS-038

Brief intimate partner violence screen: how often a partner Hurts, Insults, Threatens or Screams at the person.

Sherin, K. M., et al. (1998). HITS: A short domestic violence screening too...

INS-037

A brief behavioral screen for ages about 4-16 covering emotional symptoms, conduct, hyperactivity, peer problems and prosocial behavior.

Goodman, R. (1997). The Strengths and Difficulties Questionnaire. Journa...

INS-040

Genogram: three or more generations of family patterns. Ecomap: the family's current links to outside systems and resources.

Hartman, A. (1978). Diagrammatic assessment of family relationships. S...

INS-039

Rates risk of intimate partner homicide. Major red flags include past strangulation, a gun in the home, recent or planned separation and escalating violence.

Campbell, J. C., Webster, D. W., & Glass, N. (2009). The Danger Assessm...

INS-041

Culturagram?

INS-042

Cultural Formulation Interview?

INS-043

WHODAS 2.0?

INS-044

DSM-5 Level 1 Cross-Cutting Symptom Measure?

INS-045

Mental status exam: mood versus affect?

INS-046

MSE: tangential versus circumstantial thinking?

INS-047

MSE: illusion versus hallucination?

INS-048

MSE: insight versus judgment?

INS-042

DSM-5-TR Section III interview (16 core questions) for any client, exploring the cultural definition, causes and help-seeking around the problem.

DSM-5-TR (APA, 2022) Section III: Cultural Formulation

INS-041

Assesses immigrant families: reasons for migrating, time in the community, legal status, language, health beliefs, holidays, crises, and values about family and work.

Congress, E. P. (1994). The use of culturagrams to assess and empower...

INS-044

A brief screen across many symptom domains; positive domains lead to more detailed Level 2 measures.

DSM-5-TR (APA, 2022) Section III: assessment measures

INS-043

DSM-5-TR Section III measure of disability across six domains: cognition, mobility, self-care, getting along, life activities, participation.

DSM-5-TR (APA, 2022) Section III: assessment measures

INS-046

Circumstantial: wanders through details but reaches the point. Tangential: drifts away and never returns to it.

Boland, Verduin & Ruiz (Eds.). Kaplan & Sadock's Synopsis of Psychiatry...

INS-045

Mood is the client's own report ('I feel empty'). Affect is what the worker observes (range, congruence, appropriateness).

Boland, Verduin & Ruiz (Eds.). Kaplan & Sadock's Synopsis of Psychiatry...

INS-048

Insight: awareness of having a problem and needing help. Judgment: ability to make sound decisions, best judged from real recent choices.

Boland, Verduin & Ruiz (Eds.). Kaplan & Sadock's Synopsis of Psychiatry...

INS-047

Illusion: misperceiving a real stimulus. Hallucination: perceiving something with no stimulus present.

Boland, Verduin & Ruiz (Eds.). Kaplan & Sadock's Synopsis of Psychiatry...