

Table 1: Comparison of clustering inertia (left) and CPU time (right) among KSKM, KSKM_E, and PCCC with no step of centroid mutation, reported as multiples of the baseline method (KSKM) across datasets. Lower inertia and smaller time multiples indicate better performance.

Dataset	Stats Method Constraint Level	Inertia			CPU Time		
		KSKM	KSKM_E	PCCC	KSKM	KSKM_E	PCCC
COL1	0.05	1.00	1.00	1.00	1.00	4.09	25.84
	0.1	1.00	1.00	1.00	1.00	5.84	27.47
	0.15	1.00	1.00	0.99	1.00	6.74	25.80
	0.2	1.00	1.00	1.00	1.00	7.96	37.80
COL2	0.05	1.00	1.00	0.99	1.00	7.57	81.68
	0.1	1.00	0.98	0.98	1.00	37.19	180.17
	0.15	1.00	0.96	0.96	1.00	59.67	286.06
	0.2	1.00	0.96	0.97	1.00	83.40	338.96
	0.25	1.00	0.95	0.97	1.00	105.16	412.30
	0.3	1.00	0.94	0.97	1.00	108.52	450.33
	0.35	1.00	0.99	0.99	1.00	101.87	372.77
	0.4	1.00	0.99	0.98	1.00	111.36	368.39
	0.45	1.00	0.98	0.97	1.00	103.70	438.68
	0.5	1.00	0.99	1.00	1.00	50.52	205.19
COL3	0.05	1.00	0.98	0.98	1.00	48.23	412.70
	0.1	1.00	0.96	0.96	1.00	329.98	1162.69
	0.15	1.00	0.96	0.96	1.00	485.59	2245.15
	0.2	1.00	0.98	0.98	1.00	425.59	1085.59