

ST437/537 – HW #01

Arnab Maity

Due date: January 17, 2019

Instructions

Please follow the instructions below when you prepare and submit your assignment.

- **Include a cover-page** with your homework. It should contain
 - i. Full name,
 - ii. Course#: ST 437/537 and
 - iii. HW-#
 - iv. Submission date
- Assignments should be submitted in class on the date specified (“due date”).
- Neatly typed or hand-written solution on standard letter-size papers (stapled on the top-left corner) should be submitted. **All R code/output should be well commented, with relevant outputs highlighted.**
- **Always staple (upper left corner) your homework before coming to class. Ten percent points will be deducted otherwise.**
- When you solve a particular problem, do not only give the final answer. Instead **show all your work** and the steps you used (with proper explanation) to arrive at your answer to get full credit.

Problems

Solve the following problems. You may use R for these problems unless I specifically instruct otherwise.

1. (10 points) Let $x = (5, 1, 3)^T$ and $y = (-1, 3, 1)^T$ be two 3×1 column vectors.

- a. Find the length of x and y (Do this by hand)
- b. Find $x^T y$ (Do this by hand)
- c. Are x and y orthogonal? Explain.
- d. Repeat (a) and (b) using R. Provide code and output.

2. (10 points) Read the section about linear combinations in the “Multivariate summary statistics” lecture notes, and then answer the following questions.

You are given a random vector $X = [X_1, X_2, X_3, X_4]^T$ with the mean vector $\mu = [4, 3, 2, 1]^T$, and the variance-covariance matrix

$$\Sigma = \begin{bmatrix} 3 & 0 & 2 & 2 \\ 0 & 1 & 1 & 0 \\ 2 & 1 & 9 & -2 \\ 2 & 0 & -2 & 4 \end{bmatrix}.$$

Also define the matrix

$$A = \begin{bmatrix} 1 & 2 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ 0 & 0 & 1 & -2 \\ 0 & 0 & 2 & -1 \end{bmatrix}.$$

- Find $E(X)$ and $E(AX)$.
- Find $cov(AX)$.

3. (20 points) Consider the `iris` data available in `R`.

- Construct side-by-side boxplots of the four quantitative variables (SL, SW, PL, PW) for each species. Do not forget to properly label the axes and give a proper title to the plots when needed. [Hint: you will have 3 plots, one for each species. Each plot will contain four boxplots. See the function `boxplot()` in `R`]
- Construct a `pairs-plot` of the four variables *for each of the three species*.
- Define the vector $\mathbf{x} = [\text{Sepal. Length}, \text{Sepal. Width}, \text{Petal. Length}, \text{Petal. Width}]^T$. The dataset have 50 observations of this vector (one for each flower), and with 3 species. Thus, we have a sample $\mathbf{x}_1, \dots, \mathbf{x}_n$ of size $n = 50$ for each of the three species. Compute the sample mean $\bar{\mathbf{x}}$ (a 4×1 vector), the sample covariance matrix \mathbf{S} (a 4×4 matrix) and the sample correlation matrix \mathbf{R} (a 4×4 matrix) *for each species*.
- Looking at the pairs plot in (b) and the correlation matrices in (c), do you see any patterns or differences among the species? Explain.

4. (20 points) Consider the `skulls` dataset in the `HSAUR3` package in `R`. You will first need to install the package in `R` to access the dataset. Use `?skulls` command to get more details on the data. A snapshot of the data is shown below.

```
library(HSAUR3)
```

```
## Loading required package: tools
```

```
head(skulls)
```

```
##      epoch  mb  bh  bl  nh
## 1 c4000BC 131 138  89  49
## 2 c4000BC 125 131  92  48
## 3 c4000BC 131 132  99  50
## 4 c4000BC 119 132  96  44
## 5 c4000BC 136 143 100  54
## 6 c4000BC 138 137  89  56
```

- Suppose we want to estimate the *population mean* of all the 4 variable for skulls with epoch `c4000BC`. Write down the population, parameter, the sample and the statistic you will use to answer the question above.
- From the `skulls` data set, provide an estimate (numeric value) of the parameter mentioned above. Explain how you obtained it.

- c. Now suppose we want to estimate the population variance-covariance matrix. When estimator will you use? Provide a numeric estimate.
- d. Compute the variance covariance matrix of the estimator of the population mean in part (b).
- e. Provide an estimate for the parameter vector $(\mu_{mb} - \mu_{nh}, \mu_{bh} - \mu_{nh})^T$ and compute the covariance matrix of the estimator.

5. (10 points) Answer the following questions.

- a. What is the basic difference between multivariate and longitudinal data?
- b. Why is investigating covariance (or correlation) between two variables important?
- c. Suppose that in your analysis, you found $cor(X, Y) = 0$. Can you say that “ X and Y does not have *any* relationship at all?” Explain.
- d. In problem 4 above, you considered the `skulls` dataset. Is this a longitudinal dataset (since the skulls are coming from different era/time as given in the epoch)? Explain.