

Please share this with a doctor or pharmacist.

Dear Healthcare Professional,

I am interested in my heart health and would like to know if Crestor as a non-prescription cholesterol treatment is right for me. I have completed the self-assessment which utilizes a web app to assess my eligibility for treatment based on an AHA/ACC cardiovascular risk calculation and the proposed Drug Facts Label for "Crestor OTC". The results indicate that I qualify for cholesterol-lowering therapy, but that I should "talk to a doctor or pharmacist" about the shaded responses in the attached summary to determine if Crestor OTC is right for me.

Questions	Response
How old are you?	28
What was your sex at birth?	Male
Which best describes your ethnicity/race?	White
Are you taking any cholesterol or triglyceride lowering prescription medicine?	No
Are you taking cyclosporine (a medicine for your immune system)?	No
Are you taking warfarin (a blood thinner)?	No
Have you ever had a heart attack?	No
Have you ever had a stroke?	No
Have you ever had an operation or procedure on your heart?	No
Have you ever had Peripheral Artery Disease (PAD)?	No
Has your father or brother had a heart attack or stroke <u>before</u> age 55?	Yes
Has your mother or sister had a heart attack or stroke <u>before</u> age 65?	No
What are your triglycerides?	140
What is your total cholesterol level?	200
What is your LDL cholesterol?	160
What is your HDL cholesterol?	39
Do you have liver disease?	No
Have you been told by a doctor that you have kidney disease?	No
Do you drink 3 or more glasses of alcohol <u>every day</u> ?	No
Have you had a previous side effect to any medicine to treat your cholesterol and a doctor stopped the medicine?	No
Are you taking any of the following medicines?	Yes

- Colchicine (a medicine to treat gout)
- HIV/AIDS or hepatitis medicines (such as lopinavir, ritonavir or atazanavir)

These medicines can raise blood levels of Crestor OTC and/or increase the risk of side effects.

Questions? Go to <website> or call toll-free 1-844-856-1864

The results of the Crestor OTC assessment requiring the consumer/patient to ask a doctor or pharmacist whether Crestor OTC is right for them is intended to inform clinical judgement. The appropriateness of Crestor OTC should be determined after discussion between the consumer/patient and their health care provider.

Drug Facts	
Active ingredient (in each tablet)	Purpose
Rosuvastatin calcium 5 mg	Lower cholesterol
Use	
<ul style="list-style-type: none"> To lower cholesterol, one of the key risk factors that can lead to heart disease, if you: <ol style="list-style-type: none"> have a LDL cholesterol of 70 to 189 mg/dL AND are female 50 to 75 years of age or are male 20 to 75 years of age AND go to <website> and complete the assessment to see if this product is right for you 	
Warnings	
<p>Allergy alert: Do not use if you ever had an allergic reaction to this product or any of the ingredients.</p> <p>Pregnancy alert: Do not use if you are pregnant or breastfeeding. If you become pregnant, stop taking and call a doctor.</p>	
Do not use if you:	
<ul style="list-style-type: none"> had a heart attack, stroke, peripheral artery disease (PAD) or an operation or procedure on your heart have liver disease are taking any of the following <ul style="list-style-type: none"> cholesterol lowering or triglyceride lowering prescription medicines cyclosporine (a medicine for your immune system) warfarin (blood thinner) 	
Ask a doctor before use if you:	
<ul style="list-style-type: none"> had a previous side effect to any medicine to treat your cholesterol have kidney disease drink 3 or more glasses of alcohol every day are taking <ul style="list-style-type: none"> colchicine (a medicine to treat gout) HIV/AIDS or hepatitis medicines (such as lopinavir, ritonavir or atazanavir) 	
When using this product:	
<ul style="list-style-type: none"> temporary mild body aches, headache, abdominal pain, or nausea may occur. Continue taking this product 	
Stop use and ask a doctor if you:	
<ul style="list-style-type: none"> get severe and unexplained muscle pain, tenderness or weakness get symptoms of liver problems (upper belly pain, dark urine, yellowing of skin or whites of eyes) 	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions	
<ul style="list-style-type: none"> Take 1 tablet every day <ul style="list-style-type: none"> swallow the tablet whole, do not crush do not take antacids within 2 hours of taking this product continue taking every day or your cholesterol number will go back up eat a healthy diet and exercise Test your LDL cholesterol within 3 months after you start taking this product and then once a year <ul style="list-style-type: none"> if your LDL cholesterol number did not go down, see a doctor 	
Male Age	Female Age
under 20	under 50
20-75	50-75
76 and over	
Do not use	
Take 1 tablet every day	
Do not start taking If already taking, continue to take 1 tablet every day.	
Other information store at 68 to 77°F (20 to 25°C)	
Inactive ingredients microcrystalline cellulose NF, lactose monohydrate NF, tribasic calcium phosphate NF, crospovidone NF, magnesium stearate NF, hypromellose NF, triacetin NF, titanium dioxide USP, yellow ferric oxide, and red ferric oxide NF	
Questions? Call toll-free 1-844-856-1864	