

Plotting Wavefields from Research Module

Salome Bachmann

March 24, 2026



What have I been working on?


- Writing introduction
- Implementing feedback for introduction
- Setup of remote machine
- Running simulations with lower source frequency
- Plotting waterfall plots, wiggle plots, single traces and frequency content

Simulation parameters used

- 30m by 3m domain
- one simulation with 150m by 3m domain
- 3 layers, all layers 1m thick
- absorbing boundaries at edges of domain
- 10Hz vector point source
- moment tensor source for skier ($m_{yy}=0$, $m_{xx}=0$, $m_{xy}=3e4$)


Wavefield Waterfall plots

Seismic waterfall plot below source location at 1m depth in snow



(a) Vector point 2D source with $f_y = -1$. Source location indicated by white line. Colorbar shows particle velocity (m/s).

Seismic waterfall plot below source location at 1m depth in snow




(b) Moment tensor source with $m_{xx} = 0$, $m_{yy} = 0$, $m_{xy} = 3 \times 10^4$. Source location indicated by white line. Colorbar shows particle velocity (m/s).


Waterfall plots interpretation

- Both sources show horizontal bands with high velocity: possible resonance between snow and air layer?
- Explained by tuning (Widess, 1973): possible constructive interference between top and bottom reflections of layer
- Vector point source shows some direct wave moveout (v-shaped pattern)
- Vector point source also shows some reflections at later times
- Problem of resonance could be solved by larger domain?

Wiggle Plots




(a) Wiggle plot for vector point source.




(b) Wiggle plot for moment tensor source.

Bandpass filtered wiggle plots



(a) Bandpass filtered wiggle plot for vector point source, Bandpass from 1 to 30HZ




(b) Bandpass filtered wiggle plot for moment tensor source, Bandpass from 5-20HZ

Wiggle Plots Interpretation

- Very noisy traces for vector point source, especially at the edges to the domain
- Bandpass filtering: Both moment tensor and vector point source show clear first breaks (large wiggles)
- Some ground roll maybe (later arrivals)
- No hyperbolic reflections, which makes sense from domain setup

Single traces & Spectrograms



Interpretation of single traces and spectrogram, power spectral density content

- Broadband signal after firing of source (indicated by black line)
- SOurce frequency shown by cyan lines in spectrogram
- Energy at all frequencies: field close to the source not dominated by source frequency




Figure: Single trace, power spectral density and spectrogram for moment tensor source

Interpretation of single traces and spectrogram, power spectral density content



Problems / Questions

- Simulations take very long to run on remote machine (2+ hours)
- Does it make sense to add topography to simulations?
- Does it make sense to add eg old snow layer (make the simulations multilayered)?

References



Widess, M. B. (1973). How thin is a thin bed? *Geophysics*, 38(6), 1176–1180.
<https://doi.org/10.1190/1.1440403> 



Salome Bachmann
Master Student
sbachmann@student.ethz.ch

ETH Zurich
D-EAPS