Rodenticide Tables

2022-08-31

Table 1: N compounds (with trace)

| | coef | param_est | std_error | 2.5CI | 97.5CI | P-value |
|----|------------------------------|-----------|--------------|-------|--------|---------|
| 2 | $Threshold0_1$ | -2.69 | 0.49 | -3.65 | -1.73 | 0.00 |
| 3 | $Threshold1_2$ | -1.35 | 0.47 | -2.27 | -0.44 | 0.00 |
| 4 | $Threshold2_3+$ | 0.52 | 0.46 | -0.38 | 1.42 | 0.00 |
| 5 | SexM | 0.69 | 0.41 | -0.12 | 1.50 | 0.00 |
| 6 | Juvenile | -1.78 | 0.34 | -2.45 | -1.12 | 0.00 |
| 7 | Subadult | -0.22 | 0.32 | -0.86 | 0.41 | 0.00 |
| 11 | $SexM \times Juv$ | -0.21 | 0.55 | -1.28 | 0.86 | 0.00 |
| 12 | $SexM \times SubAd$ | -0.45 | 0.56 | -1.55 | 0.65 | 0.00 |
| 8 | Pasture (30km2) | -0.04 | 0.12 | -0.29 | 0.20 | 0.04 |
| 9 | Intermix (100m, 30km2) | 0.43 | 0.37 | -0.30 | 1.15 | 0.00 |
| 10 | Lagged BMI $(30 \text{km}2)$ | 0.13 | 0.22 | -0.30 | 0.57 | 0.00 |