

Characteristics		Number of patients (all)	Percentage of all patients (%)	Number of patients (Sub-cohort for which the microbiota was characterized)	Percentage of sub-cohort (%)
Transplant recipients		37	100	30*	81.1
Average recipient age in years		8.2 (Range: 1.1-18.0)	NA	7.8 (Range: 1.1-16.5)	NA
Intestinal microbiome characterized	At 1-2 timepoints	NA	NA	8	26.7
	At 3-4 timepoints	NA	NA	15	50
	At 5-6 timepoints	NA	NA	8	26.7
Patient sex	Female	15	40.5	12	40
	Male	22	59.5	18	60
Disease at transplantation	malignant hematologic diseases	23	62.2	17	56.7
	Severe aplastic anemia	5	13.5	5	16.7
	Other benign disorders including immunodeficiencies	9	24.3	8	26.7
Donor type	HLA-matched sibling	7	18.9	6	20
	HLA-matched unrelated donor (9/10 or 10/10 match)	27	73	22	73.3
	HLA-mismatched umbilical cord blood donor (8/10 match)	3	8.1	2	6.7
Stem cell source	Bone marrow	30	81.1	25	83.3
	Umbilical cord blood	4	10.8	3	10
	Peripheral blood	2	5.4	1	3.3
	Bone marrow and umbilical cord blood	1	2.7	1	3.3
Conditioning regimen	TBI + CY / TBI + VP16	10	27.1	10	33.3
	Combinations of BU, CY, VP16 and MEL	18	48.7	12	40
	Fludarabine-based conditioning	9	24.3	8	26.7

Anti-thymocyte globulin treatment		28	75.7	23	76.7
Antibiotics pre- and post-HSCT		37	100	30	100
Sex mismatch (female donor to male recipient)		4	10	3	10
Acute GvHD	Grade 0-I	26	70.3	22	73.3
	Grade II-IV	11	29.2	8	26.7
Chronic GvHD within 24 months		8	18.9	5	16.7
Bacterial infections	At least one registered bacterial infection	25	67.6	19	63.3
	No registered bacterial infection	12	32.4	11	36.7
Overall Survival**	Alive	30	81.1	24	80
	Dead	7	18.9	6	20
Relapse of primary malignancy		6	16.2	7	23.3
Non-relapse mortality		2	5.4	1	3.3
Re-transplantation		3	8.1	4	13.3

*29/37 +1, one patient's microbiome was characterized but for which hBD2 was not measured

** Mean follow-up time after HSCT: 62.1 months (range: 48.9 – 75.8 months)