Koen Essers • Master Internship • October 12, 2026


Notes Week 32

a. Grid

Based on the simple binary evolution grids I showed last week, I have decided to simulate a high dimensional grid that looks at a broad mass range, broad initial mass-ratio range, and initial Roche lobe radii such that I simulate pretty much the complete RLOF regime with some dredge up.

I do this for two RLOF accretion efficiencies, each with three different ratios for isotropic re-emission and circumbinary mass loss. The ratios are (CCT, isotropic re-emission) = (0.1, 0.9) ,(0.2, 0.8) ,(0.3, 0.7). The accretion efficiencies are 0 and 0.1.

The plot below shows the complete grid.



Notice in naive simple binary case, the different ratios between isotropic re-emission and mass loss to a circumbinary torus would result in the same accretion and thus in the same abundances. Therefore, each dot is really 3 dots stacked on top of each other.


In the detailed MESA simulation, there will most likely be small differences between the final abundances for these systems. However, the biggest differences between these will be in the final orbital properties of these systems. More mass loss to a circumbinary torus will generally lead to much tighter final binary systems.

The panels with $\varepsilon_{\mathrm{RLOF}} = 0.1$ and $\varepsilon_{\mathrm{RLOF}} = 0.0$ at first sight might look very similar. This is because in both these cases, *most* of the mass is transferred via winds, which are not affected by the efficiency parameter. However, in the regime where winds are not very strong, differences between the two settings can be seen. This is on the left side of the plot. Here we see that the systems with *some* RLOF accretion are generally able to create barium stars with slightly smaller initial Roche lobe radii.

b. Mean Molecular Weight


b.1 Giant Branch Dip

Using the `EOS_plotter` routine from *MESA*, it is possible to clearly plot the different EOS tables that MESA uses, and in which regions in the ρ-T diagram. Below, I show this, and I plot the ρ-T profile for an evolved RGB star over these tables.



The RGB profile is colored, where very red regions correspond to regions with relatively large mean molecular weight μ. Blue regions correspond to the lowest mean molecular weights, and white corresponds to an average mean molecular weight of the envelope. I think this figure quite clearly shows why we see a pit in the mean molecular weight. The core boundary happens inside the Skye EoS table (Jermyn et al., 2021). This table *assumes* complete ionization. Once the profile crosses over to the boundary region, we see that the mean molecular weight increases until it crosses the FreeEOS regime, which does *not assume* complete ionization.

Now that the regions of the different equations of state tables are clear, and we know that our giant branch models cross into the Skye EOS just before the core, we can again make a plot of the μ as a function of depth in the star. For the sake of clarity, I chose $\log R$ for the horizontal axis, but obviously one could chose various different quantities such as internal mass, temperature or pressure.




This plot *clearly* shows that the pit in the μ profile is not real. The Skye EOS (Jermyn et al., 2021) *assumes* complete ionization, which, according to the FreeEOS, is clearly not true up until the deepest parts of its influence. The cyan blending region just ensures the two different tables do not produce discontinuous profiles.

We also clearly see that my naive complete ionization prescription to compute the mean molecular weight perfectly matches the Skye (Jermyn et al., 2021) EOS, and starts to diverge in the blending region, just as I would expect based on the fact that Skye assumes full ionization and FreeEOS does not.


b.2 Main Sequence

Now, let's look at the main sequence. Maybe here we see a similar crossing between EOS regimes.



I once again plot the ρ-T plane and the regions of the different EOS tables. I then plot 3 different MS models with different masses and $X(\mathrm{H})_{\mathrm{center}} = 0.65$. I also indicate where the hydrogen burning core boundary is with a black dot.

The plot clearly shows that the heaviest model here crosses into the Skye EOS (Jermyn et al., 2021) in its core. The entirety of the 1 $M_\odot$ model uses the FreeEOS table and the 0.4 $M_\odot$ crosses into the OPAL/SCVH region (Rogers & Nayfonov, 2002).




Again, we see that the mean molecular weight of the Skye EOS agrees perfectly with my fully ionized computation. As soon as we enter the blended region, the mean molecular weight starts to deviate.

The weird thing is that the Asplund et al. (2009) mean molecular weight agrees with the FreeEOS nearly perfectly.

It took me an embarrassingly long time, but I *think* I have finally been able to figure out what causes the differences in μ between the Skye EOS and FreeEOS. The differences can be explained by the masses of the elements that are used in the computation of μ. My naive approach for complete ionization for the *MESA* abundances was to simply take integer mass numbers equal to the number of protons plus neutrons. *However*, for the Asplund et al. (2009) mean molecular weight computations, I use the Standard atomic weight. This is a bit different.

It seems like Skye (Jermyn et al., 2021) uses the mass number, while the FreeEOS uses the atomic mass. In fact, when I compute the *MESA* mean molecular weight using both methods, we can clearly see that the computed mean molecular weights match depending on which mass is used.




Clearly, the "mass number" mean molecular weights match the Skye (Jermyn et al., 2021) EOS mean molecular weights. The "atomic mass" mean molecular weights match the FreeEOS mean molecular weights.

This is not necessarily a problem. For all main sequence models, the blending region, if it even exists, is *inside* the core. This means that for the entire mixing region, the FreeEOS mean molecular weight is used. I use this same "atomic mass" prescription to compute μ for the accreted material, so to compute the mixing mass, we can simply use the full ionization with atomic masses. There are then no inconsistencies.

I have now implemented the fully ionized mean molecular weight computation into the accretor code. Since it agrees so well with the normal *MESA* EOS, there are no noticeable differences in the abundances or mixing masses.

b.3 Giant Branch Dip 2


I also plot the two different computation methods for the late RGB model.



Again, clearly the "mass number" mean molecular weights match the Skye (Jermyn et al., 2021) EOS mean molecular weights. The "atomic mass" mean molecular weights match the FreeEOS mean molecular weights.

c. Roche lobe radius Grid

Below I will compare some of the orbital properties of the broad Roche lobe radius grid that I simulated a couple of weeks ago.




This plot shows the period of the various models. The gray line shows the initial period of the models as a function of the initial Roche lobe radius. This is simply a power law based on the Eggleton (1983) Roche lobe radius approximation and Kepler's Third Law.

I also plot the final period, which is in some cases *smaller* than the initial period, and in some cases, especially towards large initial Roche lobe radii, the final period is *larger* than the initial period.


The minimum period get reached during RLOF and is always smaller than the final Roche lobe radius.

It is more obvious if we plot this compared to the initial period.



The differences with respect to the initial period increases with the Roche lobe radius. This is in part due to the tides in combination with the winds, and due to the RLOF, which seems to decrease the periods *more* for the larger masses.


d. 1.4 Solar mass grid



Here, I show the final [s/Fe] ratios for $q = 0.85$, $M_{\rm TPAGB} = 1.4\,M_\odot$ detailed *MESA* binary simulations and the simple binary simulation.

I think the *MESA* models and the simple binary models correspond quite well, but there are definitely differences. Mainly, the detailed MESA simulations seem to predict slightly larger observed abundances in many cases. For one model however, the abundances are smaller. Then, for the wind only regime, the *MESA* models agree very well with the simple binary models.


Let me first focus on the set of models with the third Roche lobe radius. These are interesting because they interact during the phase of strong winds. We can recognise this in the figure above by the clear upwards diagonal in the $\varepsilon = 0$ models. The downwards sloping diagonal line that follows it contains models that interact during the last thermal pulse.



Here, I plot the Roche lobe radius and the star radius as a function of time for the models with an initial Roche lobe radius of approximately $R_{\rm RL} = 520\,R_\odot$. The most obvious differences between the models is caused by the differences in f_δ, which is to be expected. We see that more mass loss to a CCT leads to smaller final periods. A second, initially less obvious effect is that these models interact *earlier*. This effects how much mass is lost through *winds*.

If we look at the total amount of accreted mass, we can clearly explain the differences we see between the simple binary and the detailed *MESA* simulations. For the $\varepsilon = 0$ case, the earlier RLOF interaction causes less mass to be transferred via winds for the $f_\delta = 0.3$ model. For the $f_\delta = 0.1$ model, there are still significant winds during and after RLOF, which invalidates the assumption of the simple binary that all mass is lost through RLOF once this starts. For the $\varepsilon = 0.1$ models, this effect causes the differences.

Now, let me look at the first Roche lobe radius, where we see some peculiar differences, especially for the $\varepsilon = 0, f_\delta = 0.1$ model.



I feel like, with this, I can explain the differences between the simple binary and the detailed *MESA* models. They originate from different RLOF moments and the assumption of complete RLOF mass transfer.

Apparently, these different mass transfer mechanisms cause these models to interact during different thermal pulses. This makes it slightly confusing to interpret the differences between the simple binary and the detailed *MESA* models. To start, we see that the $f_\delta = 0.3$ models interact first. This means these models will transfer mass, independent of how much, with the *lowest* enrichment. In fact, the envelope is barely enriched at all during this thermal pulse and therefore, both of these models do not show any enhanced [s/Fe]. Then, the orange $f_\delta = 0.2$, models. These interact one pulse later, but still earlier than the simple binary. This means that these models will have lower, but still slightly enhanced envelope abundances, which explains why the model with $\varepsilon = 0.1$ predicts a slightly enhanced [s/Fe], but still lower than the simple binary. Lastly, we have the $f_\delta = 0.1$ models. These undergo RLOF during the same thermal pulse, *but* they do not expel their entire envelope during RLOF, unlike what we assume for the simple binary. Because the envelope becomes enriched due to dredge up *after* the RLOF episode, wind mass losses after this episode are especially important. Because the $\varepsilon = 0.0$ model experiences more of this strongly enhanced wind, it predicts the highest [s/Fe] value.

References

Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. (2009) *The Chemical Composition of the Sun*. ARA&A47, 481.

Eggleton, P. P. (1983) *Aproximations to the radii of Roche lobes.*. ApJ268, 368.

Jermyn, A. S., Schwab, J., Bauer, E., Timmes, F. X., & Potekhin, A. Y. (2021) *Skye: A Differentiable Equation of State*. ApJ913, 72.

Rogers, F. J. & Nayfonov, A. (2002) *Updated and Expanded OPAL Equation-of-State Tables: Implications for Helioseismology*. ApJ576, 1064.