

# Combinatorial Models for Type $A$ Specialized Non-symmetric Macdonald Polynomials and Affine Demazure Crystals

Sam Spellman

March 7, 2023

# Motivation

Many known connections between:

- representation theory of semisimple Lie algebras
- Schubert calculus and geometry of flag manifolds

# Motivation

Many known connections between:

- representation theory of semisimple Lie algebras
- Schubert calculus and geometry of flag manifolds
- We want to develop a quantum/affine version of these connections.

# Motivation

Many known connections between:

- representation theory of semisimple Lie algebras
- Schubert calculus and geometry of flag manifolds
- We want to develop a quantum/affine version of these connections.

Part of this story involves nonsymmetric Macdonald polynomials, affine Demazure crystals, and combinatorics.

# Macdonald Polynomials

Polynomials in  $(x_1, x_2, \dots, x_n)$  associated to an irreducible rank  $n$  affine root system, with coefficients a rational function in  $q$  and  $t$ .

# Macdonald Polynomials

Polynomials in  $(x_1, x_2, \dots, x_n)$  associated to an irreducible rank  $n$  affine root system, with coefficients a rational function in  $q$  and  $t$ .

There are two versions:

- symmetric (under the Weyl group action)  $P_\lambda(x, q, t)$ , for a dominant weight  $\lambda$ .

# Macdonald Polynomials

Polynomials in  $(x_1, x_2, \dots, x_n)$  associated to an irreducible rank  $n$  affine root system, with coefficients a rational function in  $q$  and  $t$ .

There are two versions:

- symmetric (under the Weyl group action)  $P_\lambda(x, q, t)$ , for a dominant weight  $\lambda$ .
- non-symmetric  $E_\mu(x, q, t)$ , for an arbitrary weight  $\mu$ .

# Specialized Macdonald Polynomials

Various specializations of these polynomials give characters of representations:

- $P_\lambda(x, 0, 0)$  are the irreducible characters of simple Lie algebras.
- $P_\lambda(x, q, 0)$  are graded characters of KR modules for affine Lie algebras.



# Specialized Macdonald Polynomials

Various specializations of these polynomials give characters of representations:

- $P_\lambda(x, 0, 0)$  are the irreducible characters of simple Lie algebras.
- $P_\lambda(x, q, 0)$  are graded characters of KR modules for affine Lie algebras.
- $E_\mu(x, 0, 0)$  are characters of Demazure modules of simple Lie algebras.
- $E_\mu(x, q, 0)$  are graded characters of Demazure submodules of KR modules.

# Crystals

- A crystal is a nonempty set  $B$  along with operators  $e_i : B \rightarrow B \cup \{0\}$  and  $f_i : B \rightarrow B \cup \{0\}$  subject to some conditions.
- Can be thought of as a finite, colored, directed graph.

# Crystals

- A crystal is a nonempty set  $B$  along with operators  $e_i : B \rightarrow B \cup \{0\}$  and  $f_i : B \rightarrow B \cup \{0\}$  subject to some conditions.
- Can be thought of as a finite, colored, directed graph.
- All the representations in the last slide have a crystal structure.

# Combinatorial Models for Macdonald Polynomials

There are two main combinatorial formulas:

1. (2008) **Ram-Yip formula**, sum over alcove walks. Valid in all Lie types.

# Combinatorial Models for Macdonald Polynomials

There are two main combinatorial formulas:

1. (2008) **Ram-Yip formula**, sum over alcove walks. Valid in all Lie types.
  - For  $P_\lambda(x, q, 0)$ ,  $E_\mu(x, q, 0)$  the quantum alcove model of [LNSSS].
  - $P_\lambda(x, q, 0)$  has crystal structure due to Lenart, Lubovsky.

# Combinatorial Models for Macdonald Polynomials

There are two main combinatorial formulas:

1. (2008) **Ram-Yip formula**, sum over alcove walks. Valid in all Lie types.
  - For  $P_\lambda(x, q, 0)$ ,  $E_\mu(x, q, 0)$  the quantum alcove model of [LNSSS].
  - $P_\lambda(x, q, 0)$  has crystal structure due to Lenart, Lubovsky.
2. (2005) **Haglund-Haiman-Loehr formula**, sum over fillings of Young diagrams subject to certain conditions. Valid in type A only.

# Combinatorial Models for Macdonald Polynomials

There are two main combinatorial formulas:

1. (2008) **Ram-Yip formula**, sum over alcove walks. Valid in all Lie types.
  - For  $P_\lambda(x, q, 0)$ ,  $E_\mu(x, q, 0)$  the quantum alcove model of [LNSSS].
  - $P_\lambda(x, q, 0)$  has crystal structure due to Lenart, Lubovsky.
2. (2005) **Haglund-Haiman-Loehr formula**, sum over fillings of Young diagrams subject to certain conditions. Valid in type A only.
  - For  $E_\mu(x, q, 0)$  the relevant fillings are the semistandard key tabloids (SSKT) defined by Assaf and Gonzalez.
  - SSKT come with a crystal structure.

We want to establish a connection between the non-symmetric type  $A$  Ram-Yip formula to the HHL formula when  $t = 0$ .



We want to establish a connection between the non-symmetric type  $A$  Ram-Yip formula to the HHL formula when  $t = 0$ .

## Main Results

- In type  $A$  for  $t = 0$  we define an explicit bijection with inverse map from alcove walks to SSKT.
- We construct an affine Demazure crystal structure on alcove walks (colored directed graph).
- **Conjecture:** the bijection is compatible with our crystal operators and the operators between semistandard key tabloids by Assaf, Gonzalez.

## Type $A_{n-1}$ notation

- $V = \mathbb{R}^n / \text{span}(1, 1, \dots, 1)$
- The finite Weyl group is  $S_n$
- The root system is  $\Phi := \{\epsilon_i - \epsilon_j \mid i, j \leq n\}$

## Type $A_{n-1}$ notation

- $V = \mathbb{R}^n / \text{span}(1, 1, \dots, 1)$
- The finite Weyl group is  $S_n$
- The root system is  $\Phi := \{\epsilon_i - \epsilon_j \mid i, j \leq n\}$
- Denote by  $(i, j)$  the root  $\alpha_{ij} = \epsilon_i - \epsilon_j$  and the transposition  $t_{ij}$ ;

## Type $A_{n-1}$ notation

- $V = \mathbb{R}^n / \text{span}(1, 1, \dots, 1)$
- The finite Weyl group is  $S_n$
- The root system is  $\Phi := \{\epsilon_i - \epsilon_j \mid i, j \leq n\}$
- Denote by  $(i, j)$  the root  $\alpha_{ij} = \epsilon_i - \epsilon_j$  and the transposition  $t_{ij}$ ;
- Identify a dominant weight  $\lambda$  with the partition  
 $\lambda = (\lambda_1 \geq \dots \lambda_{n-1} \geq \lambda_n = 0)$

## Type $A_{n-1}$ notation

- $V = \mathbb{R}^n / \text{span}(1, 1, \dots, 1)$
- The finite Weyl group is  $S_n$
- The root system is  $\Phi := \{\epsilon_i - \epsilon_j \mid i, j \leq n\}$
- Denote by  $(i, j)$  the root  $\alpha_{ij} = \epsilon_i - \epsilon_j$  and the transposition  $t_{ij}$ ;
- Identify a dominant weight  $\lambda$  with the partition  $\lambda = (\lambda_1 \geq \dots \lambda_{n-1} \geq \lambda_n = 0)$
- an arbitrary weight  $\mu$  with the weak composition  $(\mu_1, \mu_2, \dots, \mu_n) \in \mathbb{Z}^n$

# The Quantum Alcove Model

For a positive root  $\alpha$  Let  $H_{\alpha,k}$  be the hyperplane defined by

$$H_{\alpha,k} = \{\lambda \in V \mid \langle \lambda, \alpha \rangle = k \text{ for } k \in \mathbb{Z}\}$$

# The Quantum Alcove Model

For a positive root  $\alpha$  Let  $H_{\alpha,k}$  be the hyperplane defined by

$$H_{\alpha,k} = \{\lambda \in V \mid \langle \lambda, \alpha \rangle = k \text{ for } k \in \mathbb{Z}\}$$

**Alcoves:** connected components of  $V \setminus (\bigcup H_{\alpha,k})$ .

# The Quantum Alcove Model

For a positive root  $\alpha$  Let  $H_{\alpha,k}$  be the hyperplane defined by

$$H_{\alpha,k} = \{\lambda \in V \mid \langle \lambda, \alpha \rangle = k \text{ for } k \in \mathbb{Z}\}$$

**Alcoves:** connected components of  $V \setminus (\bigcup H_{\alpha,k})$ .

**Fundamental alcove**  $A_o$ , with a vertex at 0.



# The Quantum Alcove Model

For a positive root  $\alpha$  Let  $H_{\alpha,k}$  be the hyperplane defined by

$$H_{\alpha,k} = \{\lambda \in V \mid \langle \lambda, \alpha \rangle = k \text{ for } k \in \mathbb{Z}\}$$

**Alcoves:** connected components of  $V \setminus (\bigcup H_{\alpha,k})$ .

**Fundamental alcove**  $A_o$ , with a vertex at 0.

An **Alcove path** is a shortest sequence of adjacent alcoves

$$\Gamma = A_o \xrightarrow{-\gamma_1} A_1 \xrightarrow{-\gamma_2} A_2 \xrightarrow{-\gamma_3} \dots \xrightarrow{-\gamma_m} A_m$$

# The Quantum Alcove Model

For a positive root  $\alpha$  Let  $H_{\alpha,k}$  be the hyperplane defined by

$$H_{\alpha,k} = \{\lambda \in V \mid \langle \lambda, \alpha \rangle = k \text{ for } k \in \mathbb{Z}\}$$

**Alcoves:** connected components of  $V \setminus (\bigcup H_{\alpha,k})$ .

**Fundamental alcove**  $A_o$ , with a vertex at 0.


An **Alcove path** is a shortest sequence of adjacent alcoves

$$\Gamma = A_o \xrightarrow{-\gamma_1} A_1 \xrightarrow{-\gamma_2} A_2 \xrightarrow{-\gamma_3} \dots \xrightarrow{-\gamma_m} A_m$$


$\Gamma$  is encoded by the sequence of roots  $(\gamma_1, \gamma_2, \dots, \gamma_m)$

Alcove picture in type  $A_2$  for  $\mu = (0, 3, 1)$

$$\Gamma = ((12)(13)|(12)|(32)(12))$$



# Quantum Bruhat Graph



Directed graph with labeled edges

$$w \xrightarrow{\alpha} ws_{\alpha}, \text{ if :}$$

$$\ell(ws_{\alpha}) = \ell(w) + 1 \text{ (covers of Bruhat order),}$$

or

$$\ell(ws_{\alpha}) = \ell(w) - 2ht(\alpha^{\vee}) + 1$$

## Special alcove paths ( $\mu$ -chains)

Fix an arbitrary weight  $\mu = (\mu_1, \dots, \mu_n)$  with  $\mu = v\lambda$ ,  $v \in S_n$  of maximal length, and  $\lambda$  a dominant weight (partition).

## Special alcove paths ( $\mu$ -chains)

Fix an arbitrary weight  $\mu = (\mu_1, \dots, \mu_n)$  with  $\mu = \nu\lambda$ ,  $\nu \in S_n$  of maximal length, and  $\lambda$  a dominant weight (partition).

Construct an alcove path

$$\Gamma := (A_0 = A_o \xrightarrow{-\gamma_1} A_1 \xrightarrow{-\gamma_2} \dots \xrightarrow{-\gamma_m} A_m = wA_o + \mu).$$

where  $w$  depends on  $\mu$ .

## Special alcove paths ( $\mu$ -chains)

Fix an arbitrary weight  $\mu = (\mu_1, \dots, \mu_n)$  with  $\mu = v\lambda$ ,  $v \in S_n$  of maximal length, and  $\lambda$  a dominant weight (partition).

Construct an alcove path

$$\Gamma := (A_0 = A_o \xrightarrow{-\gamma_1} A_1 \xrightarrow{-\gamma_2} \dots \xrightarrow{-\gamma_m} A_m = wA_o + \mu).$$

where  $w$  depends on  $\mu$ .

We will need a closely related path to  $\Gamma$ :

$$\Gamma' := -v^{-1}\Gamma = (\beta_1, \beta_2, \dots, \beta_m)$$

The  $\beta$  roots are all positive

## $v$ -admissible Subsets

From the path  $\Gamma' = (\beta_1, \beta_2, \dots, \beta_m)$ , select a subset of positions

$$J = \{j_1 < j_2 < \dots < j_s\} \subseteq [m]$$



## $v$ -admissible Subsets

From the path  $\Gamma' = (\beta_1, \beta_2, \dots, \beta_m)$ , select a subset of positions

$$J = \{j_1 < j_2 < \dots < j_s\} \subseteq [m]$$

We say  $J$  is  **$v$ -admissible** if we have a path in the quantum Bruhat graph:

$$v \leftarrow v s_{\beta_{j_1}} \leftarrow v s_{\beta_{j_1}} s_{\beta_{j_2}} \leftarrow \dots \leftarrow v s_{\beta_{j_1}} s_{\beta_{j_2}} \dots s_{\beta_{j_s}}$$

where  $s_{\beta_i}$  is the transposition  $(i_1, i_2)$  if  $\beta_i = \epsilon_{i_1} - \epsilon_{i_2}$ .

## $v$ -admissible Subsets

From the path  $\Gamma' = (\beta_1, \beta_2, \dots, \beta_m)$ , select a subset of positions

$$J = \{j_1 < j_2 < \dots < j_s\} \subseteq [m]$$

We say  $J$  is  **$v$ -admissible** if we have a path in the quantum Bruhat graph:

$$v \leftarrow v s_{\beta_{j_1}} \leftarrow v s_{\beta_{j_1}} s_{\beta_{j_2}} \leftarrow \dots \leftarrow v s_{\beta_{j_1}} s_{\beta_{j_2}} \dots s_{\beta_{j_s}}$$


where  $s_{\beta_i}$  is the transposition  $(i_1, i_2)$  if  $\beta_i = \epsilon_{i_1} - \epsilon_{i_2}$ .

$J$  can be viewed geometrically as a folding of  $\Gamma$  along hyperplanes.


Example for  $\mu = (0, 3, 1)$ ,  $\nu = 231$ ,  $J = \{1, 2, 5\}$

$$\Gamma = ((12)(13)|(12)|(32)(12))$$

$$\Gamma' = ((13)(23)|(13)|(12)(13))$$



# Quantum Bruhat graph for $W = S_3$




# Semistandard Key Tabloids ( $\text{SSKT}(\mu)$ )

# Semistandard Key Tabloids ( $\text{SSKT}(\mu)$ )

A **diagram**  $\text{dg}(\mu)$  of the weak composition  $\mu$  consists of  $\mu_i$  boxes left-justified in row  $i$ .

For example,  $\text{dg}(\mu)$  is shown below for  $\mu = (1, 0, 3, 1)$ :



# Semistandard Key Tabloids ( $\text{SSKT}(\mu)$ )

A **semistandard key tabloid** of shape  $\mu$  is a filling of  $\text{dg}(\mu)$  with positive integers from 1 to  $n$ , subject to some conditions on entries in pairs and triples of boxes.

$$\mu = (0, 2, 1, 2)$$

4	4
3	
2	2

3	3
2	
1	4

4	2
3	
1	1

# Filling map from $v$ -admissible subsets to $\text{SSKT}(\mu)$

From an alcove path  $\Gamma'$  and a  $v$ -admissible subset  $J$ :

- Start from the permutation  $v$ .
- $v^j$  is the permutation obtained by applying to  $v$  all transpositions in  $J$  up to the  $j$ -th segment of  $\Gamma'$ .
- The entry in row  $i$ , column  $j$  is

$$\text{fill}(J)(i, j) := v^j v^{-1}(i)$$




# Filling map from $v$ -admissible subsets to $\text{SSKT}(\mu)$

From an alcove path  $\Gamma'$  and a  $v$ -admissible subset  $J$ :

- Start from the permutation  $v$ .
- $v^j$  is the permutation obtained by applying to  $v$  all transpositions in  $J$  up to the  $j$ -th segment of  $\Gamma'$ .
- The entry in row  $i$ , column  $j$  is

$$\text{fill}(J)(i, j) := v^j v^{-1}(i)$$

Recall  $\Gamma' = ((13)(23)|(13)|(12)(13))$ ,  $J = \{1, 2, 5\}$ ,  $v = 231$




# Filling map from $v$ -admissible subsets to $\text{SSKT}(\mu)$

From an alcove path  $\Gamma'$  and a  $v$ -admissible subset  $J$ :

- Start from the permutation  $v$ .
- $v^j$  is the permutation obtained by applying to  $v$  all transpositions in  $J$  up to the  $j$ -th segment of  $\Gamma'$ .
- The entry in row  $i$ , column  $j$  is

$$\text{fill}(J)(i, j) := v^j v^{-1}(i)$$

Recall  $\Gamma' = ((13)(23)|(13)|(12)(13))$ ,  $J = \{1, 2, 5\}$ ,  $v = 231$



$$\text{fill}(J) = \begin{array}{|c|c|c|} \hline 2 & & \\ \hline 1 & 1 & 3 \\ \hline \end{array}$$

## Theorem

*Given a  $v$ -admissible subset  $J$ ,  $\text{fill}(J)$  is a semistandard key tabloid of shape  $\mu$  with inverse map given by a "greedy" algorithm.*

## Theorem

*The set of  $v$ -admissible subsets of  $\Gamma'$ , together with operators  $e_0, e_1, \dots, e_{n-1}$  and  $f_0, f_1, \dots, f_{n-1}$  defined by*

$$e_i(J) = J \setminus \{m\} \cup \{k\}$$

$$f_i(J) = J \setminus \{k\} \cup \{m\}$$

*form an affine Demazure crystal, where  $k$  and  $m$  are positions in the alcove path  $\Gamma$  (or possibly  $\infty$ ), and are determined by  $J$  and  $i$ .*

# Affine Demazure Crystal for $\mu = (0, 3, 1)$

