





► **Data processing inequality:** for any kernel K and $\mu, \mu' \in \mathcal{P}(X)$

$$D_f(\mu' K \| \mu K) \leq D_f(\mu' \| \mu)$$

► Therefore, if K is π -invariant, then $\pi = \pi K$ and,

$$D_f(\pi || K\mu) \leq D_f(\pi || \mu)$$

► A π -Invariant kernel can not push you away from target!

► If $X \sim \mu$ and K is π -invariant, is $X' \sim K(X, dx')$ closer to π than X ?



BUT invariance alone does not guarantee that it will bring you closer either.

► Example: if $K(x, dx') = \delta_x(dx')$ is the identity kernel, then $\mu K = \mu$ and hence,

$$D_f(\pi || K\mu) = D_f(\pi || \mu)$$

- ▶ If $X \sim \mu$ and K is π -invariant, is $X' \sim K(X, \mathrm{d}x')$ closer to π than X ?
- ▶ **Data processing inequality:** for any kernel K and $\mu, \mu' \in \mathcal{P}(\mathbb{X})$

$$D_f(\mu'K \| \mu K) \leq D_f(\mu' \| \mu)$$

- ▶ Therefore, if K is π -invariant, then $\pi = \pi K$ and,

$$D_f(\pi \| K\mu) \leq D_f(\pi \| \mu)$$

- ▶ A π -Invariant kernel can not push you away from target!
- ▶ **BUT** invariance alone does not guarantee that it will bring you closer either.
 - ▶ Example: if $K(x, \mathrm{d}x) = \delta_x(\mathrm{d}x')$ is the identity kernel, then $\mu K = \mu$ and hence,

$$D_f(\pi \| K\mu) = D_f(\pi \| \mu)$$

CONVERGENCE OF MARKOV CHAINS