



# SAMPLING: POTENTIALS

1

0

- The target  $\pi$  is a Boltzmann-Gibbs distribution

$$\pi(x) = \frac{\exp(-\beta U(x))}{Z(\beta)}$$

►  $U : \mathcal{X} \rightarrow \mathbb{R}$  is the potential energy, encoding the physics of a system

$$\gamma(x) = \exp(-\beta U(x))$$

- Normalising constant is the partition function

$$Z(\beta) = \int_{\mathbb{X}} \exp(-\beta U(x)) dx$$



Original formulation of sampling problem from statistical mechanics

►  $\beta$  is the inverse-temperature of a system



▶ We will often interpret between density and log-space

# SAMPLING: POTENTIALS

10

- ▶ Original formulation of sampling problem came from statistical mechanics
- ▶ The target  $\pi$  is a Boltzmann-Gibbs distribution

$$\pi(x) = \frac{\exp(-\beta U(x))}{Z(\beta)}$$

- ▶  $U : \mathbb{X} \rightarrow \mathbb{R}$  is the potential energy, encoding the physics of a system

$$\gamma(x) = \exp(-\beta U(x))$$

- ▶  $\beta$  is the inverse-temperature of a system
- ▶ Normalising constant is the partition function

$$Z(\beta) = \int_{\mathbb{X}} \exp(-\beta U(x)) dx$$

- ▶ We will often interpolate between density and log-space

# GOALS