



18

► A Markov chain is μ -irreducible if $\forall x \in X$, and $\mu(A) > 0$, there exists $t \geq 0$ such that

$$K^t(x, A) > 0$$

► μ -irreducible Markov chain is **aperiodic** if for all $\mu(A) > 0$,

$$\gcd(t : \mathbb{P}[X_t \in A \mid X_0 \in A] > 0) = 1$$

► **Theorem:** If K is a π -irreducible, π -invariant and aperiodic, then for any μ

$$\lim_{t \rightarrow \infty} \|\mu K^t - \pi\|_{\text{TV}} = 0$$

► Implies X_t is an approximate sample from π when t is large enough



reduces chains always bring us closer to the target, but not always quickly

► E.g. multi-modal distribution...

CONVERGENCE OF MARKOV CHAINS

18

- ▶ A Markov chain is μ -irreducible if $\forall x \in \mathbb{X}$, and $\mu(A) > 0$, there exists $t \geq 0$ such that $K^t(x, A) > 0$

- ▶ μ -irreducible Markov chain is **aperiodic** if for all $\mu(A) > 0$,

$$\gcd(t : \mathbb{P}[X_t \in A \mid X_0 \in A] > 0) = 1$$

- ▶ **Theorem:** If K is a π -irreducible, π -invariant and aperiodic, then for any μ

$$\lim_{t \rightarrow \infty} \|\mu K^t - \pi\|_{\text{TV}} = 0$$

- ▶ Implies X_t is an approximate sample from π when t is large enough
- ▶ Irreducible chains always bring you closer to the target, but not always quickly
 - ▶ E.g. multi-modal distribution...

ERGODIC THEOREM