



MAXING-TIME

2

0

► For $\epsilon > 0$, we define the **mixing time** for a π -invariant kernel K equals

$$\tau_{\text{mix}}(\epsilon) = \inf \left\{ t : \sup_{\mu \in \mathcal{P}(\mathbb{X})} \text{TV}(\mu K^t, \pi) < \epsilon \right\}$$

- The mixing time for a uniformly ergodic chain satisfies:

$$\tau_{\text{mix}}(\epsilon) < \frac{\log \frac{\epsilon}{M}}{\log \rho}$$

► τ_{mix} measures how long it takes to achieve stationarity and forget μ

► A Markov chain is **geometrically ergodic** if there exists a $M(x) > 0$ and $\rho \in [0, 1]$

$$\forall x \in \mathbb{X}, \quad \text{TV}(K^t(x, \cdot), \pi) \leq M(x)\rho^t$$

▶ A Markov chain is uniformly ergodic if $M(x) \leq M$ for some $M < \infty$

- ▶ For $\epsilon > 0$, we define the **mixing time** for a π -invariant kernel K equals

$$\tau_{\text{mix}}(\epsilon) = \inf \left\{ t : \sup_{\mu \in \mathcal{P}(\mathbb{X})} \text{TV}(\mu K^t, \pi) < \epsilon \right\}$$

- ▶ τ_{mix} measures how long it takes to achieve stationarity and forget μ
- ▶ A Markov chain is **geometrically ergodic** if there exists a $M(x) > 0$ and $\rho \in [0,1]$

$$\forall x \in \mathbb{X}, \quad \text{TV}(K^t(x, \cdot), \pi) \leq M(x)\rho^t$$

- ▶ A Markov chain is **uniformly ergodic** if $M(x) \leq M$ for some $M < \infty$
- ▶ The mixing time for a uniformly ergodic chain satisfies:

$$\tau_{\text{mix}}(\epsilon) < \frac{\log \frac{\epsilon}{M}}{\log \rho}$$

CENTRAL LIMIT THEOREM