

Sets and Set Theory

CS1F

IM Lecture 7

Craig Macdonald

Overview

2

This lecture

- Sets and Set Theory
- Relations and the Cartesian Product

Next lecture (lecture 8)

- Relational Algebra
 - The foundations for SQL

Where to go for more info...

3

- Rosen, Discrete Mathematics and Its Applications
 - sets - sections 1.4 & 1.5
 - relations - sections 6.1 & 6.2
 - <http://www.mhhe.com/math/advmath/rosen/>
- Rolland, section 3.3


Sets and Set Theory

4

Set theory is the branch of mathematics that studies **sets**

Sets are collections of objects

- The set of all numbers
- the set of all animals with tails
- the set of letters in the alphabet
- the set of all students in a class
-




Georg Cantor (1845-1918)


Sets

5

Often all members of a set have similar properties



Odd numbers
less than 10



Students in a
Tutorial Group

Set Theory - Vocabulary

6

- Objects in a set are called '**elements**' or '**members**' of a set
- A set is said to 'contain' its elements
- In databases
 - all exam scores make up a '**set**' of exam scores.
 - all employees of a company make up a '**set**' of employees

Describing Sets

7

- Describing a set
 - List all the members between braces
 - ✦ E.g. **{a, b, c, d}**
 - ✦ Represents the set with the four elements a, b, c, and d.

Describing Sets

8

- E.g. The set V of all vowels in the English alphabet
- E.g. The set O of positive integers less than 10

Describing Sets

9

- E.g. The set V of all vowels in the English alphabet

○ $V = \{a, e, i, o, u\}$

- E.g. The set O of positive integers less than 10

○ $O = \{1, 3, 5, 7, 9\}$

- $|$ denotes the *cardinality* of a set

○ $|V| = 5, |O| = 5$

Set Equality

10

- Two sets are **equal** if and only if they have the same elements

○ **Order doesn't matter**

✦ $\{1, 3, 5\} = \{1, 5, 3\} = \{3, 1, 5\} = \{3, 5, 1\} = \{5, 1, 3\} = \{5, 3, 1\}$

○ **Repetition doesn't matter**

✦ $\{1, 2\} = \{1, 1, 2\} = \{1, 2, 2, 2, 2\}$

Set Equality

11

$$A = \{1,2,3\}$$

$$B = \{3,2,1\} \quad C = \{1,1,2,2,2,3\} \quad D = \{1,2,3\}$$

Which set(s) are equal to A?

Set Equality

12

$$A = \{1,2,3\}$$

$$B = \{3,2,1\} \quad C = \{1,1,2,2,2,3\} \quad D = \{1,2,3\}$$

A = B, C and D

Sets

13

- Sets *usually* group together elements with associated properties
 - but seemingly unrelated properties can also be listed as a set
 - {2, e, Fred, Paris} is also a set
 - ✦ We just don't know much about exactly how they are related to each other

Predicates and Sets

14

- It is sometimes inconvenient or impossible to describe a set by listing all of its elements
 - What is the set of all integers less than 1 million?

Predicates and Sets

15

- It is sometimes inconvenient or impossible to describe a set by listing all of its elements
 - What is the set of all integers less than 1 million?
 - ✦ {1,2,3,4,5.....!!!!!!}

Set Builder Notation

16

- Characterise all those elements in the set by stating the properties they must have to be members

E.g.

- The set O of all positive integers less than 10 in set builder notation is:
 - $O = \{X \mid X \text{ is an odd integer less than } 10\}$
- More mathematical definitions are also OK:
 - $O = \{X \mid X \in \mathbb{N} \wedge x < 10 \wedge x \% 2 == 1 \}$

Predicates and Sets

17

- A **predicate** is sometimes used to indicate **set membership**
- A predicate $F(x)$ will be true or false, depending on whether x belongs to a set

Predicates and set membership

18

An example

$\{x \mid x \text{ is a positive integer less than } 4\}$
is the set $\{1, 2, 3\}$

If t is an element of the set $\{x \mid F(x)\}$
then the statement $F(t)$ is *true*

So if $F(x)$ is defined as $x \% 2 = 0$
 $\{x \mid F(x)\}$ contains.... the set of all even numbers

Here, $F(x)$ is referred to as the **predicate**, and x the *subject* of the *proposition*

Some Notation

20

- $a \in A$
 - a is an element of set A
- $a \notin A$
 - a is not an element of set A
- \emptyset
 - The empty or null set
 - Also represented by $\{ \}$

Graphical representation of sets


21

- Sets can be represented graphically using **Venn diagrams**
- The **universal set U** (which contains all of the objects under consideration) is represented by a rectangle
- Inside the rectangle, circles are used to represent sets
- Sometimes points are used to represent the particular elements of the set

A Example Set

22


U is the Universe
– the set of all
possible items



The set V of vowels from all letters U

A Example Set


23



The set T of people in tutorial group from all Students S

Subsets

24




A is a subset of B

$$A \subset B$$

A test that returns true iff $A \subset B$

Subsets (2)

25



A is a subset or equal to B

$$A \subseteq B$$

A test that returns true iff $A \subseteq B$

The Power Set

26

- Given a set S , the **power set** is the set of all subsets of the set S
 - Denoted by $P(S)$ or $\mathbb{P}(S)$
- E.g. the power set of $\{0,1,2\}$ is

The Power Set

27

- E.g. the power set of $\{0,1,2\}$ is
 - $P(\{0,1,2\}) = \{\emptyset, \{0\}, \{1\}, \{2\}, \{0,1\}, \{0,2\}, \{1,2\}, \{0,1,2\}\}$
 - NB - the **empty set** and the **set itself** are members of this set of subsets
- If a set has n elements, its power set has 2^n elements
- The power set does not contain numbers, it contains SETs


Set Operations

28

- Two sets can be combined in many different ways
 - The following illustrates some such combinations
- ✦ See Rosen, 1.5 for further explanations

Union

29



Symbol
like Union

The union of A and B


$A \cup B$

$$A \cup B = \{x \mid x \in A \vee x \in B\}$$

The set that contains those elements that are either in A, B, or in both

Intersection

30



The **intersection** of A and B


$A \cap B$

$$A \cap B = \{x \mid x \in A \wedge x \in B\}$$

Symbol
like aNd

Difference

31




The **difference** of A and B

$A - B$

$$A - B = \{x \mid x \in A \wedge x \notin B\}$$

Complement

32




The complement of A

\bar{A}

$$\bar{A} = \{x \mid x \notin A\}$$

Symmetric Difference

33



The symmetric difference of A and B

$$A \oplus B = (A - B) \cup (B - A)$$

$$A \oplus B = \{x \mid (x \in A \wedge x \notin B) \vee (x \in B \wedge x \notin A)\}$$

Summary

34

- What are sets?
- Notation for making sets, comparing sets
- Operators: making new sets from other sets
 - \cup **union**
 - \cap **intersection**
 - $-$ **difference**
 - \overline{A} **complement**
 - \oplus **symmetric difference**

From past exam(s)

35

Given the following sets:

$$A = \{1,3,7,9\}, B = \{2,4,6\}, C = \{7,9\}$$

determine the following (assume that P is the powerset operator):

- $|B|$ = 3
- $P(B)$ = $\{ \{ \}, \{2\}, \{4\}, \{6\}, \{2,4\}, \{4,6\}, \{2,4,6\}, \{2,6\} \}$
- $|P(C)|$ = 4
- $A \cup B$ = $\{1,2,3,4,6,7,9\}$
- $A \cap C$ = $\{7,9\}$
- Which of the following are true?
 - ✕ $C \subset A$ = TRUE
 - ✕ $C \subset B$ = FALSE

Summary of Sets

36

- What are sets? $\{1,3,5\}$
- Set builder notation for making sets; comparing sets
- Operators: making new sets from other sets
 - \cup **union**
 - \cap **intersection**
 - $-$ **difference**
 - \overline{A} **complement**
 - \oplus **symmetric difference**
- So how does this help with databases?

So why is this useful?

37

Consider a query:

“What are the grades of student 8187491?”

How can we query the database to obtain this information?

So why is this useful?

38

Two ways of querying a database:

- **procedural** (relational algebra, Pandas)

- ✦ sequence of operations
- ✦ the output of each operation is the input to the next operation

result ← **F4 (F2 (F1(tableA), tableB), F3(tableC))**

This is based on set theory

- **declarative** (SQL)

- ✦ describes the desired results (in terms of conditions)
- ✦ the DBMS works out the operations

result ← **CONDITIONS** (tableA, tableB, tableC)

This is internally implemented as RA operations

RA is key to understanding SQL query processing!

A Relation instance

39

STUDENT

name	matric	exam1	exam2
Gedge	891023	12	58
Kerr	892361	66	90
Fraser	880123	50	65

Schema

- This Student relation instance has:
 - Degree 4 and Cardinality 3
- A **relation** is a **set of tuples** (or **n-tuples**)
 - As each tuple has n values (same as degree of the relation)
 - E.g. <Fraser,880123,50,65> is a 4-tuple.

N-tuples are not sets

40

- The order of elements in a collection is sometimes important
 - But sets are unordered, so a different structure is needed
- This is provided by ***ordered n-tuples***
 - $\langle 2, 1, 5 \rangle$ is an 3-tuple

N-tuples

41

- Two ordered n-tuples are equal if and only if each corresponding pair of their elements is equal
 - $\langle a_1, a_2, \dots, a_n \rangle = \langle b_1, b_2, \dots, b_n \rangle$ if and only if $a_i = b_i$ for $i=1, 2, \dots, n$
 - $\{1, 3, 5\} = \{3, 1, 5\} = \text{TRUE}$ for SETS
 - $\langle 1, 3, 5 \rangle = \langle 3, 1, 5 \rangle = \text{FALSE}$ for N-TUPLES
- NB: We can use $\langle \rangle$ or $()$ to denote tuples, but not $\{\}$

Cartesian Product

42

Let A and B be sets

- The **cartesian product** of A and B ($A \times B$) is
 - the set of all **ordered pairs** (i.e. tuples)
 $\langle a, b \rangle$ where $a \in A$ and $b \in B$

$A = \{0, 1\}$, $B = \{a, b, c\}$

$A \times B =$

Cartesian Product

43

Let A and B be sets

- The **cartesian product** of A and B ($A \times B$) is
 - the set of all **ordered pairs** (i.e. tuples)
 $\langle a, b \rangle$ where $a \in A$ and $b \in B$

$A = \{0, 1\}$, $B = \{a, b, c\}$

$A \times B =$
 $\{\langle 0, a \rangle, \langle 0, b \rangle, \langle 0, c \rangle, \langle 1, a \rangle, \langle 1, b \rangle, \langle 1, c \rangle\}$


Connecting to Databases: Representing a relation

44

Forename = {Marilyn, Quintin, Simon, Henrik}
Surname = {McGee, Cutts, Gay}

Domains


Names = {<Marilyn, McGee>, <Quintin, Cutts>, <Simon, Gay>}



R	Cutts	McGee	Gay
Marilyn		x	
Quintin	x		
Simon			x
Henrik			

A relationship and its equivalent relation

45



Lecturer

Initials
MMcG
QC
SGay
HRight
RInnis

Teaches

Lecturer	Course
MMcG	CS1Q
QC	CS1CT
SGay	CS1P
SGay	CS1Q

Course

Code
CS1Q
CS1CT
CS1P


The table as a relation

46

Lecturers = {MM, QC, SG, HR, RI}

Courses = {CS1P, CS1Q, CS1CT}

Teaches = {<MM, CS1Q>, <QC, CS1CT>, <SG, CS1P>, <SG, CS1Q>}



Teaches	CS1P	CS1Q	CS1CT
MM		x	
QC			x
SG	x	x	
HR			
RI			

This Week

47

- A tutorial on Sets
- Next lecture:
 - Relations, and Relational Algebra
- You can also continue to work on your database **in your own time...**
 - In the lab, or from home/laptop
 - NEXT WEEK'S LAB – Populating DB with Data

Where to go for more info...

48

- Rosen, Discrete Mathematics and Its Applications
 - Sets - sections 1.4 & 1.5
 - Relations - sections 6.1 & 6.2
 - see <http://www.mhhe.com/math/advmath/rosen/>