

Your Report Title

Your Name

January 25, 2026

Abstract

Your abstract goes here.

1 Introduction

1.1 Whittaker Smoothing vs. Asymmetric Least Squares (AsLS) for Spectroscopy

Whittaker smoothing is a classic method for smoothing data, formulated as a minimization problem:

$$\min_z \sum_i (y_i - z_i)^2 + \lambda \sum_i (D^2 z_i)^2$$

where y is the observed signal, z is the smoothed signal, D^2 is the second difference operator, and λ controls the trade-off between fidelity to the data and smoothness. Whittaker smoothing is linear, treats positive and negative residuals equally (symmetric), and has a single trade-off parameter λ . The solution is closed-form.

Asymmetric Least Squares (AsLS) modifies only the data fidelity term by introducing weights:

$$\min_z \sum_i w_i (y_i - z_i)^2 + \lambda \sum_i (D^2 z_i)^2$$

where $w_i = p$ if $y_i > z_i$ and $w_i = 1 - p$ if $y_i < z_i$, with p typically close to zero. The weights depend on the current solution z , making the problem iterative and asymmetric. AsLS actively ignores peaks by down-weighting positive residuals, focusing on fitting the lower envelope of the signal—ideal for baseline removal in spectroscopy.

In AsLS, λ is typically chosen by heuristics (e.g., $\lambda \sim 10^4$ – 10^8 for spectroscopy) and visual inspection. In summary, AsLS uses Whittaker as a subproblem but adds asymmetric, adaptive weighting, making it more suitable for baseline correction in spectroscopy.

2 Methods

Describe your methods here.

3 Results

Present your results here.

4 Discussion

Discuss your findings here.

5 Conclusion

Summarize your conclusions here.