

Hilbert spaces

Exercises marked by “-B” are optional. If you solve such exercises, you will earn bonus points.

3.1. Show that the norm on the spaces $(\mathbb{C}^n, \|\cdot\|_p)$, ℓ^p , $(C[a, b], \|\cdot\|_p)$, $L^p(X, \mu)$ (where (X, μ) is a measure space containing at least two disjoint measurable sets of positive measure) is not generated by an inner product (unless $n = 1$, $p = 2$).

3.2. Generalize the parallelogram rule to n vectors.

3.3. Show that the norm on the spaces ℓ^p , $(C[a, b], \|\cdot\|_p)$, $L^p(X, \mu)$ (where (X, μ) is a measure space containing infinitely many disjoint measurable sets of positive measure) is not equivalent to a norm generated by an inner product (unless $p = 2$).

3.4. Consider the vector space $H = C[-1, 1]$ with the inner product $\langle f | g \rangle = \int_{-1}^1 f(t) \overline{g(t)} dt$. Let

$$H_0 = \left\{ f \in H : \int_{-1}^0 f(t) dt = \int_0^1 f(t) dt \right\}.$$

(a) Prove that H_0 is a closed vector subspace of H .

(b) Does the equality $H = H_0 \oplus H_0^\perp$ hold?

3.5. Prove that every incomplete inner product space H has a closed vector subspace H_0 such that $H_0 \oplus H_0^\perp \neq H$.

3.6. Let $C_c^\infty(a, b)$ be the space of smooth compactly supported functions on the interval (a, b) . Prove that for each $p \in [1, \infty)$ $C_c^\infty(a, b)$ is dense in $L^p[a, b]$.

Definition 3.1. Let $f \in L^2[a, b]$. A function $f' \in L^2[a, b]$ is a *weak derivative* of f if

$$\int_a^b f' \varphi dt = - \int_a^b f \varphi' dt$$

for all $\varphi \in C_c^\infty(a, b)$.

3.7. Prove that if $f \in L^2[a, b]$ has a weak derivative f' , then f' is unique (as an element of $L^2[a, b]$).

3.8 (the Sobolev space). Let $W^{1,2}(a, b)$ denote the space of all $f \in L^2[a, b]$ that have a weak derivative $f' \in L^2[a, b]$. Prove that $W^{1,2}(a, b)$ is a Hilbert space with respect to the inner product

$$\langle f | g \rangle = \int_a^b (f \bar{g} + f' \bar{g}') dt.$$

3.9 (the Hardy space). Let $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$, and let H^2 denote the space of holomorphic functions $f: \mathbb{D} \rightarrow \mathbb{C}$ satisfying the following condition:

$$\|f\| = \sup_{0 < r < 1} \left(\frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\varphi})|^2 d\varphi \right)^{1/2} < \infty.$$

Show that the map $f \mapsto (c_n(f))_{n \geq 0}$ (where $c_n(f)$ is the n th Taylor coefficient of f at 0) is an isometric isomorphism of $(H^2, \|\cdot\|)$ onto $\ell^2(\mathbb{Z}_{\geq 0})$. Hence H^2 is a Hilbert space.

3.10-B (the Bergman space). Let $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$, and let $L_a^2(\mathbb{D})$ denote the space of holomorphic functions $f: \mathbb{D} \rightarrow \mathbb{C}$ satisfying the following condition:

$$\|f\| = \left(\int_{\mathbb{D}} |f(x + iy)|^2 dx dy \right)^{1/2} < \infty.$$

Show that $L_a^2(\mathbb{D})$ is a closed vector subspace of $L^2(\mathbb{D})$. Hence $L_a^2(\mathbb{D})$ is a Hilbert space.