

Matematiske metoder (MM529)

Stephan Brandt

Syddansk Universitet, Odense

10. 09. 2013

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

The number a_n is called the **leading coefficient**.

The **degree** $n = \deg p$ of p is the index of the leading coefficient.

The **zero-polynomial** $p(x) = 0$ is the function mapping every $x \in \mathbb{R}$ to 0.

Its degree is defined to be $-\infty$.

Example: Multiplication of two polynomials

$$(x^3 - 3x^2 + 3x + 1)(2x^2 + x + 4) = 2x^5 - 5x^4 + 7x^3 - 7x^2 + 13x + 4.$$

Example: Multiplication of two polynomials

$$(x^3 - 3x^2 + 3x + 1)(2x^2 + x + 4) = 2x^5 - 5x^4 + 7x^3 - 7x^2 + 13x + 4.$$

Two polynomials:

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

$$q(x) = b_m x^m + b_{m-1} x^{m-1} + \dots + b_1 x + b_0.$$

Example: Multiplication of two polynomials

$$(x^3 - 3x^2 + 3x + 1)(2x^2 + x + 4) = 2x^5 - 5x^4 + 7x^3 - 7x^2 + 13x + 4.$$

Two polynomials:

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

$$q(x) = b_m x^m + b_{m-1} x^{m-1} + \dots + b_1 x + b_0.$$

Product of polynomials

$$p(x) \cdot q(x) = c_{n+m} x^{n+m} + c_{n+m-1} x^{n+m-1} + \dots + c_1 x + c_0,$$

where $c_k = a_0 b_k + a_1 b_{k-1} + \dots + a_{k-1} b_1 + a_k b_0$ (telescope sum).

Example: Multiplication of two polynomials

$$(x^3 - 3x^2 + 3x + 1)(2x^2 + x + 4) = 2x^5 - 5x^4 + 7x^3 - 7x^2 + 13x + 4.$$

Two polynomials:

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

$$q(x) = b_m x^m + b_{m-1} x^{m-1} + \dots + b_1 x + b_0.$$

Product of polynomials

$$p(x) \cdot q(x) = c_{n+m} x^{n+m} + c_{n+m-1} x^{n+m-1} + \dots + c_1 x + c_0,$$

where $c_k = a_0 b_k + a_1 b_{k-1} + \dots + a_{k-1} b_1 + a_k b_0$ (telescope sum).

Agreement: $a_i = 0$ if $i > \deg p$ and $b_i = 0$ if $i > \deg q$.

Example: Multiplication of two polynomials

$$(x^3 - 3x^2 + 3x + 1)(2x^2 + x + 4) = 2x^5 - 5x^4 + 7x^3 - 7x^2 + 13x + 4.$$

Two polynomials:

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

$$q(x) = b_m x^m + b_{m-1} x^{m-1} + \dots + b_1 x + b_0.$$

Product of polynomials

$$p(x) \cdot q(x) = c_{n+m} x^{n+m} + c_{n+m-1} x^{n+m-1} + \dots + c_1 x + c_0,$$

where $c_k = a_0 b_k + a_1 b_{k-1} + \dots + a_{k-1} b_1 + a_k b_0$ (telescope sum).

Agreement: $a_i = 0$ if $i > \deg p$ and $b_i = 0$ if $i > \deg q$.

Degree $\deg(p \cdot q) = \deg p + \deg q$.

Rational function

A **rational function** is a function of the form

$$r(x) = \frac{p(x)}{q(x)}, \quad \text{where } p(x) \text{ and } q(x) \text{ are polynomials.}$$

Rational function

A **rational function** is a function of the form

$$r(x) = \frac{p(x)}{q(x)}, \quad \text{where } p(x) \text{ and } q(x) \text{ are polynomials.}$$

If $S \subseteq \mathbb{R}$ is the set $\{x \in \mathbb{R} \mid q(x) = 0\}$ (i.e. the set of "zeroes" of q) then the domain of $r(x)$ is $\mathbb{R} \setminus S$.

Rational function

A **rational function** is a function of the form

$$r(x) = \frac{p(x)}{q(x)}, \quad \text{where } p(x) \text{ and } q(x) \text{ are polynomials.}$$

If $S \subseteq \mathbb{R}$ is the set $\{x \in \mathbb{R} \mid q(x) = 0\}$ (i.e. the set of "zeroes" of q) then the domain of $r(x)$ is $\mathbb{R} \setminus S$.

Famous rational function

$$\begin{aligned} f : \mathbb{R} \setminus \{0\} &\rightarrow \mathbb{R} \\ f(x) &= \frac{1}{x}. \end{aligned}$$

Power function $f(x) = x^k$: Variable basis x , fixed exponent k .

Exponential functions

Power function $f(x) = x^k$: Variable basis x , fixed exponent k .

Exponential f'n $g(x) = k^x$: Fixed basis $k > 0$, variable exponent x .

Power laws valid, e.g. $k^x \cdot k^y = k^{x+y}$.

Exponential functions

Power function $f(x) = x^k$: Variable basis x , fixed exponent k .

Exponential f'n $g(x) = k^x$: Fixed basis $k > 0$, variable exponent x .

Power laws valid, e.g. $k^x \cdot k^y = k^{x+y}$.

Examples of exponential functions

$$2^x, \quad \left(\frac{1}{2}\right)^x, \quad 10^x, \quad (1.1)^x, \dots$$

Exponential functions

Power function $f(x) = x^k$: Variable basis x , fixed exponent k .

Exponential f'n $g(x) = k^x$: Fixed basis $k > 0$, variable exponent x .

Power laws valid, e.g. $k^x \cdot k^y = k^{x+y}$.

Examples of exponential functions

$$2^x, \quad \left(\frac{1}{2}\right)^x, \quad 10^x, \quad (1.1)^x, \dots$$

"The" exponential function

$\exp(x) = e^x$, where $e = 2.718\dots$ (Eulerian constant).

Power function $f(x) = x^k$: Variable basis x , fixed exponent k .

Exponential f'n $g(x) = k^x$: Fixed basis $k > 0$, variable exponent x .

Power laws valid, e.g. $k^x \cdot k^y = k^{x+y}$.

Examples of exponential functions

$$2^x, \quad \left(\frac{1}{2}\right)^x, \quad 10^x, \quad (1.1)^x, \dots$$

"The" exponential function

$\exp(x) = e^x$, where $e = 2.718\dots$ (Eulerian constant).

Example: Radioactive decay

Radioactive isotope, amount $W(t)$ at time $t \geq 0$,
 $W_0 = W(0)$ initial amount.

Power function $f(x) = x^k$: Variable basis x , fixed exponent k .

Exponential f'n $g(x) = k^x$: Fixed basis $k > 0$, variable exponent x .

Power laws valid, e.g. $k^x \cdot k^y = k^{x+y}$.

Examples of exponential functions

$$2^x, \quad \left(\frac{1}{2}\right)^x, \quad 10^x, \quad (1.1)^x, \dots$$

"The" exponential function

$\exp(x) = e^x$, where $e = 2.718\dots$ (Eulerian constant).

Example: Radioactive decay

Radioactive isotope, amount $W(t)$ at time $t \geq 0$,

$W_0 = W(0)$ initial amount.

Law of radioactive decay: $W(t) = W_0 e^{-\lambda t}$.

Decay rate λ only depending on the isotope.

Properties of functions

A function $f : A \rightarrow B$ is called

- **injective**, if every $b \in B$ has at most one preimage,
- **surjective**, if every $b \in B$ has at least one preimage,
- **bijective**, if every $b \in B$ has exactly one preimage.

Properties of functions

A function $f : A \rightarrow B$ is called

- **injective**, if every $b \in B$ has at most one preimage,
- **surjective**, if every $b \in B$ has at least one preimage,
- **bijective**, if every $b \in B$ has exactly one preimage.

Observation: f is bijective if and only if it is injective and surjective.

Properties of functions

A function $f : A \rightarrow B$ is called

- **injective**, if every $b \in B$ has at most one preimage,
- **surjective**, if every $b \in B$ has at least one preimage,
- **bijective**, if every $b \in B$ has exactly one preimage.

Observation: f is bijective if and only if it is injective and surjective.

Interpretation in the graph

A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is

- **injective**, if every horizontal line intersects the graph in at most one point,
- **surjective**, if every horizontal line intersects the graph in at least one point,
- **bijective**, if every horizontal line intersects the graph in exactly one point.

Properties of functions

A function $f : A \rightarrow B$ is called

- **injective**, if every $b \in B$ has at most one preimage,
- **surjective**, if every $b \in B$ has at least one preimage,
- **bijective**, if every $b \in B$ has exactly one preimage.

Properties of functions

A function $f : A \rightarrow B$ is called

- **injective**, if every $b \in B$ has at most one preimage,
- **surjective**, if every $b \in B$ has at least one preimage,
- **bijective**, if every $b \in B$ has exactly one preimage.

Examples:

$f : \mathbb{R} \rightarrow \mathbb{R}$

- $f(x) = x^3$ is bijective.
- $f(x) = x^2$ is none of the three,
- $f(x) = e^x$ is injective, but not surjective, but

Properties of functions

A function $f : A \rightarrow B$ is called

- **injective**, if every $b \in B$ has at most one preimage,
- **surjective**, if every $b \in B$ has at least one preimage,
- **bijective**, if every $b \in B$ has exactly one preimage.

Examples:

$f : \mathbb{R} \rightarrow \mathbb{R}$

- $f(x) = x^3$ is bijective.
- $f(x) = x^2$ is none of the three,
- $f(x) = e^x$ is injective, but not surjective, but

$f : \mathbb{R} \rightarrow (0, \infty)$, $f(x) = e^x$ is bijective.

Properties of functions

A function $f : A \rightarrow B$ is called

- **injective**, if every $b \in B$ has at most one preimage,
- **surjective**, if every $b \in B$ has at least one preimage,
- **bijective**, if every $b \in B$ has exactly one preimage.

Examples:

$f : \mathbb{R} \rightarrow \mathbb{R}$

- $f(x) = x^3$ is bijective.
- $f(x) = x^2$ is none of the three,
- $f(x) = e^x$ is injective, but not surjective, but

$f : \mathbb{R} \rightarrow (0, \infty)$, $f(x) = e^x$ is bijective.

Note: Every injective function with the codomain restricted to the range is bijective.

Inverse function

For a **bijective** function $f : A \rightarrow B$, $f(x) = y$, there is a unique function $f^{-1} : B \rightarrow A$ with

$$f^{-1}(y) = x, \text{ if and only if } f(x) = y.$$

Inverse function

For a **bijective** function $f : A \rightarrow B$, $f(x) = y$, there is a unique function $f^{-1} : B \rightarrow A$ with

$$f^{-1}(y) = x, \text{ if and only if } f(x) = y.$$

graph $f = \{(x, y) : f(x) = y\}$.

graph $f^{-1} = \{(y, x) : f(x) = y\}$.

Inverse function

For a **bijective** function $f : A \rightarrow B$, $f(x) = y$, there is a unique function $f^{-1} : B \rightarrow A$ with

$$f^{-1}(y) = x, \text{ if and only if } f(x) = y.$$

graph $f = \{(x, y) : f(x) = y\}$.

graph $f^{-1} = \{(y, x) : f(x) = y\}$.

Geometric interpretation: Mirroring at angle bisector.

Inverse function

For a **bijective** function $f : A \rightarrow B$, $f(x) = y$, there is a unique function $f^{-1} : B \rightarrow A$ with

$$f^{-1}(y) = x, \text{ if and only if } f(x) = y.$$

graph $f = \{(x, y) : f(x) = y\}$.

graph $f^{-1} = \{(y, x) : f(x) = y\}$.

Geometric interpretation: Mirroring at angle bisector.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = 2x. \text{ Inverse: } f^{-1}(x) = \frac{1}{2}x.$$

Inverse function

For a **bijective** function $f : A \rightarrow B$, $f(x) = y$, there is a unique function $f^{-1} : B \rightarrow A$ with

$$f^{-1}(y) = x, \text{ if and only if } f(x) = y.$$

graph $f = \{(x, y) : f(x) = y\}$.

graph $f^{-1} = \{(y, x) : f(x) = y\}$.

Geometric interpretation: Mirroring at angle bisector.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = 2x. \text{ Inverse: } f^{-1}(x) = \frac{1}{2}x.$$

$$f(x) = x^2? \text{ Not bijective, but...}$$

Inverse function

For a **bijective** function $f : A \rightarrow B$, $f(x) = y$, there is a unique function $f^{-1} : B \rightarrow A$ with

$$f^{-1}(y) = x, \text{ if and only if } f(x) = y.$$

graph $f = \{(x, y) : f(x) = y\}$.

graph $f^{-1} = \{(y, x) : f(x) = y\}$.

Geometric interpretation: Mirroring at angle bisector.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = 2x. \text{ Inverse: } f^{-1}(x) = \frac{1}{2}x.$$

$f(x) = x^2$? Not bijective, but...

$f : [0, \infty) \rightarrow [0, \infty)$, $f(x) = x^2$ is bijective. Inverse?

$$f^{-1} : [0, \infty) \rightarrow [0, \infty), f^{-1}(x) = \sqrt{x}.$$

Inverse function

For a **bijective** function $f : A \rightarrow B$, $f(x) = y$, there is a unique function $f^{-1} : B \rightarrow A$ with

$$f^{-1}(y) = x, \text{ if and only if } f(x) = y.$$

graph $f = \{(x, y) : f(x) = y\}$.

graph $f^{-1} = \{(y, x) : f(x) = y\}$.

Geometric interpretation: Mirroring at angle bisector.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = 2x. \text{ Inverse: } f^{-1}(x) = \frac{1}{2}x.$$

$f(x) = x^2$? Not bijective, but...

$f : [0, \infty) \rightarrow [0, \infty)$, $f(x) = x^2$ is bijective. Inverse?

$$f^{-1} : [0, \infty) \rightarrow [0, \infty), f^{-1}(x) = \sqrt{x}.$$

What is the inverse of f^{-1} ?

Inverse function

For a **bijective** function $f : A \rightarrow B$, $f(x) = y$, there is a unique function $f^{-1} : B \rightarrow A$ with

$$f^{-1}(y) = x, \text{ if and only if } f(x) = y.$$

graph $f = \{(x, y) : f(x) = y\}$.

graph $f^{-1} = \{(y, x) : f(x) = y\}$.

Geometric interpretation: Mirroring at angle bisector.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = 2x. \text{ Inverse: } f^{-1}(x) = \frac{1}{2}x.$$

$f(x) = x^2$? Not bijective, but...

$f : [0, \infty) \rightarrow [0, \infty)$, $f(x) = x^2$ is bijective. Inverse?

$$f^{-1} : [0, \infty) \rightarrow [0, \infty), f^{-1}(x) = \sqrt{x}.$$

What is the inverse of f^{-1} ? $(f^{-1})^{-1} = f$.

Identity function: $\text{id}_A : A \rightarrow A, x \mapsto x$.

Identity function: $\text{id}_A : A \rightarrow A, x \mapsto x$.

$f : A \rightarrow B$, with inverse $f^{-1} : B \rightarrow A$.

Identity function: $\text{id}_A : A \rightarrow A, x \mapsto x$.

$f : A \rightarrow B$, with inverse $f^{-1} : B \rightarrow A$.

Composition

$$f^{-1} \circ f = \text{id}_A.$$

Identity function: $\text{id}_A : A \rightarrow A, x \mapsto x$.

$f : A \rightarrow B$, with inverse $f^{-1} : B \rightarrow A$.

Composition

$$f^{-1} \circ f = \text{id}_A.$$

$$f \circ f^{-1} = \text{id}_B.$$

Identity function: $\text{id}_A : A \rightarrow A, x \mapsto x$.

$f : A \rightarrow B$, with inverse $f^{-1} : B \rightarrow A$.

Composition

$$f^{-1} \circ f = \text{id}_A.$$

$$f \circ f^{-1} = \text{id}_B.$$

What is the inverse function of an exponential function?

Identity function: $\text{id}_A : A \rightarrow A, x \mapsto x$.

$f : A \rightarrow B$, with inverse $f^{-1} : B \rightarrow A$.

Composition

$$f^{-1} \circ f = \text{id}_A.$$

$$f \circ f^{-1} = \text{id}_B.$$

What is the inverse function of an exponential function?

Logarithm

Exponential function $f : \mathbb{R} \rightarrow (0, \infty)$, $f(x) = k^x$, $k > 0$, $k \neq 1$, is bijective.

Identity function: $\text{id}_A : A \rightarrow A, x \mapsto x$.

$f : A \rightarrow B$, with inverse $f^{-1} : B \rightarrow A$.

Composition

$$f^{-1} \circ f = \text{id}_A.$$

$$f \circ f^{-1} = \text{id}_B.$$

What is the inverse function of an exponential function?

Logarithm

Exponential function $f : \mathbb{R} \rightarrow (0, \infty)$, $f(x) = k^x$, $k > 0$, $k \neq 1$, is bijective.

Inverse function: $f^{-1} = \log_k : (0, \infty) \rightarrow \mathbb{R}$,
 $y = \log_k(x) = \log_k x$, if and only if $k^y = x$.

"Logarithm to the basis k of x ."

Identity function: $\text{id}_A : A \rightarrow A, x \mapsto x$.

$f : A \rightarrow B$, with inverse $f^{-1} : B \rightarrow A$.

Composition

$$f^{-1} \circ f = \text{id}_A.$$

$$f \circ f^{-1} = \text{id}_B.$$

What is the inverse function of an exponential function?

Logarithm

Exponential function $f : \mathbb{R} \rightarrow (0, \infty)$, $f(x) = k^x$, $k > 0$, $k \neq 1$, is bijective.

Inverse function: $f^{-1} = \log_k : (0, \infty) \rightarrow \mathbb{R}$,
 $y = \log_k(x) = \log_k x$, if and only if $k^y = x$.

"Logarithm to the basis k of x ."

Most common bases:

2 (computer sci.), 10 (economics), $e = 2.718\dots$ (natural sci.).

Natural logarithm

$$\ln = \log_e;$$

Natural logarithm

$$\ln = \log_e;$$

$$\ln \circ \exp = \text{id}_{\mathbb{R}};$$

$$\exp \circ \ln = \text{id}_{(0,\infty)}.$$

Natural logarithm

$$\ln = \log_e;$$

$$\ln \circ \exp = \text{id}_{\mathbb{R}};$$

$$\exp \circ \ln = \text{id}_{(0,\infty)}.$$

Rules for logarithms

$$\log_b(x \cdot y) = \log_b x + \log_b y.$$

Natural logarithm

$$\ln = \log_e;$$

$$\ln \circ \exp = \text{id}_{\mathbb{R}};$$

$$\exp \circ \ln = \text{id}_{(0,\infty)}.$$

Rules for logarithms

$$\log_b(x \cdot y) = \log_b x + \log_b y.$$

$$\log_b x = \frac{\log_a x}{\log_a b}, \text{ in particular}$$

Natural logarithm

$$\ln = \log_e;$$

$$\ln \circ \exp = \text{id}_{\mathbb{R}};$$

$$\exp \circ \ln = \text{id}_{(0,\infty)}.$$

Rules for logarithms

$$\log_b(x \cdot y) = \log_b x + \log_b y.$$

$$\log_b x = \frac{\log_a x}{\log_a b}, \text{ in particular}$$

$$\log_b x = \frac{\ln x}{\ln b}.$$

For many real life applications, a linear scale is problematic.

For many real life applications, a linear scale is problematic.
Example: unit **bel** for loudness of noise

For many real life applications, a linear scale is problematic.

Example: unit **bel** for loudness of noise

commonly known as **decibel**, one tenth of a bel

(e.g. 80 decibel = 8 bel).

For many real life applications, a linear scale is problematic.

Example: unit **bel** for loudness of noise

commonly known as **decibel**, one tenth of a bel

(e.g. 80 decibel = 8 bel).

One bel more represents a noise whose amplitude is ten times bigger.

For many real life applications, a linear scale is problematic.

Example: unit **bel** for loudness of noise

commonly known as **decibel**, one tenth of a bel

(e.g. 80 decibel = 8 bel).

One bel more represents a noise whose amplitude is ten times bigger.

(e.g. 100 decibel is a thousand times louder than 70 decibel).

Angles can be measured in **degrees** and in **radians** (the length of the circular arc on the unit circle between the two sides of the angle).

Angles can be measured in **degrees** and in **radians** (the length of the circular arc on the unit circle between the two sides of the angle). Direction: counter-clockwise (mathematically positive).

Angles can be measured in **degrees** and in **radians** (the length of the circular arc on the unit circle between the two sides of the angle). Direction: counter-clockwise (mathematically positive).
Angle $\theta = \theta + k \cdot 2\pi$, $k \in \mathbb{Z}$.

Angles can be measured in **degrees** and in **radians** (the length of the circular arc on the unit circle between the two sides of the angle). Direction: counter-clockwise (mathematically positive).

Angle $\theta = \theta + k \cdot 2\pi$, $k \in \mathbb{Z}$.

Right-angled triangle:

long side **hypotenuse**,

short side adjacent to the angle **adjacent**,

short side opposite to the angle **opposite**.

Trigonometric functions

Angles can be measured in **degrees** and in **radians** (the length of the circular arc on the unit circle between the two sides of the angle). Direction: counter-clockwise (mathematically positive).

Angle $\theta = \theta + k \cdot 2\pi$, $k \in \mathbb{Z}$.

Right-angled triangle:

long side **hypotenuse**,

short side adjacent to the angle **adjacent**,

short side opposite to the angle **opposite**.

Trigonometric functions (defined by the lengths of the sides):

$$\sin : \mathbb{R} \rightarrow [-1, 1], \quad \sin \theta = \frac{\text{opposite}}{\text{hypotenuse}};$$

Trigonometric functions

Angles can be measured in **degrees** and in **radians** (the length of the circular arc on the unit circle between the two sides of the angle). Direction: counter-clockwise (mathematically positive).

Angle $\theta = \theta + k \cdot 2\pi$, $k \in \mathbb{Z}$.

Right-angled triangle:

long side **hypotenuse**,

short side adjacent to the angle **adjacent**,

short side opposite to the angle **opposite**.

Trigonometric functions (defined by the lengths of the sides):

$$\sin : \mathbb{R} \rightarrow [-1, 1], \quad \sin \theta = \frac{\text{opposite}}{\text{hypotenuse}};$$

$$\cos : \mathbb{R} \rightarrow [-1, 1], \quad \cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}};$$

Angles can be measured in **degrees** and in **radians** (the length of the circular arc on the unit circle between the two sides of the angle). Direction: counter-clockwise (mathematically positive).

Angle $\theta = \theta + k \cdot 2\pi$, $k \in \mathbb{Z}$.

Right-angled triangle:

long side **hypotenuse**,

short side adjacent to the angle **adjacent**,

short side opposite to the angle **opposite**.

Trigonometric functions (defined by the lengths of the sides):

$$\sin : \mathbb{R} \rightarrow [-1, 1], \quad \sin \theta = \frac{\text{opposite}}{\text{hypotenuse}};$$

$$\cos : \mathbb{R} \rightarrow [-1, 1], \quad \cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}};$$

$$\tan : A \rightarrow \mathbb{R}, \quad \tan \theta = \frac{\text{opposite}}{\text{adjacent}},$$

where $A = \mathbb{R} \setminus \{(2k+1)\frac{\pi}{2} \mid k \in \mathbb{Z}\}$.

Angle defined by the points $(0, 0)$, $(1, 0)$ and a third point (x, y) on the unit circle around the origin (circle with center $(0, 0)$ and radius 1):

Angle defined by the points $(0, 0)$, $(1, 0)$ and a third point (x, y) on the unit circle around the origin (circle with center $(0, 0)$ and radius 1):

Trigonometric functions:

Angle defined by the points $(0, 0)$, $(1, 0)$ and a third point (x, y) on the unit circle around the origin (circle with center $(0, 0)$ and radius 1):

Trigonometric functions:

$$\sin : \mathbb{R} \rightarrow [-1, 1], \quad \sin \theta = \frac{y}{1} = y;$$

Angle defined by the points $(0, 0)$, $(1, 0)$ and a third point (x, y) on the unit circle around the origin (circle with center $(0, 0)$ and radius 1):

Trigonometric functions:

$$\sin : \mathbb{R} \rightarrow [-1, 1], \quad \sin \theta = \frac{y}{1} = y;$$

$$\cos : \mathbb{R} \rightarrow [-1, 1], \quad \cos \theta = \frac{x}{1} = x;$$

Angle defined by the points $(0, 0)$, $(1, 0)$ and a third point (x, y) on the unit circle around the origin (circle with center $(0, 0)$ and radius 1):

Trigonometric functions:

$$\sin : \mathbb{R} \rightarrow [-1, 1], \quad \sin \theta = \frac{y}{1} = y;$$

$$\cos : \mathbb{R} \rightarrow [-1, 1], \quad \cos \theta = \frac{x}{1} = x;$$

$$\tan : A \rightarrow \mathbb{R}, \quad \tan \theta = \frac{y}{x},$$

where $A = \mathbb{R} \setminus \{(2k + 1)\frac{\pi}{2} \mid k \in \mathbb{Z}\}$ (the cases where $x = 0$).

Identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta}.$$

Identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta}.$$

$$\sin^2 \theta + \cos^2 \theta = 1 \quad (\text{Pythagoras' Theorem}).$$

Identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta}.$$

$$\sin^2 \theta + \cos^2 \theta = 1 \quad (\text{Pythagoras' Theorem}).$$

Sums of angles:

$$\sin(\theta + \varphi) = \sin \theta \cos \varphi + \sin \varphi \cos \theta$$

$$\cos(\theta + \varphi) = \cos \theta \cos \varphi - \sin \theta \sin \varphi$$

Trigonometric functions are not bijective, but restricting the domain makes them bijective:

Trigonometric functions are not bijective, but restricting the domain makes them bijective:

$$\sin : \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow [-1, 1]; \quad \cos : [0, \pi] \rightarrow [-1, 1], ;$$

$$\tan : \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \rightarrow \mathbb{R}.$$

Trigonometric functions are not bijective, but restricting the domain makes them bijective:

$$\sin : \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow [-1, 1]; \quad \cos : [0, \pi] \rightarrow [-1, 1], ;$$

$$\tan : \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \rightarrow \mathbb{R}.$$

Inverse functions

$$\arcsin : [-1, 1] \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right];$$

$$\arccos : [-1, 1] \rightarrow [0, \pi];$$

$$\arctan : \mathbb{R} \rightarrow \left(-\frac{\pi}{2}, \frac{\pi}{2}\right).$$

Example

$(0, 1, 1, 2, 3, 5, 8, 13, 21, \dots)$

Example

$(0, 1, 1, 2, 3, 5, 8, 13, 21, \dots)$ Fibonacci sequence

Example

$(0, 1, 1, 2, 3, 5, 8, 13, 21, \dots)$ Fibonacci sequence

$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$ (recursive definition).

Example

$(0, 1, 1, 2, 3, 5, 8, 13, 21, \dots)$ Fibonacci sequence

$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$ (recursive definition).

Two interpretations:

Example

(0, 1, 1, 2, 3, 5, 8, 13, 21, ...) Fibonacci sequence

$F_0 = 0$, $F_1 = 1$, $F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$ (recursive definition).

Two interpretations:

Rabbit population

Example

$(0, 1, 1, 2, 3, 5, 8, 13, 21, \dots)$ Fibonacci sequence

$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$ (recursive definition).

Two interpretations:

Rabbit population

Family tree of a drone

Example

$(0, 1, 1, 2, 3, 5, 8, 13, 21, \dots)$ Fibonacci sequence

$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$ (recursive definition).

Two interpretations:

Rabbit population

Family tree of a drone

Sequence (of real numbers)

$$a : \mathbb{N}_0 \rightarrow \mathbb{R}$$

Written:

$$(a(0), a(1), a(2), a(3), \dots) = (a_0, a_1, a_2, a_3, \dots) = (a_n)_{n \in \mathbb{N}_0}.$$

Example

$(0, 1, 1, 2, 3, 5, 8, 13, 21, \dots)$ Fibonacci sequence

$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$ (recursive definition).

Two interpretations:

Rabbit population

Family tree of a drone

Sequence (of real numbers)

$$a : \mathbb{N}_0 \rightarrow \mathbb{R}$$

Written:

$$(a(0), a(1), a(2), a(3), \dots) = (a_0, a_1, a_2, a_3, \dots) = (a_n)_{n \in \mathbb{N}_0}.$$

Further examples:

$$\left(\frac{1}{2^n}\right)_{n \in \mathbb{N}_0} = \left(1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots\right)$$

Example

$(0, 1, 1, 2, 3, 5, 8, 13, 21, \dots)$ Fibonacci sequence

$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$ (recursive definition).

Two interpretations:

Rabbit population

Family tree of a drone

Sequence (of real numbers)

$$a : \mathbb{N}_0 \rightarrow \mathbb{R}$$

Written:

$$(a(0), a(1), a(2), a(3), \dots) = (a_0, a_1, a_2, a_3, \dots) = (a_n)_{n \in \mathbb{N}_0}.$$

Further examples:

$$\left(\frac{1}{2^n}\right)_{n \in \mathbb{N}_0} = \left(1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots\right)$$

$$\left((-1)^n\right)_{n \in \mathbb{N}_0} = (1, -1, 1, -1, 1, \dots)$$

Convergence and the limit of a sequence

A sequence $(a_n)_{n \in \mathbb{N}_0}$ is **convergent** to a **limit** $\gamma \in \mathbb{R}$ if for every $\varepsilon > 0$ there is an $n_0 \in \mathbb{N}_0$ such that $|\gamma - a_n| < \varepsilon$ for all $n > n_0$.

$$\lim_{n \rightarrow \infty} a_n = \gamma \quad \text{or short} \quad (a_n) \rightarrow \gamma \quad \text{or} \quad a_n \rightarrow \gamma.$$

Convergence and the limit of a sequence

A sequence $(a_n)_{n \in \mathbb{N}_0}$ is **convergent** to a **limit** $\gamma \in \mathbb{R}$ if for every $\varepsilon > 0$ there is an $n_0 \in \mathbb{N}_0$ such that $|\gamma - a_n| < \varepsilon$ for all $n > n_0$.

$$\lim_{n \rightarrow \infty} a_n = \gamma \quad \text{or short} \quad (a_n) \rightarrow \gamma \quad \text{or} \quad a_n \rightarrow \gamma.$$

Otherwise the sequence is **divergent**.

Convergence and the limit of a sequence

A sequence $(a_n)_{n \in \mathbb{N}_0}$ is **convergent** to a **limit** $\gamma \in \mathbb{R}$ if for every $\varepsilon > 0$ there is an $n_0 \in \mathbb{N}_0$ such that $|\gamma - a_n| < \varepsilon$ for all $n > n_0$.

$$\lim_{n \rightarrow \infty} a_n = \gamma \quad \text{or short} \quad (a_n) \rightarrow \gamma \quad \text{or} \quad a_n \rightarrow \gamma.$$

Otherwise the sequence is **divergent**.

Example: $\lim_{n \rightarrow \infty} \frac{1}{2^n} = 0$. Choose $n_0 > \log_2 \frac{1}{\varepsilon}$.

Convergence and the limit of a sequence

A sequence $(a_n)_{n \in \mathbb{N}_0}$ is **convergent** to a **limit** $\gamma \in \mathbb{R}$ if for every $\varepsilon > 0$ there is an $n_0 \in \mathbb{N}_0$ such that $|\gamma - a_n| < \varepsilon$ for all $n > n_0$.

$$\lim_{n \rightarrow \infty} a_n = \gamma \quad \text{or short} \quad (a_n) \rightarrow \gamma \quad \text{or} \quad a_n \rightarrow \gamma.$$

Otherwise the sequence is **divergent**.

Example: $\lim_{n \rightarrow \infty} \frac{1}{2^n} = 0$. Choose $n_0 > \log_2 \frac{1}{\varepsilon}$.

Infinite limits

Let (a_n) be a divergent sequence. If for every $K \in \mathbb{R}$ there is an $n_0 \in \mathbb{N}_0$ such that for every $n > n_0$

- $a_n > K$ then we say $\lim_{n \rightarrow \infty} a_n = \infty$;
- $a_n < K$ then we say $\lim_{n \rightarrow \infty} a_n = -\infty$.

Convergence and the limit of a sequence

A sequence $(a_n)_{n \in \mathbb{N}_0}$ is **convergent** to a **limit** $\gamma \in \mathbb{R}$ if for every $\varepsilon > 0$ there is an $n_0 \in \mathbb{N}_0$ such that $|\gamma - a_n| < \varepsilon$ for all $n > n_0$.

$$\lim_{n \rightarrow \infty} a_n = \gamma \quad \text{or short} \quad (a_n) \rightarrow \gamma \quad \text{or} \quad a_n \rightarrow \gamma.$$

Otherwise the sequence is **divergent**.

Example: $\lim_{n \rightarrow \infty} \frac{1}{2^n} = 0$. Choose $n_0 > \log_2 \frac{1}{\varepsilon}$.

Infinite limits

Let (a_n) be a divergent sequence. If for every $K \in \mathbb{R}$ there is an $n_0 \in \mathbb{N}_0$ such that for every $n > n_0$

- $a_n > K$ then we say $\lim_{n \rightarrow \infty} a_n = \infty$;
- $a_n < K$ then we say $\lim_{n \rightarrow \infty} a_n = -\infty$.

Examples: $\lim_{n \rightarrow \infty} n = \infty$, $\lim_{n \rightarrow \infty} \ln \frac{1}{n} = -\infty$.

$$(a_0, a_1, a_2, a_3, a_4, \dots) = (0, 0.3, 0.33, 0.333, 0.3333, \dots) \rightarrow \frac{1}{3}.$$

$(a_0, a_1, a_2, a_3, a_4, \dots) = (0, 0.3, 0.33, 0.333, 0.3333, \dots) \rightarrow \frac{1}{3}$.
If $\varepsilon \geq 10^{-k} = \frac{1}{10^k}$ take $n_0 = k + 1$.

$(a_0, a_1, a_2, a_3, a_4, \dots) = (0, 0.3, 0.33, 0.333, 0.3333, \dots) \rightarrow \frac{1}{3}$.
If $\varepsilon \geq 10^{-k} = \frac{1}{10^k}$ take $n_0 = k + 1$.

Fibonacci sequence:

$(F_0, F_1, F_2, F_3, F_4, F_5, F_6, F_7, \dots) = (0, 1, 1, 2, 3, 5, 8, 13, \dots)$:

$$\lim_{n \rightarrow \infty} F_n = \infty.$$

$(a_0, a_1, a_2, a_3, a_4, \dots) = (0, 0.3, 0.33, 0.333, 0.3333, \dots) \rightarrow \frac{1}{3}$.
If $\varepsilon \geq 10^{-k} = \frac{1}{10^k}$ take $n_0 = k + 1$.

Fibonacci sequence:

$(F_0, F_1, F_2, F_3, F_4, F_5, F_6, F_7, \dots) = (0, 1, 1, 2, 3, 5, 8, 13, \dots)$:

$$\lim_{n \rightarrow \infty} F_n = \infty.$$

Given K , any $n_0 > K + 1$ will do.

$(a_0, a_1, a_2, a_3, a_4, \dots) = (0, 0.3, 0.33, 0.333, 0.3333, \dots) \rightarrow \frac{1}{3}$.
If $\varepsilon \geq 10^{-k} = \frac{1}{10^k}$ take $n_0 = k + 1$.

Fibonacci sequence:

$(F_0, F_1, F_2, F_3, F_4, F_5, F_6, F_7, \dots) = (0, 1, 1, 2, 3, 5, 8, 13, \dots)$:

$$\lim_{n \rightarrow \infty} F_n = \infty.$$

Given K , any $n_0 > K + 1$ will do.

$((-1)^n)_{n \in \mathbb{N}_0} = (1, -1, 1, -1, 1, \dots)$ is not convergent (take $\varepsilon = \frac{1}{2}$)
and has no infinite limit (take $K = 0$).

Limit laws for convergent series

Suppose that $\lim_{n \rightarrow \infty} a_n = a$ and $\lim_{n \rightarrow \infty} b_n = b$. Then

(1) $\lim_{n \rightarrow \infty} (a_n + b_n) = a + b;$

(2) $\lim_{n \rightarrow \infty} (a_n \cdot b_n) = a \cdot b;$

(3) $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{a}{b},$ if $b_n \neq 0$ for all $n \in \mathbb{N}_0;$

(4) $\lim_{n \rightarrow \infty} (a \cdot b_n) = a \cdot b$ for $a \in \mathbb{R}.$

Limit laws for convergent series

Suppose that $\lim_{n \rightarrow \infty} a_n = a$ and $\lim_{n \rightarrow \infty} b_n = b$. Then

$$(1) \lim_{n \rightarrow \infty} (a_n + b_n) = a + b;$$

$$(2) \lim_{n \rightarrow \infty} (a_n \cdot b_n) = a \cdot b;$$

$$(3) \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{a}{b}, \text{ if } b_n \neq 0 \text{ for all } n \in \mathbb{N}_0;$$

$$(4) \lim_{n \rightarrow \infty} (a \cdot b_n) = a \cdot b \text{ for } a \in \mathbb{R}.$$

Example: $\lim_{n \rightarrow \infty} \left[\left(1 + \frac{1}{n}\right) \cdot \left(1 - \frac{1}{n}\right) \right] = \quad .$

Limit laws for convergent series

Suppose that $\lim_{n \rightarrow \infty} a_n = a$ and $\lim_{n \rightarrow \infty} b_n = b$. Then

$$(1) \lim_{n \rightarrow \infty} (a_n + b_n) = a + b;$$

$$(2) \lim_{n \rightarrow \infty} (a_n \cdot b_n) = a \cdot b;$$

$$(3) \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{a}{b}, \text{ if } b_n \neq 0 \text{ for all } n \in \mathbb{N}_0;$$

$$(4) \lim_{n \rightarrow \infty} (a \cdot b_n) = a \cdot b \text{ for } a \in \mathbb{R}.$$

Example: $\lim_{n \rightarrow \infty} \left[\left(1 + \frac{1}{n}\right) \cdot \left(1 - \frac{1}{n}\right) \right] =$.

$$\lim_{n \rightarrow \infty} 1 + \frac{1}{n} = \lim_{n \rightarrow \infty} 1 + \lim_{n \rightarrow \infty} \frac{1}{n} = 1 + 0 = 1,$$

Limit laws for convergent series

Suppose that $\lim_{n \rightarrow \infty} a_n = a$ and $\lim_{n \rightarrow \infty} b_n = b$. Then

- (1) $\lim_{n \rightarrow \infty} (a_n + b_n) = a + b$;
- (2) $\lim_{n \rightarrow \infty} (a_n \cdot b_n) = a \cdot b$;
- (3) $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{a}{b}$, if $b_n \neq 0$ for all $n \in \mathbb{N}_0$;
- (4) $\lim_{n \rightarrow \infty} (a \cdot b_n) = a \cdot b$ for $a \in \mathbb{R}$.

Example: $\lim_{n \rightarrow \infty} \left[\left(1 + \frac{1}{n}\right) \cdot \left(1 - \frac{1}{n}\right) \right] =$.

$$\lim_{n \rightarrow \infty} 1 + \frac{1}{n} = \lim_{n \rightarrow \infty} 1 + \lim_{n \rightarrow \infty} \frac{1}{n} = 1 + 0 = 1,$$

$$\lim_{n \rightarrow \infty} 1 - \frac{1}{n} = \lim_{n \rightarrow \infty} 1 - \lim_{n \rightarrow \infty} \frac{1}{n} = 1 - 0 = 1.$$

Limit laws for convergent series

Suppose that $\lim_{n \rightarrow \infty} a_n = a$ and $\lim_{n \rightarrow \infty} b_n = b$. Then

$$(1) \lim_{n \rightarrow \infty} (a_n + b_n) = a + b;$$

$$(2) \lim_{n \rightarrow \infty} (a_n \cdot b_n) = a \cdot b;$$

$$(3) \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{a}{b}, \text{ if } b_n \neq 0 \text{ for all } n \in \mathbb{N}_0;$$

$$(4) \lim_{n \rightarrow \infty} (a \cdot b_n) = a \cdot b \text{ for } a \in \mathbb{R}.$$

Example: $\lim_{n \rightarrow \infty} \left[\left(1 + \frac{1}{n}\right) \cdot \left(1 - \frac{1}{n}\right) \right] = 1 \cdot 1 = 1.$

$$\lim_{n \rightarrow \infty} 1 + \frac{1}{n} = \lim_{n \rightarrow \infty} 1 + \lim_{n \rightarrow \infty} \frac{1}{n} = 1 + 0 = 1,$$

$$\lim_{n \rightarrow \infty} 1 - \frac{1}{n} = \lim_{n \rightarrow \infty} 1 - \lim_{n \rightarrow \infty} \frac{1}{n} = 1 - 0 = 1.$$