

Matematiske metoder (MM529)

Stephan Brandt

Syddansk Universitet, Odense

03. 09. 2013

Georg Cantor 1895:

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$,

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$,

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$, $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$, $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

Elements: $1 \in \{1, 2, 3, 4\}$,

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$, $\{\clubsuit, \diamondsuit, \heartsuit, \spadesuit\}$.

Elements: $1 \in \{1, 2, 3, 4\}$, $1 \in \{1, a, B\}$,

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$, $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

Elements: $1 \in \{1, 2, 3, 4\}$, $1 \in \{1, a, B\}$, $1 \notin \{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$, $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

Elements: $1 \in \{1, 2, 3, 4\}$, $1 \in \{1, a, B\}$, $1 \notin \{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.
 $a \in \{1, a, B\}$,

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$, $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

Elements: $1 \in \{1, 2, 3, 4\}$, $1 \in \{1, a, B\}$, $1 \notin \{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.
 $a \in \{1, a, B\}$, $A \notin \{1, a, B\}$.

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$, $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

Elements: $1 \in \{1, 2, 3, 4\}$, $1 \in \{1, a, B\}$, $1 \notin \{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

$a \in \{1, a, B\}$, $A \notin \{1, a, B\}$.

Read \in : "is an element of"

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$, $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

Elements: $1 \in \{1, 2, 3, 4\}$, $1 \in \{1, a, B\}$, $1 \notin \{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

$a \in \{1, a, B\}$, $A \notin \{1, a, B\}$.

Read \in : "is an element of"

Strictly speaking **no** set:

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$, $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

Elements: $1 \in \{1, 2, 3, 4\}$, $1 \in \{1, a, B\}$, $1 \notin \{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

$a \in \{1, a, B\}$, $A \notin \{1, a, B\}$.

Read \in : "is an element of"

Strictly speaking **no** set: $\{1, 1, 2, 3, 3, 3\}$.

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$, $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

Elements: $1 \in \{1, 2, 3, 4\}$, $1 \in \{1, a, B\}$, $1 \notin \{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.
 $a \in \{1, a, B\}$, $A \notin \{1, a, B\}$.

Read \in : "is an element of"

Strictly speaking **no** set: $\{1, 1, 2, 3, 3, 3\}$.

Convention: $\{1, 1, 2, 3, 3, 3\} = \{1, 2, 3\}$.

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$, $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

Elements: $1 \in \{1, 2, 3, 4\}$, $1 \in \{1, a, B\}$, $1 \notin \{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

$a \in \{1, a, B\}$, $A \notin \{1, a, B\}$.

Read \in : "is an element of"

Strictly speaking **no** set: $\{1, 1, 2, 3, 3, 3\}$.

Convention: $\{1, 1, 2, 3, 3, 3\} = \{1, 2, 3\}$.

Set equality: Two sets are equal, if they contain the same elements.

Georg Cantor 1895:

*A **set** is a gathering together into a whole of definite, **distinct** objects of our perception or of our thought – which are called **elements** of the set.*

Sets: $\{1, 2, 3, 4\}$, $\{1, a, B\}$, $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

Elements: $1 \in \{1, 2, 3, 4\}$, $1 \in \{1, a, B\}$, $1 \notin \{\clubsuit, \diamond, \heartsuit, \spadesuit\}$.

$a \in \{1, a, B\}$, $A \notin \{1, a, B\}$.

Read \in : "is an element of"

Strictly speaking **no** set: $\{1, 1, 2, 3, 3, 3\}$.

Convention: $\{1, 1, 2, 3, 3, 3\} = \{1, 2, 3\}$.

Set equality: Two sets are equal, if they contain the same elements.

$\{1, 2, 3, 4\} = \{2, 1, 4, 3\}$.

Large sets:

Large sets:

$\{A, B, C, \dots, Z\}$ slightly ambiguous (ancient Roman? No J, no U!)

Large sets:

$\{A, B, C, \dots, Z\}$ slightly ambiguous (ancient Roman? No J, no U!)

More precise: $\{x \mid x \text{ is a capital letter of the English alphabet}\}$

Large sets:

$\{A, B, C, \dots, Z\}$ slightly ambiguous (ancient Roman? No J, no U!)

More precise: $\{x \mid x \text{ is a capital letter of the English alphabet}\}$

Special sets:

Large sets:

$\{A, B, C, \dots, Z\}$ slightly ambiguous (ancient Roman? No J, no U!)

More precise: $\{x \mid x \text{ is a capital letter of the English alphabet}\}$

Special sets:

Empty set:

$\emptyset = \{\}$, the unique set with no element (remember set equality).

Large sets:

$\{A, B, C, \dots, Z\}$ slightly ambiguous (ancient Roman? No J, no U!)

More precise: $\{x \mid x \text{ is a capital letter of the English alphabet}\}$

Special sets:

Empty set:

$\emptyset = \{\}$, the unique set with no element (remember set equality).

Integers:

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\} = \{x \mid x \text{ is an integer}\}$

Large sets:

$\{A, B, C, \dots, Z\}$ slightly ambiguous (ancient Roman? No J, no U!)

More precise: $\{x \mid x \text{ is a capital letter of the English alphabet}\}$

Special sets:

Empty set:

$\emptyset = \{\}$, the unique set with no element (remember set equality).

Integers:

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\} = \{x \mid x \text{ is an integer}\}$

Natural numbers:

$\mathbb{N} = \{1, 2, 3, \dots\} = \{x \in \mathbb{Z} \mid x > 0\}$

Large sets:

$\{A, B, C, \dots, Z\}$ slightly ambiguous (ancient Roman? No J, no U!)

More precise: $\{x \mid x \text{ is a capital letter of the English alphabet}\}$

Special sets:

Empty set:

$\emptyset = \{\}$, the unique set with no element (remember set equality).

Integers:

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\} = \{x \mid x \text{ is an integer}\}$

Natural numbers:

$\mathbb{N} = \{1, 2, 3, \dots\} = \{x \in \mathbb{Z} \mid x > 0\}$

Rational numbers:

$\mathbb{Q} = \{\frac{p}{q} \mid p \in \mathbb{Z} \text{ and } q \in \mathbb{N}\}$ (remember convention)

Large sets:

$\{A, B, C, \dots, Z\}$ slightly ambiguous (ancient Roman? No J, no U!)

More precise: $\{x \mid x \text{ is a capital letter of the English alphabet}\}$

Special sets:

Empty set:

$\emptyset = \{\}$, the unique set with no element (remember set equality).

Integers:

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\} = \{x \mid x \text{ is an integer}\}$

Natural numbers:

$\mathbb{N} = \{1, 2, 3, \dots\} = \{x \in \mathbb{Z} \mid x > 0\}$

Rational numbers:

$\mathbb{Q} = \{\frac{p}{q} \mid p \in \mathbb{Z} \text{ and } q \in \mathbb{N}\}$ (remember convention)

Real numbers:

$\mathbb{R} = \{x \mid x \text{ has a decimal representation}\}$.

Real numbers:

$$\mathbb{R} = \{x \mid x \text{ has a decimal representation}\}$$

Real numbers:

$$\mathbb{R} = \{x \mid x \text{ has a decimal representation}\}$$

Decimal Representation: $x = a_n a_{n-1} \dots a_2 a_1 a_0 . b_1 b_2 b_3 \dots$

$$a_i, b_j \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$$

Real numbers:

$$\mathbb{R} = \{x \mid x \text{ has a decimal representation}\}$$

Decimal Representation: $x = a_n a_{n-1} \dots a_2 a_1 a_0 . b_1 b_2 b_3 \dots$

$$a_i, b_j \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$$

e.g.: $\pi = 3.1415926 \dots$

Real numbers:

$$\mathbb{R} = \{x \mid x \text{ has a decimal representation}\}$$

Decimal Representation: $x = a_n a_{n-1} \dots a_2 a_1 a_0 . b_1 b_2 b_3 \dots$

$$a_i, b_j \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$$

e.g.: $\pi = 3.1415926 \dots$

Finitely many digits in front of decimal mark, (possibly) infinitely many behind.

Real numbers:

$$\mathbb{R} = \{x \mid x \text{ has a decimal representation}\}$$

Decimal Representation: $x = a_n a_{n-1} \dots a_2 a_1 a_0 . b_1 b_2 b_3 \dots$

$$a_i, b_j \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$$

e.g.: $\pi = 3.1415926 \dots$

Finitely many digits in front of decimal mark, (possibly) infinitely many behind.

Attention: In general, decimal representation not unique:

Real numbers:

$$\mathbb{R} = \{x \mid x \text{ has a decimal representation}\}$$

Decimal Representation: $x = a_n a_{n-1} \dots a_2 a_1 a_0 . b_1 b_2 b_3 \dots$

$$a_i, b_j \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$$

e.g.: $\pi = 3.1415926 \dots$

Finitely many digits in front of decimal mark, (possibly) infinitely many behind.

Attention: In general, decimal representation not unique:

$$0.99999 \dots = 1.0000 \dots = 1 \text{ (why later!)}$$

Real numbers:

$$\mathbb{R} = \{x \mid x \text{ has a decimal representation}\}$$

Decimal Representation: $x = a_n a_{n-1} \dots a_2 a_1 a_0 . b_1 b_2 b_3 \dots$

$$a_i, b_j \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$$

e.g.: $\pi = 3.1415926 \dots$

Finitely many digits in front of decimal mark, (possibly) infinitely many behind.

Attention: In general, decimal representation not unique:

$$0.99999 \dots = 1.0000 \dots = 1 \text{ (why later!)}$$

Real numbers visualized as points on the [real line](#).

A, B sets:

A, B sets:

Union of sets: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$.

A, B sets:

Union of sets: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$.

Intersection of sets: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$.

A, B sets:

Union of sets: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$.

Intersection of sets: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$.

Difference of sets:

$A \setminus B = \{x \mid x \in A \text{ and } x \notin B\} = \{x \in A \mid x \notin B\}$.

A, B sets:

Union of sets: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$.

Intersection of sets: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$.

Difference of sets:

$A \setminus B = \{x \mid x \in A \text{ and } x \notin B\} = \{x \in A \mid x \notin B\}$.

Examples:

$\{1, 2, 3, 4\} \cup \{1, a, B\} = \{1, 2, 3, 4, a, B\}$,

A, B sets:

Union of sets: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$.

Intersection of sets: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$.

Difference of sets:

$A \setminus B = \{x \mid x \in A \text{ and } x \notin B\} = \{x \in A \mid x \notin B\}$.

Examples:

$\{1, 2, 3, 4\} \cup \{1, a, B\} = \{1, 2, 3, 4, a, B\}$,

$\{1, 2, 3, 4\} \cap \{1, a, B\} = \{1\}$,

A, B sets:

Union of sets: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$.

Intersection of sets: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$.

Difference of sets:

$$A \setminus B = \{x \mid x \in A \text{ and } x \notin B\} = \{x \in A \mid x \notin B\}.$$

Examples:

$$\{1, 2, 3, 4\} \cup \{1, a, B\} = \{1, 2, 3, 4, a, B\},$$

$$\{1, 2, 3, 4\} \cap \{1, a, B\} = \{1\},$$

$$\{1, 2, 3, 4\} \setminus \{1, a, B\} = \{2, 3, 4\},$$

A, B sets:

Union of sets: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$.

Intersection of sets: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$.

Difference of sets:

$A \setminus B = \{x \mid x \in A \text{ and } x \notin B\} = \{x \in A \mid x \notin B\}$.

Examples:

$$\{1, 2, 3, 4\} \cup \{1, a, B\} = \{1, 2, 3, 4, a, B\},$$

$$\{1, 2, 3, 4\} \cap \{1, a, B\} = \{1\},$$

$$\{1, 2, 3, 4\} \setminus \{1, a, B\} = \{2, 3, 4\},$$

$$\{1, 2, 3, 4\} \cap \{\clubsuit, \diamondsuit, \heartsuit, \spadesuit\} = \emptyset,$$

A, B sets:

Union of sets: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$.

Intersection of sets: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$.

Difference of sets:

$$A \setminus B = \{x \mid x \in A \text{ and } x \notin B\} = \{x \in A \mid x \notin B\}.$$

Examples:

$$\{1, 2, 3, 4\} \cup \{1, a, B\} = \{1, 2, 3, 4, a, B\},$$

$$\{1, 2, 3, 4\} \cap \{1, a, B\} = \{1\},$$

$$\{1, 2, 3, 4\} \setminus \{1, a, B\} = \{2, 3, 4\},$$

$$\{1, 2, 3, 4\} \cap \{\clubsuit, \diamond, \heartsuit, \spadesuit\} = \emptyset,$$

$$\{1, 2, 3, 4\} \cap \mathbb{R} = \{1, 2, 3, 4\}.$$

A, B sets:

A, B sets:

$A \subseteq B$ (A is a **subset** of B), if B contains every element of A .

A, B sets:

$A \subseteq B$ (A is a subset of B), if B contains every element of A .

Examples:

A, B sets:

$A \subseteq B$ (A is a **subset** of B), if B contains every element of A .

Examples:

$$\{1, 3\} \subseteq \{1, 2, 3, 4\},$$

A, B sets:

$A \subseteq B$ (A is a **subset** of B), if B contains every element of A .

Examples:

$$\{1, 3\} \subseteq \{1, 2, 3, 4\},$$

$$\mathbb{N} \subseteq \mathbb{Q},$$

A, B sets:

$A \subseteq B$ (A is a **subset** of B), if B contains every element of A .

Examples:

$$\{1, 3\} \subseteq \{1, 2, 3, 4\},$$

$$\mathbb{N} \subseteq \mathbb{Q},$$

$$\mathbb{Q} \not\subseteq \mathbb{N},$$

A, B sets:

$A \subseteq B$ (A is a **subset** of B), if B contains every element of A .

Examples:

$$\{1, 3\} \subseteq \{1, 2, 3, 4\},$$

$$\mathbb{N} \subseteq \mathbb{Q},$$

$$\mathbb{Q} \not\subseteq \mathbb{N},$$

$$\emptyset \subseteq A \text{ for every set } A.$$

A, B sets:

$A \subseteq B$ (A is a **subset** of B), if B contains every element of A .

Examples:

$$\{1, 3\} \subseteq \{1, 2, 3, 4\},$$

$$\mathbb{N} \subseteq \mathbb{Q},$$

$$\mathbb{Q} \not\subseteq \mathbb{N},$$

$$\emptyset \subseteq A \text{ for every set } A.$$

$$A \subseteq A \text{ for every set } A.$$

A, B sets:

$A \subseteq B$ (A is a **subset** of B), if B contains every element of A .

Examples:

$$\{1, 3\} \subseteq \{1, 2, 3, 4\},$$

$$\mathbb{N} \subseteq \mathbb{Q},$$

$$\mathbb{Q} \not\subseteq \mathbb{N},$$

$$\emptyset \subseteq A \text{ for every set } A.$$

$$A \subseteq A \text{ for every set } A.$$

Proper subset: $A \subsetneq B$, if $A \subseteq B$ and $A \neq B$.

A, B sets:

$A \subseteq B$ (A is a **subset** of B), if B contains every element of A .

Examples:

$$\{1, 3\} \subseteq \{1, 2, 3, 4\},$$

$$\mathbb{N} \subseteq \mathbb{Q},$$

$$\mathbb{Q} \not\subseteq \mathbb{N},$$

$$\emptyset \subseteq A \text{ for every set } A.$$

$$A \subseteq A \text{ for every set } A.$$

Proper subset: $A \subsetneq B$, if $A \subseteq B$ and $A \neq B$.

Example:

$$\emptyset \subsetneq A \text{ for every set } A \neq \emptyset$$

Cartesian product: $A \times B = \{(x, y) \mid x \in A \text{ and } y \in B\}$.

Cartesian product: $A \times B = \{(x, y) \mid x \in A \text{ and } y \in B\}$.

Set of all ordered pairs (x, y) , where $x \in A$ and $y \in B$.

Cartesian product: $A \times B = \{(x, y) \mid x \in A \text{ and } y \in B\}$.

Set of all ordered pairs (x, y) , where $x \in A$ and $y \in B$.

$$\{1, 2\} \times \{1, a, B\} = \{(1, 1), (1, a), (1, B), (2, 1), (2, a), (2, B)\},$$

Cartesian product: $A \times B = \{(x, y) \mid x \in A \text{ and } y \in B\}$.

Set of all ordered pairs (x, y) , where $x \in A$ and $y \in B$.

$$\{1, 2\} \times \{1, a, B\} = \{(1, 1), (1, a), (1, B), (2, 1), (2, a), (2, B)\},$$
$$(1, a) \in \{1, 2\} \times \{1, a, B\},$$

Cartesian product: $A \times B = \{(x, y) \mid x \in A \text{ and } y \in B\}$.

Set of all ordered pairs (x, y) , where $x \in A$ and $y \in B$.

$$\{1, 2\} \times \{1, a, B\} = \{(1, 1), (1, a), (1, B), (2, 1), (2, a), (2, B)\},$$

$$(1, a) \in \{1, 2\} \times \{1, a, B\},$$

$$(a, 1) \notin \{1, 2\} \times \{1, a, B\}.$$

Cartesian product: $A \times B = \{(x, y) \mid x \in A \text{ and } y \in B\}$.

Set of all ordered pairs (x, y) , where $x \in A$ and $y \in B$.

$$\{1, 2\} \times \{1, a, B\} = \{(1, 1), (1, a), (1, B), (2, 1), (2, a), (2, B)\},$$

$$(1, a) \in \{1, 2\} \times \{1, a, B\},$$

$$(a, 1) \notin \{1, 2\} \times \{1, a, B\}.$$

More than two sets: $A_1 \times A_2 \times A_3 \times \dots \times A_n =$

$$\{(a_1, a_2, a_3, \dots, a_n) \mid a_i \in A_i \text{ for all } i \in \{1, 2, 3, \dots, n\}\}.$$

Cartesian product: $A \times B = \{(x, y) \mid x \in A \text{ and } y \in B\}$.

Set of all ordered pairs (x, y) , where $x \in A$ and $y \in B$.

$$\{1, 2\} \times \{1, a, B\} = \{(1, 1), (1, a), (1, B), (2, 1), (2, a), (2, B)\},$$

$$(1, a) \in \{1, 2\} \times \{1, a, B\},$$

$$(a, 1) \notin \{1, 2\} \times \{1, a, B\}.$$

More than two sets: $A_1 \times A_2 \times A_3 \times \dots \times A_n =$

$$\{(a_1, a_2, a_3, \dots, a_n) \mid a_i \in A_i \text{ for all } i \in \{1, 2, 3, \dots, n\}\}.$$

Set of all ordered n -tuples,

Cartesian product: $A \times B = \{(x, y) \mid x \in A \text{ and } y \in B\}$.

Set of all ordered pairs (x, y) , where $x \in A$ and $y \in B$.

$$\{1, 2\} \times \{1, a, B\} = \{(1, 1), (1, a), (1, B), (2, 1), (2, a), (2, B)\},$$

$$(1, a) \in \{1, 2\} \times \{1, a, B\},$$

$$(a, 1) \notin \{1, 2\} \times \{1, a, B\}.$$

More than two sets: $A_1 \times A_2 \times A_3 \times \dots \times A_n =$

$$\{(a_1, a_2, a_3, \dots, a_n) \mid a_i \in A_i \text{ for all } i \in \{1, 2, 3, \dots, n\}\}.$$

Set of all ordered n -tuples,

Important Cartesian product:

$$\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \dots \times \mathbb{R} = \{(a_1, a_2, a_3, \dots, a_n) \mid a_i \in \mathbb{R} \text{ for all } i \in \{1, 2, 3, \dots, n\}\}.$$

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$ (where A, B are sets).

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$ (where A, B are sets). Written:

$$\begin{array}{lll} f : A & \rightarrow & B \\ a & \mapsto & f(a) \end{array} \quad \text{or} \quad \begin{array}{lll} f : A & \rightarrow & B \\ f(a) & = & b \end{array}$$

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$ (where A, B are sets). Written:

$$\begin{array}{ccc} f : A & \rightarrow & B \\ a & \mapsto & f(a) \end{array} \quad \text{or} \quad \begin{array}{ccc} f : A & \rightarrow & B \\ f(a) & = & b \end{array}$$

b is the **image** (or function value) of a under f

Functions

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$ (where A, B are sets). Written:

$$\begin{array}{ccc} f : A & \rightarrow & B \\ a & \mapsto & f(a) \end{array} \quad \text{or} \quad \begin{array}{ccc} f : A & \rightarrow & B \\ f(a) & = & b \end{array}$$

b is the **image** (or function value) of a under f
Domain A , **Codomain** B ,

Functions

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$ (where A, B are sets). Written:

$$\begin{array}{ccc} f : A & \rightarrow & B \\ a & \mapsto & f(a) \end{array} \quad \text{or} \quad \begin{array}{ccc} f : A & \rightarrow & B \\ f(a) & = & b \end{array}$$

b is the **image** (or function value) of a under f

Domain A , **Codomain** B ,

Range of f : $\mathcal{R}(f) = f(A) = \{y \in B \mid f(x) = y \text{ for an } x \in A\}$.

Functions

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$ (where A, B are sets). Written:

$$\begin{array}{ccc} f : A & \rightarrow & B \\ a & \mapsto & f(a) \end{array} \quad \text{or} \quad \begin{array}{ccc} f : A & \rightarrow & B \\ f(a) & = & b \end{array}$$

b is the **image** (or function value) of a under f

Domain A , **Codomain** B ,

Range of f : $\mathcal{R}(f) = f(A) = \{y \in B \mid f(x) = y \text{ for an } x \in A\}$.

Preimage of $y \in B$: $f^{-1}(y) = \{x \in A \mid f(x) = y\}$.

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$ (where A, B are sets). Written:

$$\begin{array}{ccc} f : A & \rightarrow & B \\ a & \mapsto & f(a) \end{array} \quad \text{or} \quad \begin{array}{ccc} f : A & \rightarrow & B \\ f(a) & = & b \end{array}$$

b is the **image** (or function value) of a under f

Domain A , **Codomain** B ,

Range of f : $\mathcal{R}(f) = f(A) = \{y \in B \mid f(x) = y \text{ for an } x \in A\}$.

Preimage of $y \in B$: $f^{-1}(y) = \{x \in A \mid f(x) = y\}$.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2.$$

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$ (where A, B are sets). Written:

$$\begin{array}{ccc} f : A & \rightarrow & B \\ a & \mapsto & f(a) \end{array} \quad \text{or} \quad \begin{array}{ccc} f : A & \rightarrow & B \\ f(a) & = & b \end{array}$$

b is the **image** (or function value) of a under f

Domain A , **Codomain** B ,

Range of f : $\mathcal{R}(f) = f(A) = \{y \in B \mid f(x) = y \text{ for an } x \in A\}$.

Preimage of $y \in B$: $f^{-1}(y) = \{x \in A \mid f(x) = y\}$.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2.$$

Domain \mathbb{R} ; codomain \mathbb{R} ; range $f(A) = \{x \in \mathbb{R} \mid x \geq 0\}$.

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$ (where A, B are sets). Written:

$$\begin{array}{ccc} f : A & \rightarrow & B \\ a & \mapsto & f(a) \end{array} \quad \text{or} \quad \begin{array}{ccc} f : A & \rightarrow & B \\ f(a) & = & b \end{array}$$

b is the **image** (or function value) of a under f

Domain A , **Codomain** B ,

Range of f : $\mathcal{R}(f) = f(A) = \{y \in B \mid f(x) = y \text{ for an } x \in A\}$.

Preimage of $y \in B$: $f^{-1}(y) = \{x \in A \mid f(x) = y\}$.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2.$$

Domain \mathbb{R} ; codomain \mathbb{R} ; range $f(A) = \{x \in \mathbb{R} \mid x \geq 0\}$.

$$g : \{1, 2, 3\} \rightarrow \{1, a, B\}$$

$$g(1) = B, g(2) = 1, g(3) = B.$$

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$ (where A, B are sets). Written:

$$\begin{array}{ccc} f : A & \rightarrow & B \\ a & \mapsto & f(a) \end{array} \quad \text{or} \quad \begin{array}{ccc} f : A & \rightarrow & B \\ f(a) & = & b \end{array}$$

b is the **image** (or function value) of a under f

Domain A , **Codomain** B ,

Range of f : $\mathcal{R}(f) = f(A) = \{y \in B \mid f(x) = y \text{ for an } x \in A\}$.

Preimage of $y \in B$: $f^{-1}(y) = \{x \in A \mid f(x) = y\}$.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2.$$

Domain \mathbb{R} ; codomain \mathbb{R} ; range $f(A) = \{x \in \mathbb{R} \mid x \geq 0\}$.

$$g : \{1, 2, 3\} \rightarrow \{1, a, B\}$$

$$g(1) = B, g(2) = 1, g(3) = B.$$

Domain $\{1, 2, 3\}$; codomain $\{1, a, B\}$; range of g : $\{1, B\}$; preimage

$$g^{-1}(a) = \emptyset; g^{-1}(B) = \{1, 3\}.$$

Graph of a function

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$.

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$.

The **graph** of the function f is the point set $\{(x, f(x)) \mid x \in A\}$.

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$.

The **graph** of the function f is the point set $\{(x, f(x)) \mid x \in A\}$.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2.$$

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$.

The **graph** of the function f is the point set $\{(x, f(x)) \mid x \in A\}$.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2.$$

Graph of f : $\{(x, x^2) \mid x \in \mathbb{R}\}$.

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$.

The **graph** of the function f is the point set $\{(x, f(x)) \mid x \in A\}$.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2.$$

Graph of f : $\{(x, x^2) \mid x \in \mathbb{R}\}$.

(for functions $f : \mathbb{R} \rightarrow \mathbb{R}$ graphically representable as point set in the plane).

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$.

The **graph** of the function f is the point set $\{(x, f(x)) \mid x \in A\}$.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2.$$

Graph of f : $\{(x, x^2) \mid x \in \mathbb{R}\}$.

(for functions $f : \mathbb{R} \rightarrow \mathbb{R}$ graphically representable as point set in the plane).

$$g : \{1, 2, 3\} \rightarrow \{1, a, B\}, g(1) = B, g(2) = 1, g(3) = B.$$

A function $f : A \rightarrow B$ is a rule that assigns a **unique** element $f(a) \in B$ to **every** element $a \in A$.

The **graph** of the function f is the point set $\{(x, f(x)) \mid x \in A\}$.

Examples:

$$f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2.$$

Graph of f : $\{(x, x^2) \mid x \in \mathbb{R}\}$.

(for functions $f : \mathbb{R} \rightarrow \mathbb{R}$ graphically representable as point set in the plane).

$$g : \{1, 2, 3\} \rightarrow \{1, a, B\}, g(1) = B, g(2) = 1, g(3) = B.$$

Graph of g : $\{(1, B), (2, 1), (3, B)\}$.

Given two functions

$$f : A \rightarrow B$$

$$a \mapsto f(a)$$

$$g : B \rightarrow C$$

$$b \mapsto g(b)$$

Given two functions

$$f : A \rightarrow B$$

$$a \mapsto f(a)$$

$$g : B \rightarrow C$$

$$b \mapsto g(b)$$

The **function composition** $g \circ f$ is the function

$$g \circ f : A \rightarrow C$$

$$a \mapsto g(f(a))$$

Given two functions

$$\begin{array}{ll} f : A \rightarrow B & g : B \rightarrow C \\ a \mapsto f(a) & b \mapsto g(b) \end{array}$$

The **function composition** $g \circ f$ is the function

$$\begin{array}{ll} g \circ f : A \rightarrow C \\ a \mapsto g(f(a)) \end{array}$$

Read: " g after f " or " g of f ".

Given two functions

$$\begin{array}{ll} f : A \rightarrow B & g : B \rightarrow C \\ a \mapsto f(a) & b \mapsto g(b) \end{array}$$

The **function composition** $g \circ f$ is the function

$$\begin{array}{ll} g \circ f : A \rightarrow C \\ a \mapsto g(f(a)) \end{array}$$

Read: "g after f" or "g of f".

Example:

$$\begin{array}{ll} f : \mathbb{R} \rightarrow \mathbb{R} & g : \mathbb{R} \rightarrow \mathbb{R} \\ x \mapsto x^2 & x \mapsto 2x \end{array}$$

Given two functions

$$\begin{array}{ll} f : A \rightarrow B & g : B \rightarrow C \\ a \mapsto f(a) & b \mapsto g(b) \end{array}$$

The **function composition** $g \circ f$ is the function

$$\begin{array}{ll} g \circ f : A \rightarrow C \\ a \mapsto g(f(a)) \end{array}$$

Read: "g after f" or "g of f".

Example:

$$\begin{array}{ll} f : \mathbb{R} \rightarrow \mathbb{R} & g : \mathbb{R} \rightarrow \mathbb{R} \\ x \mapsto x^2 & x \mapsto 2x \end{array}$$

$$g \circ f(x) = g(f(x)) = g(x^2) = 2x^2.$$

Given two functions

$$\begin{array}{ll} f : A \rightarrow B & g : B \rightarrow C \\ a \mapsto f(a) & b \mapsto g(b) \end{array}$$

The **function composition** $g \circ f$ is the function

$$\begin{array}{ll} g \circ f : A \rightarrow C \\ a \mapsto g(f(a)) \end{array}$$

Read: "g after f" or "g of f".

Example:

$$\begin{array}{ll} f : \mathbb{R} \rightarrow \mathbb{R} & g : \mathbb{R} \rightarrow \mathbb{R} \\ x \mapsto x^2 & x \mapsto 2x \end{array}$$

$$g \circ f(x) = g(f(x)) = g(x^2) = 2x^2.$$

$$\text{But: } f \circ g(x) = f(g(x)) = f(2x) = (2x)^2 = 4x^2.$$

Given two functions

$$\begin{array}{ll} f : A \rightarrow B & g : B \rightarrow C \\ a \mapsto f(a) & b \mapsto g(b) \end{array}$$

The **function composition** $g \circ f$ is the function

$$\begin{array}{ll} g \circ f : A \rightarrow C \\ a \mapsto g(f(a)) \end{array}$$

Read: "g after f" or "g of f".

Example:

$$\begin{array}{ll} f : \mathbb{R} \rightarrow \mathbb{R} & g : \mathbb{R} \rightarrow \mathbb{R} \\ x \mapsto x^2 & x \mapsto 2x \end{array}$$

$$g \circ f(x) = g(f(x)) = g(x^2) = 2x^2.$$

$$\text{But: } f \circ g(x) = f(g(x)) = f(2x) = (2x)^2 = 4x^2.$$

Not commutative in general!

*k*th power of x

Function $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = x^k = \underbrace{x \cdot x \cdots x}_{k \text{ times}}$.

k th power of x

Function $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = x^k = \underbrace{x \cdot x \cdots x}_{k \text{ times}}$.

x : basis. k : exponent.

k th power of x

Function $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = x^k = \underbrace{x \cdot x \cdots x}_{k \text{ times}}$.

x : **basis**. k : **exponent**.

$x \in \mathbb{R}$, $x > 0$:

Power rules

$$x^n \cdot x^m = x^{n+m}.$$

$$x^n : x^m = x^{n-m}.$$

$$x^{-n} = \frac{1}{x^n}.$$

$$x^{\frac{n}{m}} = \sqrt[m]{x^n} = (\sqrt[m]{x})^n, \text{ if } m > 0.$$

$$x^0 = 1.$$

Multiplication/division of powers with the same basis:
addition/subtraction of the exponents.

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

The number a_n is called the **leading coefficient**.

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

The number a_n is called the **leading coefficient**.

The **degree** $n = \deg p$ of p is the index of the leading coefficient.

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

The number a_n is called the **leading coefficient**.

The **degree** $n = \deg p$ of p is the index of the leading coefficient.

The **zero-polynomial** $p(x) = 0$ is the function mapping every $x \in \mathbb{R}$ to 0.

Its degree is defined to be $-\infty$.

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

The number a_n is called the **leading coefficient**.

The **degree** $n = \deg p$ of p is the index of the leading coefficient.

The **zero-polynomial** $p(x) = 0$ is the function mapping every $x \in \mathbb{R}$ to 0.

Its degree is defined to be $-\infty$.

Example:

$$p(x) = x^4 - 4x^3 + \frac{3}{2}x + 1.$$

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

The number a_n is called the **leading coefficient**.

The **degree** $n = \deg p$ of p is the index of the leading coefficient.

The **zero-polynomial** $p(x) = 0$ is the function mapping every $x \in \mathbb{R}$ to 0.

Its degree is defined to be $-\infty$.

Example:

$$p(x) = x^4 - 4x^3 + \frac{3}{2}x + 1.$$

Coefficients:

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

The number a_n is called the **leading coefficient**.

The **degree** $n = \deg p$ of p is the index of the leading coefficient.

The **zero-polynomial** $p(x) = 0$ is the function mapping every $x \in \mathbb{R}$ to 0.

Its degree is defined to be $-\infty$.

Example:

$$p(x) = x^4 - 4x^3 + \frac{3}{2}x + 1.$$

Coefficients: $a_4 = 1$,

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

The number a_n is called the **leading coefficient**.

The **degree** $n = \deg p$ of p is the index of the leading coefficient.

The **zero-polynomial** $p(x) = 0$ is the function mapping every $x \in \mathbb{R}$ to 0.

Its degree is defined to be $-\infty$.

Example:

$$p(x) = x^4 - 4x^3 + \frac{3}{2}x + 1.$$

Coefficients: $a_4 = 1$, $a_3 = -4$,

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

The number a_n is called the **leading coefficient**.

The **degree** $n = \deg p$ of p is the index of the leading coefficient.

The **zero-polynomial** $p(x) = 0$ is the function mapping every $x \in \mathbb{R}$ to 0.

Its degree is defined to be $-\infty$.

Example:

$$p(x) = x^4 - 4x^3 + \frac{3}{2}x + 1.$$

Coefficients: $a_4 = 1$, $a_3 = -4$, $a_2 = 0$,

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

The number a_n is called the **leading coefficient**.

The **degree** $n = \deg p$ of p is the index of the leading coefficient.

The **zero-polynomial** $p(x) = 0$ is the function mapping every $x \in \mathbb{R}$ to 0.

Its degree is defined to be $-\infty$.

Example:

$$p(x) = x^4 - 4x^3 + \frac{3}{2}x + 1.$$

Coefficients: $a_4 = 1$, $a_3 = -4$, $a_2 = 0$, $a_1 = \frac{3}{2}$,

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

The number a_n is called the **leading coefficient**.

The **degree** $n = \deg p$ of p is the index of the leading coefficient.

The **zero-polynomial** $p(x) = 0$ is the function mapping every $x \in \mathbb{R}$ to 0.

Its degree is defined to be $-\infty$.

Example:

$$p(x) = x^4 - 4x^3 + \frac{3}{2}x + 1.$$

Coefficients: $a_4 = 1$, $a_3 = -4$, $a_2 = 0$, $a_1 = \frac{3}{2}$, $a_0 = 1$.

Polynomial

A (real) **polynomial** is a function $p : \mathbb{R} \rightarrow \mathbb{R}$

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where the coefficients $a_i \in \mathbb{R}$ for $i \in \{0, 1, \dots, n-1, n\}$ and $a_n \neq 0$.

The number a_n is called the **leading coefficient**.

The **degree** $n = \deg p$ of p is the index of the leading coefficient.

The **zero-polynomial** $p(x) = 0$ is the function mapping every $x \in \mathbb{R}$ to 0.

Its degree is defined to be $-\infty$.

Example:

$$p(x) = x^4 - 4x^3 + \frac{3}{2}x + 1.$$

Coefficients: $a_4 = 1$, $a_3 = -4$, $a_2 = 0$, $a_1 = \frac{3}{2}$, $a_0 = 1$. $\deg p = 4$.

Two polynomials:

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

$$q(x) = b_m x^m + b_{m-1} x^{m-1} + \dots + b_1 x + b_0,$$

$$\deg p = n \geq m = \deg q.$$

Two polynomials:

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

$$q(x) = b_m x^m + b_{m-1} x^{m-1} + \dots + b_1 x + b_0,$$

$$\deg p = n \geq m = \deg q.$$

Addition

$$\begin{aligned} p(x) + q(x) = \\ a_n x^n + \dots + a_{m+1} x^{m+1} + (a_m + b_m) x^m + \dots + (a_1 + b_1) x + a_0 + b_0. \end{aligned}$$

Two polynomials:

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

$$q(x) = b_m x^m + b_{m-1} x^{m-1} + \dots + b_1 x + b_0,$$

$$\deg p = n \geq m = \deg q.$$

Addition

$$\begin{aligned} p(x) + q(x) = \\ a_n x^n + \dots + a_{m+1} x^{m+1} + (a_m + b_m) x^m + \dots + (a_1 + b_1) x + a_0 + b_0. \end{aligned}$$

$$\text{Degree } \deg(p + q) \leq \max\{\deg p, \deg q\}.$$

Two polynomials:

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

$$q(x) = b_m x^m + b_{m-1} x^{m-1} + \dots + b_1 x + b_0,$$

$$\deg p = n \geq m = \deg q.$$

Addition

$$\begin{aligned} p(x) + q(x) = \\ a_n x^n + \dots + a_{m+1} x^{m+1} + (a_m + b_m) x^m + \dots + (a_1 + b_1) x + a_0 + b_0. \end{aligned}$$

$$\text{Degree } \deg(p + q) \leq \max\{\deg p, \deg q\}.$$

$$\deg(p + q) < \max\{\deg p, \deg q\} \text{ if } m = n \text{ and } a_n = -b_m.$$

Two polynomials:

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

$$q(x) = b_m x^m + b_{m-1} x^{m-1} + \dots + b_1 x + b_0,$$

$$\deg p = n \geq m = \deg q.$$

Addition

$$\begin{aligned} p(x) + q(x) = \\ a_n x^n + \dots + a_{m+1} x^{m+1} + (a_m + b_m) x^m + \dots + (a_1 + b_1) x + a_0 + b_0. \end{aligned}$$

$$\text{Degree } \deg(p + q) \leq \max\{\deg p, \deg q\}.$$

$$\deg(p + q) < \max\{\deg p, \deg q\} \text{ if } m = n \text{ and } a_n = -b_m.$$

Example:

$$\begin{aligned} p(x) &= x^4 - 4x^3 + \frac{3}{2}x + 1, \quad q(x) = 2x^2 + x + 4, \\ r(x) &= -2x^2 - x + 1. \end{aligned}$$