

Online Appendix

Inferential Approaches for Network Analysis: AMEN for Latent Factor Models

Shahryar Minhas¹, Peter D. Hoff¹, Michael D. Ward¹

^a*Department of Political Science, Michigan State University, East Lansing, MI 48824, USA*

^b*Department of Political Science, Duke University, Durham, NC 27701, USA*

^c*Departments of Statistics, Duke University, Durham, NC 27701, USA*

Abstract

Many network approaches have been developed in descriptive fashion, but attention to inferential approaches to network analysis has been growing. We introduce a new approach that models interdependencies among observations using additive and multiplicative effects (AME). This approach can be applied to binary, ordinal, and continuous network data, and provides a set of tools for inference from longitudinal networks as well. The AME approach is shown a) to be easy to implement; b) interpretable in a general linear model framework; c) computationally straightforward; d) not prone to degeneracy; e) captures 1st, 2nd, and 3rd order network dependencies; and f) notably outperforms multiple regression quadratic assignment procedures, exponential random graph models, and alternative latent space approaches on a variety of metrics both in- and out-of-sample.

Word Count: xxxx

[☆]This research was partially supported by the National Science Foundation Award 1259266.

*Corresponding author

1. Appendix

1.1. Limitations of Standard Framework

Generalized linear models are typically expressed via a stochastic and systematic component. The stochastic component reflects assumptions about the probability distribution from which the data are generated: $y_{ij} \sim P(Y|\theta_{ij})$, with a probability density or mass function such as the normal, binomial, or Poisson. Each dyad in the sample is independently drawn from a particular distribution, given θ_{ij} . The systematic component characterizes the model for the parameters of that distribution and describes how θ_{ij} varies as a function of a set of nodal and dyadic covariates, \mathbf{X}_{ij} : $\theta_{ij} = \beta^T \mathbf{X}_{ij}$. A fundamental assumption we make when applying this modeling technique is that given \mathbf{X}_{ij} and the parameters of the distribution, each of the dyadic observations is conditionally independent.

The importance of this assumption becomes clearer in the process of estimating a GLM via maximum likelihood. After having chosen a set of covariates and specifying a distribution, the joint density function over all dyads using the observations from Table 1 is given:

$$P(y_{ij}, y_{ik}, \dots, y_{lk} | \theta_{ij}, \theta_{ik}, \dots, \theta_{lk}) = P(y_{ij} | \theta_{ij}) \times P(y_{ik} | \theta_{ik}) \times \dots \times P(y_{lk} | \theta_{lk})$$

$$P(\mathbf{Y} | \boldsymbol{\theta}) = \prod_{\alpha=1}^{n \times (n-1)} P(y_{\alpha} | \theta_{\alpha}) \quad (1)$$

The joint probability is converted into a likelihood: $\mathcal{L}(\boldsymbol{\theta} | \mathbf{Y}) = \prod_{\alpha=1}^{n \times (n-1)} P(y_{\alpha} | \theta_{\alpha})$.

The parameters are estimated by maximizing the likelihood. The likelihood as defined above is only valid if y_{ij} is independent of y_{ji} and y_{ik} given the set of covariates we specified, or the values of θ_{ij} .

1.2. AME Model Convergence

Trace plot for AME model presented in paper.




Figure 1: Trace plot for AME model presented in paper. In this model, we utilize the SRM to account for first and second-order dependence. To account for third order dependencies we use the latent factor approach with $K = 2$.

1.3. Other Network Goodness of Fit Tests

Below we show a standard set of statistics upon which comparisons are usually conducted:¹

Variable	Description
Dyad-wise shared partners	Number of dyads in the network with exactly i shared partners.
Edge-wise shared partners	Similar to above except this counts the number of dyads with the same number of edges.
Geodesic distances	The proportion of pairs of nodes whose shortest connecting path is of length k , for $k = 1, 2, \dots$. Also, pairs of nodes that are not connected are classified as $k = \infty$.
Incoming k-star	Propensities for individuals to have connections with multiple network partners.
Indegree	Proportion of nodes with the same value of the attribute as the receiving node.
Outdegree	Proportion of nodes with the same value of the attribute as the sending node.

Table 1: Description of a set of standard statistics used to assess whether a model captures network dependencies.

We simulate 1,000 networks from the LSM, ERGM, and AME model and compare how well they align with the observed network in terms of the statistics described in Table ???. The results are shown in Figure ???. Values for the observed network are indicated by a gray bar and average values from the simulated networks for the AME, ERGM, and LSM are represented by a diamond, triangle, and square, respectively. The densely shaded interval around each point represents the 95% interval from the simulations and the taller, less dense the 90% interval.² Looking across the panels in Figure ??? it is clear that there is little difference between the ERGM and AME models in terms of how well they capture network dependencies. The LSM model, however, does perform somewhat worse in comparison here as well. Particularly, when it comes to assessing the number of edge-wise shared partners and in terms of capturing the in-degree and outdegree distributions of the collaboration network.

¹See [?] for details on each of these parameters. If one was to examine goodness of fit in the **ergm** package these parameters would be calculated by default.

²Calculation for the incoming k-star statistic is not currently supported by the **latentnet** package.




Figure 2: Goodness of fit statistics to assess how well the LSM, ERGM, and AME approaches account for network dependencies.

1.4. Comparison of **amen** & **latentnet** R Packages

Here we provide a comparison of the AME model we present in the paper with a variety of parameterizations from the **latentnet** package. The number of dimensions in the latent space in each of these cases is set to 2. LSM (SR) represents a model in which random sender and receiver effects are included. LSM (Bilinear) represents a model in which a bilinear latent model term is used instead of the default Euclidean distance term. A bilinear latent model with sender and receiver random effects is not equivalent to the AME approach that we introduce here for reasons that we have already discussed in the paper.

	LSM	LSM (Bilinear)	LSM (SR)	LSM (Bilinear + SR)	AME
Intercept/Edges	0.94* [0.09; 1.82]	-2.66* [-3.53; -1.87]	0.60 [-1.10; 2.37]	-2.50* [-4.14; -0.88]	-3.39* [-4.38; -2.50]
Conflicting policy preferences					
Business vs. NGO	-1.37* [-2.42; -0.41]	-2.64* [-4.61; -0.96]	-3.07* [-4.77; -1.56]	-2.87* [-4.63; -1.29]	-1.37* [-2.44; -0.47]
Opposition/alliance	0.00 [-0.40; 0.39]	0.04 [-0.44; 0.54]	0.31 [-0.24; 0.86]	0.24 [-0.36; 0.82]	1.08* [0.72; 1.47]
Preference dissimilarity	-1.76* [-2.62; -0.90]	-2.00* [-3.01; -1.03]	-1.88* [-3.07; -0.68]	-2.20* [-3.46; -0.96]	-0.79* [-1.55; -0.08]
Transaction costs					
Joint forum participation	1.51* [0.86; 2.17]	1.24* [0.53; 1.93]	1.56* [0.69; 2.41]	1.62* [0.70; 2.52]	0.92* [0.40; 1.47]
Influence					
Influence attribution	0.08 [-0.40; 0.55]	-0.08 [-0.62; 0.46]	0.30 [-0.37; 0.96]	0.28 [-0.42; 0.97]	1.09* [0.69; 1.53]
Alter's influence indegree	0.01 [-0.03; 0.04]	-0.05* [-0.09; -0.01]	0.06 [-0.03; 0.14]	0.05 [-0.04; 0.13]	0.11* [0.07; 0.15]
Influence absolute diff.	0.04 [-0.01; 0.09]	0.02 [-0.03; 0.07]	-0.08* [-0.14; -0.02]	-0.08* [-0.14; -0.02]	-0.07* [-0.11; -0.03]
Alter = Government actor	-0.46 [-1.08; 0.14]	-0.80 [-1.67; 0.04]	-0.11 [-1.91; 1.76]	-0.20 [-2.14; 1.74]	0.55 [-0.07; 1.15]
Functional requirements					
Ego = Environmental NGO	-0.60 [-1.32; 0.09]	-1.90* [-3.10; -0.86]	-1.69 [-3.74; 0.23]	-1.84 [-4.02; 0.11]	0.67 [-0.38; 1.71]
Same actor type	1.17* [0.63; 1.71]	1.40* [0.85; 1.95]	1.82* [1.10; 2.54]	1.90* [1.19; 2.62]	1.04* [0.63; 1.50]

Table 2: * $p < 0.05$. 95% posterior credible intervals are provided in brackets.




Figure 3: Assessments of out-of-sample predictive performance using ROC curves, separation plots, and PR curves. AUC statistics are provided as well for both curves.




Figure 4: Network goodness of fit summary using **amen**.

1.5. Comparison with other AME Parameterizations

Here we provide a comparison of the AME model we present in the paper that uses $K = 2$ for multiplicative effects and show how results change when we use $K = \{1, 3, 4\}$. Trace plots for $K = \{1, 3, 4\}$ are available upon request.

	AME (k=1)	AME (k=2)	AME (k=3)	AME (k=4)
Intercept/Edges	-3.08* [-3.91; -2.30]	-3.39* [-4.38; -2.50]	-3.72* [-4.84; -2.73]	-3.93* [-5.12; -2.87]
Conflicting policy preferences				
Business vs. NGO	-1.28* [-2.20; -0.47]	-1.37* [-2.44; -0.47]	-1.48* [-2.63; -0.49]	-1.51* [-2.69; -0.47]
Opposition/alliance	0.95* [0.64; 1.27]	1.08* [0.72; 1.47]	1.19* [0.80; 1.64]	1.28* [0.86; 1.77]
Preference dissimilarity	-0.65* [-1.30; -0.03]	-0.79* [-1.55; -0.08]	-0.89* [-1.71; -0.12]	-0.95* [-1.80; -0.14]
Transaction costs				
Joint forum participation	0.84* [0.38; 1.31]	0.92* [0.40; 1.47]	1.01* [0.44; 1.62]	1.06* [0.43; 1.72]
Influence				
Influence attribution	1.00* [0.63; 1.39]	1.09* [0.69; 1.53]	1.21* [0.75; 1.71]	1.28* [0.80; 1.84]
Alter's influence indegree	0.10* [0.07; 0.14]	0.11* [0.07; 0.15]	0.12* [0.08; 0.17]	0.13* [0.09; 0.18]
Influence absolute diff.	-0.06* [-0.10; -0.03]	-0.07* [-0.11; -0.03]	-0.07* [-0.12; -0.04]	-0.08* [-0.12; -0.04]
Alter = Government actor	0.52 [-0.04; 1.07]	0.55 [-0.07; 1.15]	0.60 [-0.07; 1.27]	0.64 [-0.07; 1.35]
Functional requirements				
Ego = Environmental NGO	0.61 [-0.31; 1.56]	0.67 [-0.38; 1.71]	0.76 [-0.38; 1.90]	0.80 [-0.40; 2.04]
Same actor type	0.97* [0.60; 1.35]	1.04* [0.63; 1.50]	1.11* [0.64; 1.59]	1.17* [0.68; 1.68]

Table 3: * $p < 0.05$. 95% posterior credible intervals are provided in brackets.




Figure 5: Assessments of out-of-sample predictive performance using ROC curves, separation plots, and PR curves. AUC statistics are provided as well for both curves.




Figure 6: Network goodness of fit summary using **amen**.