

## Chapter 5

### Problems

1. (a)  $c \int_2^4 \left(x - \frac{3}{x^2}\right) dx = 1 \Rightarrow c = \frac{4}{21}$   
 (b)  $F(x) = \frac{4}{21} \int_2^x \left(x - \frac{3}{x^2}\right) dx = \frac{4}{21} \left(\frac{3}{x} + \frac{x^2}{2} - \frac{7}{2}\right), 2 < x < 4$
  
4. (a)  $\int_{20}^{\infty} \frac{10}{x^2} dx = \frac{-10}{x} \Big|_{20}^{\infty} = 1/2.$   
 (b)  $F(y) = \int_{10}^y \frac{10}{x^2} dx = 1 - \frac{10}{y}, y > 10. F(y) = 0 \text{ for } y < 10.$   
 (c)  $\sum_{i=3}^6 \binom{6}{i} \left(\frac{2}{3}\right)^i \left(\frac{1}{3}\right)^{6-i}$  since  $\bar{F}(15) = \frac{10}{15}$ . Assuming independence of the events that the devices exceed 15 hours.
  
7.  $\int_1^3 (a + bx^3) dx = 1 \Rightarrow 2a + 20b = 1$   
 $\int_1^3 x(a + bx^3) dx = 5 \Rightarrow 4a + \frac{242b}{5} = 5 \Rightarrow 20a + 242b = 25$   
 Solving the two simultaneous equations gives us the following:  $a = -\frac{43}{14}, b = \frac{5}{14}.$
  
10. (a)  $P\{\text{goes to } A\} = P\{5 < X < 15 \text{ or } 20 < X < 30 \text{ or } 35 < X < 45 \text{ or } 50 < X < 60\}.$   
 $= 2/3 \text{ since } X \text{ is uniform } (0, 60).$   
 (b) same answer as in (a).
  
13. (a) Let  $X$  be the random variable represent the waiting time of the passenger.  
 Then,  $P(X > 6) = \int_6^{15} \frac{1}{15} dx = \frac{3}{5}.$   
 (b)  $P(X > 10 \mid X > 8) = \frac{P(X > 10)}{P(X > 8)} = \frac{\int_{10}^{15} \frac{1}{15} dx}{\int_8^{15} \frac{1}{15} dx} = \frac{15}{21}.$
  
16.  $P\{X > 50\} = P\left\{\frac{X - 40}{4} > \frac{10}{4}\right\} = 1 - \Phi(2.5) = 1 - .9938$   
 Hence,  $(P\{X < 50\})^{10} = (.9938)^{10}$

23. (a) Let  $R_n$  denote the number of times a red card appears in  $n$  picks and  $Z$  be a standard normal random variable. Then,

$$P(249.5 < R_{100} < 300.5) = P\left(\frac{249.5-500(0.5)}{\sqrt{500(0.5)(0.5)}} < \frac{R_{100}-500(0.5)}{\sqrt{500(0.5)(0.5)}} < \frac{300.5-500(0.5)}{\sqrt{500(0.5)(0.5)}}\right) \approx$$

$$P\left(\frac{249.5-500(0.5)}{\sqrt{500(0.5)(0.5)}} < Z < \frac{300.5-500(0.5)}{\sqrt{500(0.5)(0.5)}}\right) = P(-0.0447 < Z < 4.5169) = \Phi(4.5169) - \Phi(-0.0447) = 0.5178.$$

(b) Let  $E_n$  denote the number of times an even card appears in  $n$  picks and  $Z$  be a standard normal random variable. Then,

$$P(E_{100} > 200) = P\left(\frac{E_{100}-500\left(\frac{20}{52}\right)}{\sqrt{500\left(\frac{20}{52}\right)\left(\frac{32}{52}\right)}} > \frac{200-500\left(\frac{20}{52}\right)}{\sqrt{500\left(\frac{20}{52}\right)\left(\frac{32}{52}\right)}}\right) \approx P\left(Z > \frac{200-500\left(\frac{20}{52}\right)}{\sqrt{500\left(\frac{20}{52}\right)\left(\frac{32}{52}\right)}}\right) = P(Z > 0.7071) = 1 - \Phi(0.7071) = 0.2398.$$

32.

$$\frac{1}{\lambda} = 1.5 \Rightarrow \lambda = \frac{2}{3}; \text{ therefore, } P(X > x) = 1 - F(x) = e^{-\frac{2}{3}x}.$$

$$(a) P(X > 2) = e^{-\frac{4}{3}} = 0.2636.$$

$$(b) \text{ By memoryless property, } P(X > 2 \mid X > 1) = P(X > 1) = e^{-\frac{2}{3}} = 0.5134.$$

38. (a)  $F_Y(y) = P(Y \leq y) = P(\log(X) \leq y) = P(X \leq e^y) = \frac{e^y - 1}{4}$  for  $\log(1) \leq y \leq \log(5)$ . Therefore, the probability density function is given by the following:

$$f_Y(y) = \frac{dF_Y(y)}{dy} = \begin{cases} \frac{1}{4}e^y, & \log(1) \leq y \leq \log(5) \\ 0, & \text{otherwise} \end{cases}$$

$$(b) P\left(\frac{1}{3} < Y < \frac{2}{3}\right) = F_Y\left(\frac{2}{3}\right) - F_Y\left(\frac{1}{3}\right) = 0.1380.$$

$$40. F_Y(y) = P(Y \leq y) = P(e^{-\lambda X} \leq y) = P\left(\log(e^{-\lambda X}) \leq \log(y)\right) = P\left(X \geq -\frac{1}{\lambda} \log(y)\right) = e^{-\lambda\left(-\frac{1}{\lambda} \log(y)\right)} = y \text{ for } 0 < y \leq 1.$$

Therefore, the probability density function is given by the following equation:

$$f_Y(y) = \frac{dF_Y(y)}{dy} = \begin{cases} 1, & y \in (0,1] \\ 0, & \text{otherwise,} \end{cases}$$

, i.e.,  $Y$  is a uniform random variable on  $(0,1]$ .

## Self-Test Problems and Exercises

3.

**5.3.** First, let us find  $c$  by using

$$1 = \int_0^2 cx^4 dx = 32c/5 \Rightarrow c = 5/32$$

$$(a) E[X] = \frac{5}{32} \int_0^2 x^5 dx = \frac{5}{32} \frac{64}{6} = 5/3$$

$$(b) E[X^2] = \frac{5}{32} \int_0^2 x^6 dx = \frac{5}{32} \frac{128}{7} = 20/7 \Rightarrow \text{Var}(X) = 20/7 - (5/3)^2 = 5/63$$

3. 首先讓我們藉由下式求  $c$

$$1 = \int_0^2 cx^4 dx = 32c/5 \Rightarrow c = 5/32$$

$$(a) E[X] = \frac{5}{32} \int_0^2 x^5 dx = \frac{5}{32} \frac{64}{6} = 5/3$$

$$(b) E[X^2] = \frac{5}{32} \int_0^2 x^6 dx = \frac{5}{32} \frac{128}{7} = 20/7 \Rightarrow \text{Var}(X) = 20/7 - (5/3)^2 = 5/63$$

10.

**5.10.** Let  $X$  be the tire life in units of one thousand, and let  $Z = (X - 34)/4$ . Note that  $Z$  is a standard normal random variable.

$$(a) P\{X > 40\} = P\{Z > 1.5\} \approx .0668$$

$$(b) P\{30 < X < 35\} = P\{-1 < Z < .25\} = P\{Z < .25\} - P\{Z > 1\} \approx .44$$

$$(c) P\{X > 40 | X > 30\} = P\{X > 40\} / P\{X > 30\} \\ = P\{Z > 1.5\} / P\{Z > -1\} \approx .079$$

10. 令  $X$  為輪胎壽命以千里為單位，且令  $Z = (X - 34) / 4$ 。注意到  $Z$  為一個標準常態隨機變數。

$$(a) P\{X > 40\} = P\{Z > 1.5\} \approx .0668$$

$$(b) P\{30 < X < 35\} = P\{-1 < Z < .25\} = P\{Z < .25\} - P\{Z > 1\} \approx .44$$

$$(c) P\{X > 40 | X > 30\} = P\{X > 40\} / P\{X > 30\} \\ = P\{Z > 1.5\} / P\{Z > -1\} \approx .079$$

13.

**5.13.** The lack of memory property of the exponential gives the result  $e^{-4/5}$ .

13. 由指數分配的無記憶性可得  $e^{-4/5}$ 。