Chapter 1

Problems

10.

10.

(a) $8!$

(b) $7! \times 2!$

(c) $4! \times \binom{2}{1} \times 4!$

(d) $4! \times 5!$

(e) $4! \times 2^4$

12. $10^3 - 10 \cdot 9 \cdot 8 = 280$ numbers have at least 2 equal values. $280 - 10 = 270$ have exactly 2 equal values.

14. (a) 30^5

(b) $30 \cdot 29 \cdot 28 \cdot 27 \cdot 26$

17. There are $\binom{10}{5}\binom{12}{5}$ possible choices of the 5 men and 5 women. They can then be paired up in 5! ways, since if we arbitrarily order the men then the first man can be paired with any of the 5 women, the next with any of the remaining 4, and so on. Hence, there are $5!\binom{10}{5}\binom{12}{5}$ possible results.

21. (a) There are $\binom{8}{3}\binom{4}{3}+\binom{8}{3}\binom{2}{1}\binom{4}{2}$ = 896 possible committees.

There are $\binom{8}{3}\binom{4}{3}$ that do not contain either of the 2 men, and there are $\binom{8}{3}\binom{2}{1}\binom{4}{2}$ that contain exactly 1 of them.

(b) There are $\binom{6}{3}\binom{6}{3}+\binom{2}{1}\binom{6}{2}\binom{6}{3}$ = 1000 possible committees.

(c) There are $\binom{7}{3}\binom{5}{3}+\binom{7}{2}\binom{5}{3}+\binom{7}{3}\binom{5}{2}$ = 910 possible committees. There are $\binom{7}{3}\binom{5}{3}$ in which neither feuding party serves; $\binom{7}{2}\binom{5}{3}$ in which the feuding women serves; and $\binom{7}{3}\binom{5}{2}$ in which the feuding man serves.

24. There are $\frac{4!}{2!2!}$ paths from A to the circled point; and $\frac{3!}{2!1!}$ paths from the circled point to B. Thus, by the basic principle, there are 18 different paths from A to B that go through the circled point.

29.

$$(x_1 + 2x_2 + 3x_3)^4$$

$$= \sum_{\substack{(n_1,n_2,n_3):\\ n_1+n_2+n_3=4}} \binom{4}{n_1, n_2, n_3} x_1^{n_1} (2x_2)^{n_2} (3x_3)^{n_3}$$

31. The total number of ways is $3^{10} = 59{,}049$.

(a) One among the 3 friends will be given 4 presents and the other 2 friends will be given 3 presents each. Therefore, the total number of ways is $\binom{10}{4,3,3} + \binom{10}{3,4,3} + \binom{10}{3,3,4} = 3\left(\frac{10!}{4!3!3!}\right) = 12{,}600.$

34. (a) number of nonnegative integer solutions of $x_1 + x_2 + x_3 + x_4 = 8$.

Hence, answer is $\binom{11}{3} = 165$

(b) here it is the number of positive solutions—hence answer is $\binom{7}{3} = 35$

35. (a) number of nonnegative solutions of $x_1 + \ldots + x_6 = 8$

answer = $\binom{13}{5}$

(b) (number of solutions of $x_1 + \ldots + x_6 = 5$) × (number of solutions of $x_1 + \ldots + x_6 = 3$) = $\binom{10}{5}\binom{8}{5}$

36. (a) $x_1 + x_2 + x_3 + x_4 = 20$, $x_1 \geq 2$, $x_2 \geq 2$, $x_3 \geq 3$, $x_4 \geq 4$

Let $y_1 = x_1 - 1$, $y_2 = x_2 - 1$, $y_3 = x_3 - 2$, $y_4 = x_4 - 3$

$y_1 + y_2 + y_3 + y_4 = 13$, $y_i > 0$

Hence, there are $\binom{12}{3} = 220$ possible strategies.

(b) there are $\binom{15}{2}$ investments only in 1, 2, 3

there are $\binom{14}{2}$ investments only in 1, 2, 4

there are $\binom{13}{2}$ investments only in 1, 3, 4

there are $\binom{13}{2}$ investments only in 2, 3, 4

$\binom{15}{2} + \binom{14}{2} + 2\binom{13}{2} + \binom{12}{3} = 552$ possibilities

37. (a) $\binom{14}{4} = 1001$

(b) $\binom{10}{3} = 120$

(c) There are $\binom{13}{3} = 286$ possible outcomes having 0 trout caught and $\binom{12}{3} = 220$ possible outcomes having 1 trout caught. Hence, using (a), there are $1001 - 286 - 220 = 495$ possible outcomes in which at least 2 of the 10 are trout.

(a) 令 X_i 分別代表第 i 種魚的補獲數量，$i=1,2,\dots,5$

$\Rightarrow X_1+X_2+\dots+X_5=10$，$X_i\geq 0$，$i=1,2,\dots,5$

$\therefore \binom{10+5-1}{5-1}=\binom{14}{4}=1001$ #

(b) 令 X_1 代表鱒魚數量 $\Rightarrow X_1=3$

因此 $X_2+X_3+X_4+X_5=10-3=7$，$X_i\geq 0$，$i=2,3,\dots,5$

$\therefore \binom{7+4-1}{4-1}=\binom{10}{3}=120$ #

(c) 令 X_1 代表鱒魚數量

10條中至少2條是鱒魚

$\Rightarrow$ 全部 $-$ 0條 $-$ 1條

0條 $\Rightarrow X_2+X_3+X_4+X_5=10 \Rightarrow \binom{10+4-1}{4-1}=\binom{13}{3}$

1條 $\Rightarrow X_2+X_3+X_4+X_5=10-1=9 \Rightarrow \binom{9+4-1}{4-1}=\binom{12}{3}$

因此 $\binom{14}{4}-\binom{13}{3}-\binom{12}{3}$

$=1001-286-220=495$ #

Self-Test Problems and Exercises

11.

11. (a) 我們可以將此視為 7 個階段的實驗。首先從團體中選出 6 對代表性的夫妻，然後再從每對中選出 1 人。依一般化的基本計數原理，共有 $\binom{10}{6}2^6$ 不同的選擇。

(b) 首先從團體中選出 6 對夫妻，然後再從其中選出 3 對來貢獻 1 個男人。所以共 $\binom{10}{6}\binom{6}{3}=\frac{10!}{4!3!3!}$ 不同的選擇。

13.

13. （方程式 $x_1+\cdots+x_5=4$ 之解的個數）（$x_1+\cdots+x_5=5$ 之解的個數）（$x_1+\cdots+x_5=6$ 之解的個數）$=\binom{8}{4}\binom{9}{4}\binom{10}{4}$

19.

19. 首先選擇 3 個位置給數字，然後放字母和數字。所以，共有 $\binom{8}{3}\cdot 26\cdot 25\cdot 24\cdot 23\cdot 22\cdot 10\cdot 9\cdot 8$ 種不同的牌照。若數字必須連續，則共有 6 種可能的位置給數字，造成了共有 $6\cdot 26\cdot 25\cdot 24\cdot 23\cdot 22\cdot 10\cdot 9\cdot 8$ 種不同的牌照。