






How Neutrino Oscillations Change Explodability of Massive Stars?

Mariam Gogilashvili

Niels Bohr International Academy and DARK, Niels Bohr Institute, university of Copenhagen, Blegdamsvej 17, 2100, Copenhagen, Denmark

VILLUM FONDEN

Which Massive Stars Explode?



The Force Explosion Condition


$$\int_{R_{NS}}^{R_S + \epsilon} \frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0$$

$$\rho \left(\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} \right) = -\nabla p - \rho \nabla \Phi$$

$$\int_{R_{NS}} \frac{\partial(\rho E)}{\partial t} + \nabla \cdot \left[\rho \mathbf{v} \left(\epsilon + \frac{p}{\rho} + \frac{v^2}{2} + \Phi \right) \right] = \rho H - \rho C$$

$$\tilde{L}_\nu \tau_g - a\tilde{\kappa} + \tilde{W}_b + c\tilde{R}_r^r > b$$

Gogilashvili et al. (2021, 2022, 2023)



Flavor Conversions and Explodability

$$n_{\nu_e(\bar{\nu}_e)}^{\text{new}} = \frac{1}{3}(n_{\nu_e(\bar{\nu}_e)}^{\text{old}} + 2n_{\nu_x(\bar{\nu}_x)}^{\text{old}})$$

$$L_{\nu_e(\bar{\nu}_e)}^{\text{new}} = \frac{1}{3}(L_{\nu_e(\bar{\nu}_e)}^{\text{old}} + 2L_{\nu_x(\bar{\nu}_x)}^{\text{old}})$$

$$\kappa^{\text{new}} = \frac{(L_{\nu_e}^{\text{old}} + 2L_{\nu_x}^{\text{old}}) \left(\frac{1}{3}\kappa_{\nu_e}^{\text{old}} + \frac{2}{3}\kappa_{\nu_x}^{\text{old}} \right) + (L_{\bar{\nu}_e}^{\text{old}} + 2L_{\bar{\nu}_x}^{\text{old}}) \left(\frac{1}{3}\kappa_{\bar{\nu}_e}^{\text{old}} + \frac{2}{3}\kappa_{\bar{\nu}_x}^{\text{old}} \right)}{(L_{\nu_e}^{\text{old}} + 2L_{\nu_x}^{\text{old}}) + (L_{\bar{\nu}_e}^{\text{old}} + 2L_{\bar{\nu}_x}^{\text{old}})}$$

Gogilashvili et al. (2025 in prep)