

## 4. Roots of equations: open methods

- fixed point iteration
- Newton-Raphson
- secant method
- modified Newton-Raphson
- MATLAB `fzero` function



## Open versus bracketing methods

## Bracketing methods

- start with  $[x_l, x_u]$  where  $f(x_l)f(x_u) < 0$  (root within interval)
- repeated halving/refinement  $\Rightarrow$  always convergent (moves closer to the true root)

## Open methods

- use formulas with one start value or two that need not bracket the root
- may *diverge* (move away from the true root)
- when they converge, they typically do so *faster* than bracketing methods



## Fixed-point iteration

rearrange  $f(x) = 0$  into *fixed-point* form

$$x = g(x)$$

- for example

- algebraic manipulation:  $x^2 - 2x + 3 = 0 \Rightarrow x = \frac{x^2 + 3}{2}$  or  $x = \sqrt{2x - 3}$

- add  $x$  to both sides:  $\sin x = 0 \Rightarrow x = x + \sin x$

- given an initial guess  $x_0$ , iterate

$$x_{i+1} = g(x_i), \quad i = 0, 1, \dots$$

called *fixed point iteration* (or *one-point iteration* or *successive substitution*)

- monitor the approximate relative error

$$\varepsilon_a = \left| \frac{x_{i+1} - x_i}{x_{i+1}} \right| \times 100\%$$

## Example

$$f(x) = e^{-x} - x$$

true root:  $x^{\star} \approx 0.567143140453502$

separate directly and iterate  $x_{i+1} = g(x_i) = e^{-x_i}$  with  $x_0 = 0$

$i$	$x_i$	$x_{i+1} = g(x_i)$	$\varepsilon_a$	$\varepsilon_t$	$\varepsilon_{t,i}/\varepsilon_{t,i-1}$
0	0.0000	1.0000		76.32%	
1	1.0000	0.3679	100%	35.13%	46.03%
2	0.3679	0.6922	171.83%	22.05%	62.76%
3	0.6922	0.5005	46.85%	11.76%	53.31%
4	0.5005	0.6062	38.31%	6.89%	58.65%
5	0.6062	0.5454	17.45%	3.83%	55.62%
6	0.5454	0.5796	11.16%	2.20%	57.34%
7	0.5796	0.5601	5.90%	1.24%	56.36%
8	0.5601	0.5711	3.48%	0.71%	56.91%
9	0.5711	0.5649	1.93%	0.40%	56.60%
10	0.5649	0.5684	1.11%	0.23%	56.78%

error proportional to that in previous iteration (about 60%), called *linear convergence*

## Example

choose alternative formulation  $x = g(x) = -\ln(x)$  with starting point  $x_0 = 0.5$

$i$	$x_i$	$g(x_i)$	$\varepsilon_a$	$\varepsilon_t$	$\varepsilon_{t,i}/\varepsilon_{t,i-1}$
0	0.5000	0.6931		11.84%	
1	0.6931	0.3665	27.87%	22.22%	1.8766
2	0.3665	1.0037	89.12%	35.38%	1.5923
3	1.0037	-0.0037	63.48%	76.98%	2.1760

diverges in this case


## Two-curve graphical method

- split  $f(x) = f_1(x) - f_2(x) = 0$  into  $y_1 = f_1(x)$  and  $y_2 = f_2(x)$
- the intersections give roots

**Example:** for  $e^{-x} - x = 0$ , take  $y_1 = x$ ,  $y_2 = e^{-x}$

$x$	$y_1 = x$	$y_2 = e^{-x}$
0.0	0.000	1.000
0.2	0.200	0.819
0.4	0.400	0.670
0.6	0.600	0.549
0.8	0.800	0.449
1.0	1.000	0.368

intersection near  $x \approx 0.57$



## Convergence via two-curve method





the two-curve method can be used to illustrate convergence and divergence

- equation  $x = g(x)$  is re-expressed as a pair of equations:  $y_1 = x$ ,  $y_2 = g(x)$
- function  $y_1 = x$  and  $y_2 = g(x)$  are plotted
- roots of  $f(x) = 0$  correspond to the abscissa at the intersection of the two curves

### Iteration procedure

- start with initial guess  $x_0$
- determine the corresponding point on  $y_2$  curve:  $(x_0, g(x_0))$
- move horizontally to  $y_1$ : the point  $(x_1, x_1)$
- this corresponds to the first fixed-point iteration:  $x_1 = g(x_0)$
- repeat starting from  $x_1$  (moving to  $(x_1, g(x_1))$  and then to  $(x_2, x_2)$ )

# Convergence via two-curve method



- (a) and (b): solution converges — estimates move closer to the root
- (c) and (d): iterations diverge away from the root



# Convergence

fixed-point iteration converges locally (in a region of interest) if

$$|g'(x)| < 1$$

meaning the slope of  $y_2 = g(x)$  is less than the slope of  $y_1 = x$

- $0 < g'(x^*) < 1$ : monotone convergence
- $-1 < g'(x^*) < 0$ : oscillatory (spiral) convergence
- $|g'(x^*)| > 1$ : divergence

**Proof:** mean-value theorem with  $x_{i+1} = g(x_i)$ ,  $x^* = g(x^*)$ :

$$x^* - x_{i+1} = g(x^*) - g(x_i) = g'(\xi) (x^* - x_i)$$

thus for the true error  $E_{t,i} = x^* - x_i$ ,

$$E_{t,i+1} = g'(\xi) E_{t,i} \implies |g'(\xi)| < 1 \implies |E_{t,i+1}| < |E_{t,i}|$$

if converges, error decreases proportional to each step (linear convergence)

## MATLAB implementation of our example

```
clear,clc,format compact
g=@(x) exp(-x);
x0 = 0;
[xr,ea,iter] = fixpt(g,x0,1e-6);
X = ['The root = ',num2str(xr),' (ea = ',num2str(ea),'% in ',...
num2str(iter),' iterations)'];
disp(X)
```

with function

```
function [x1,ea,iter] = fixpt(g,x0,es,maxit)
% fixpt: fixed point root locator
if nargin < 2, error('at least 2 arguments required'), end
if nargin < 3||isempty(es),es = 1e-6;end % if es is blank set to 1e-6
if nargin < 4||isempty(maxit),maxit = 50;end % if maxit is blank set to 50
iter = 0; ea = 100;
while (1)
x1 = g(x0);
iter = iter + 1;
if x1 ~= 0, ea = abs((x1 - x0)/x1)*100; end
if (ea <= es || iter >= maxit), break, end
x0 = x1;
end
end
```

running code gives

The root = 0.56714 (ea = 7.6775e-07% in 35 iterations)

# Outline

- fixed point iteration
- **Newton-Raphson**
- secant method
- modified Newton-Raphson
- MATLAB `fzero` function

# Newton-Raphson method

- first-order Taylor approximation around  $x_i$


$$f(x) \approx \hat{f}(x) = f(x_i) + f'(x_i) (x - x_i)$$

- set to zero  $\hat{f}(x) = 0$ :

$$0 = f(x_i) + f'(x_i) (x - x_i)$$

- solution is our next estimate:

$$x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}$$



called *Newton-Raphson method* or just *Newton method*

## Example

use the Newton-Raphson method to estimate the root of

$$f(x) = e^{-x} - x$$

starting with an initial guess of  $x_0 = 0$  (true root:  $x^* \approx 0.56714329$ )

- first derivative:

$$f'(x) = -e^{-x} - 1$$

- iterative equation:

$$x_{i+1} = x_i - \frac{e^{-x_i} - x_i}{-e^{-x_i} - 1}$$

## Example

first iteration:

$$x_1 = x_0 - \frac{e^{-x_0} - x_0}{-e^{-x_0} - 1} = 0 - \frac{e^{-0} - 0}{-e^{-0} - 1} = 0.5$$

repeating gives the results

$i$	$x_i$	$\varepsilon_t(\%)$	$E_t$
0	0.000000000	100.0	0.567143290
1	0.500000000	11.8	0.067143290
2	0.566311003	0.147	0.000832287
3	0.567143165	0.0000221	0.000000125
4	0.567143290	$< 10^{-8}$	0.000000000

converges much faster than fixed-point iteration

## Error analysis of Newton-Raphson

second-order Taylor approximation:

$$f(x) = f(x_i) + f'(x_i)(x - x_i) + \frac{f''(\xi)}{2!}(x - x_i)^2$$

setting  $x = x^\star$  and using  $f(x^\star) = 0$  gives

$$0 = f(x_i) + f'(x_i)(x^\star - x_i) + \frac{f''(\xi)}{2!}(x^\star - x_i)^2$$

subtracting from Newton update  $0 = f(x_i) + f'(x_i)(x_{i+1} - x_i)$  gives

$$0 = f'(x_i)(x^\star - x_{i+1}) + \frac{f''(\xi)}{2!}(x^\star - x_i)^2$$

define error  $E_{t,i+1} = x^\star - x_{i+1}$  and assume convergence  $(x_i, \xi \rightarrow x^\star)$ ;; we get

$$E_{t,i+1} \approx -\frac{f''(x^\star)}{2f'(x^\star)} E_{t,i}^2$$

- error decreases quadratically (called *quadratic convergence*)
- number of correct digits roughly doubles with each iteration

## Example

examine error of Newton-Raphson for  $f(x) = e^{-x} - x$ , at  $x^{\star} = 0.56714329$

we have

$$f'(x^{\star}) = -1.56714329, \quad f''(x^{\star}) = 0.56714329$$

- error factor:  $E_{t,i+1} \approx -\frac{f''(x^{\star})}{2f'(x^{\star})} E_{t,i}^2 = 0.18095 E_{t,i}^2$
- initial error with  $x_0 = 0$ :  $E_{t,0} = 0.56714329$
- predicted errors using  $E_{t,i+1} \approx 0.18095 E_{t,i}^2$ :

$$E_{t,1} \approx 0.0582 \quad (\text{true: } 0.0671)$$

$$E_{t,2} \approx 8.16 \times 10^{-4} \quad (\text{true: } 8.32 \times 10^{-4})$$

$$E_{t,3} \approx 1.25 \times 10^{-7}$$

$$E_{t,4} \approx 2.84 \times 10^{-15}$$

- confirms quadratic convergence: error shrinks  $\propto E_{t,i}^2$



## Example


determine the positive root of  $f(x) = x^{10} - 1$  using Newton-Raphson with  $x_0 = 0.5$

### Newton-Raphson update

$$x_{i+1} = x_i - \frac{x_i^{10} - 1}{10x_i^9}$$

### Iterations


$i$	$x_i$	$ \varepsilon_a $ (%)
0	0.5	
1	51.65	99.032
2	46.485	11.111
3	41.8365	11.111
4	37.65285	11.111
$\vdots$	$\vdots$	$\vdots$
40	1.002316	2.130
41	1.000024	0.229
42	1	0.002




- very slow convergence
- first guess is in a region where the slope is near zero (flings the solution far away from the initial guess)

# Pitfalls of Newton-Raphson


- can be slow and may diverge
- *inflection points*: if  $f''(x) = 0$  near a root, iterations may diverge (figure a)
- *oscillations*: iterations can bounce around local maxima/minima (figure b)
- *near-zero slopes*: lead to very large jumps, possibly to other roots (figure c)
- *division by zero*: if  $f'(x) = 0$ , iteration fails completely (figure d)




(a)



(c)



(b)



(d)

# Convergence insights

- no universal convergence guarantee
- success depends on:
  - nature of the function
  - quality of the initial guess
- remedies:
  - choose initial guesses close to the root
  - use graphical insight or physical intuition
  - use small stepsize  $\alpha > 0$ :
$$x_{i+1} = x_i - \alpha \frac{f(x_i)}{f'(x_i)}$$
  - design software to detect slow convergence or divergence

## MATLAB implementation of Newton-Raphson

```
function [root,ea,iter] = newtraph(func,dfunc,xr,es,maxit,varargin)
% newtraph: Newton - Raphson root location zeroes
% input:
% func = name of function
% dfunc = name of derivative of function
% xr = initial guess
% es = desired relative error (default = 0.0001%)
% maxit = maximum allowable iterations (default = 50)
% p1,p2,... = additional parameters used by function
% output:
% root = real root
% ea = approximate relative error (%)
% iter = number of iterations
if nargin<3,error('at least 3 input arguments required'),end
if nargin<4 || isempty(es),es=0.0001;end
if nargin<5 || isempty(maxit),maxit = 50;end
iter = 0;
while (1)
xrold = xr;
xr = xr - func(xr)/dfunc(xr);
iter = iter + 1;
if xr ~ 0, ea = abs((xr - xrold)/xr) * 100; end
if ea <= es || iter >= maxit, break, end
end
root = xr;
```

## Example

find mass of bungee jumper with a drag coefficient 0.25 kg/m to have a velocity

$$v(t) = \sqrt{\frac{gm}{c_d}} \tanh\left(\sqrt{\frac{gc_d}{m}} t\right)$$

of 36 m/s after 4 s of free fall

the function to be evaluated is

$$f(m) = \sqrt{\frac{gm}{c_d}} \tanh\left(\sqrt{\frac{gc_d}{m}} t\right) - 36$$

the derivative of this function must be evaluated with respect to the unknown  $m$ :

$$\frac{df(m)}{dm} = \frac{1}{2} \sqrt{\frac{g}{mc_d}} \tanh\left(\sqrt{\frac{gc_d}{m}} t\right) - \frac{g}{2m} t \operatorname{sech}^2\left(\sqrt{\frac{gc_d}{m}} t\right)$$

code:

```
>> y = @(m) sqrt(9.81*m/0.25)*tanh(sqrt(9.81*0.25/m)*4)-36;
>> dy = @(m) 1/2*sqrt(9.81/(m*0.25))*tanh((9.81*0.25/m) ...
^(1/2)*4)-9.81/(2*m)*sech(sqrt(9.81*0.25/m)*4)^2;
>> newtraph(y,dy,140,0.00001)
ans =
142.7376
```

# Outline

- fixed point iteration
- Newton-Raphson
- **secant method**
- modified Newton-Raphson
- MATLAB `fzero` function


## Secant method

- some functions have derivatives that are difficult or inconvenient to evaluate
- the derivative can be approximated by a backward finite divided difference

$$f'(x_i) \approx \frac{f(x_{i-1}) - f(x_i)}{x_{i-1} - x_i}$$

- this approximation can be substituted into to yield the **secant method**:

$$x_{i+1} = x_i - \frac{f(x_i)(x_{i-1} - x_i)}{f(x_{i-1}) - f(x_i)}$$



requires two initial estimates

## Example

use the secant method on  $f(x) = e^{-x} - x$  with starting points  $x_{-1} = 0, x_0 = 1.0$   
true root  $\approx 0.56714329$

### First iteration

$$x_{-1} = 0 \quad f(x_{-1}) = 1.00000$$

$$x_0 = 1 \quad f(x_0) = -0.63212$$

so

$$x_1 = 1 - \frac{-0.63212(0 - 1)}{1 - (-0.63212)} = 0.61270 \quad \varepsilon_t = 8.0\%$$



## Example

### Second iteration

$$x_0 = 1 \quad f(x_0) = -0.63212$$

$$x_1 = 0.61270 \quad f(x_1) = -0.07081$$

note both estimates are now on the same side of the root; we have

$$x_2 = 0.61270 - \frac{-0.07081(1 - 0.61270)}{-0.63212 - (-0.07081)} = 0.56384 \quad \varepsilon_t = 0.58\%$$

### Third iteration

$$x_1 = 0.61270 \quad f(x_1) = -0.07081$$

$$x_2 = 0.56384 \quad f(x_2) = 0.00518$$

so

$$x_3 = 0.56384 - \frac{0.00518(0.61270 - 0.56384)}{-0.07081 - (-0.00518)} = 0.56717 \quad \varepsilon_t = 0.0048\%$$

## Secant method versus false-position method

false position

$$x_r = x_u - \frac{f(x_u)(x_l - x_u)}{f(x_l) - f(x_u)}$$

secant

$$x_{i+1} = x_i - \frac{f(x_i)(x_{i-1} - x_i)}{f(x_{i-1}) - f(x_i)}$$

- identical first iterate if  $x_u = x_0$  and  $x_l = x_{-1}$
- difference
  - false-position: always brackets the root by replacing the value with same sign as  $f(x_r)$
  - secant: updates sequentially ( $x_{i+1}$  replaces  $x_i$  and  $x_i$  replaces  $x_{i-1}$ )
- consequence
  - false-position: guaranteed convergence (root stays in bracket)
  - secant: may converge faster but can also diverge

## Example

use false-position and secant methods to estimate the root of  $f(x) = \ln x$

initial guesses:  $x_l = x_{-1} = 0.5$ ,  $x_u = x_0 = 5.0$

### False-position iterations

iteration	$x_l$	$x_u$	$x_r$
1	0.5	5.0	1.8546
2	0.5	1.8546	1.2163
3	0.5	1.2163	1.0585


converges to true root  $x = 1$

### Secant iterations

iteration	$x_{i-1}$	$x_i$	$x_{i+1}$
1	0.5	5.0	1.8546
2	5.0	1.8546	-0.10438

result: divergence


## Example



- for secant both values lie on same side of the root
- false-position is robust but slower because one endpoint stays fixed
- secant usually converges faster when it works

## Example: numerical comparison

comparison for  $f(x) = e^{-x} - x$



## Modified secant method

to avoid evaluating derivatives and two guesses

approximate derivative with fractional perturbation  $\delta$ :

$$f'(x_i) \approx \frac{f(x_i + \delta x_i) - f(x_i)}{\delta x_i}$$

iterative formula

$$x_{i+1} = x_i - \frac{\delta x_i f(x_i)}{f(x_i + \delta x_i) - f(x_i)}$$

called *modified secant method*

- $\delta$  must be chosen carefully
  - too small  $\rightarrow$  round-off errors due to subtractive cancellation
  - too large  $\rightarrow$  inefficiency or divergence
- useful when derivatives are difficult or inconvenient

## Example: modified secant

use modified secant method to solve  $f(x) = e^{-x} - x$ ,  $\delta = 0.01$ ,  $x_0 = 1.0$

- *first iteration:*

$$x_0 = 1, \quad f(x_0) = -0.63212$$

$$x_0 + \delta x_0 = 1.01, \quad f(x_0 + \delta x_0) = -0.64578$$

$$x_1 = 1 - \frac{0.01(-0.63212)}{-0.64578 - (-0.63212)} = 0.537263, \quad |\varepsilon_t| = 5.3\%$$

- *second iteration:*

$$x_1 = 0.537263, \quad f(x_1) = 0.047083$$

$$x_1 + \delta x_1 = 0.542635, \quad f(x_1 + \delta x_1) = 0.038579$$

$$x_2 = 0.537263 - \frac{0.005373(0.047083)}{0.038579 - 0.047083} = 0.56701, \quad |\varepsilon_t| = 0.0236\%$$

- *third iteration:*



$$x_2 = 0.56701, \quad f(x_2) = 0.000209$$

$$x_2 + \delta x_2 = 0.572680, \quad f(x_2 + \delta x_2) = -0.00867$$

$$x_3 = 0.56701 - \frac{0.00567(0.000209)}{-0.00867 - 0.000209} = 0.567143, \quad |\varepsilon_t| = 2.365 \times 10^{-5}\%$$

- method converges rapidly to true root 0.56714329

# Summary

Method	Formulation	Graphical Interpretation	Errors and Stopping Criteria
Newton-Raphson	$x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}$		Stopping criterion: $\left  \frac{x_{i+1} - x_i}{x_{i+1}} \right  100\% \leq e_s$ Error: $E_{i+1} = O(E_i^2)$
Secant	$x_{i+1} = x_i - \frac{f(x_i)(x_{i-1} - x_i)}{f(x_{i-1}) - f(x_i)}$		Stopping criterion: $\left  \frac{x_{i+1} - x_i}{x_{i+1}} \right  100\% \leq e_s$



# Outline

- fixed point iteration
- Newton-Raphson
- secant method
- **modified Newton-Raphson**
- MATLAB `fzero` function

## Multiple roots


a **multiple root** occurs when a function is tangent to the  $x$ -axis at the root

- example: double root (touches the axis at  $x = 1$  without crossing)


$$f(x) = (x - 3)(x - 1)(x - 1) = x^3 - 5x^2 + 7x - 3$$

- example: triple root (tangent and crosses the axis at  $x = 1$ )


$$f(x) = (x - 3)(x - 1)^3 = x^4 - 6x^3 + 12x^2 - 10x + 3$$



(a)



(b)



(c)

- rule of thumb:** even multiplicities touch but do not cross; odd multiplicities cross

## Why multiple roots are tricky?

1. **bracketing methods fail for even multiplicities:**  $f$  does not change sign, so bracketing tests may not detect the root
2. **derivative vanishes at the root:**  $f'(x^\star) = 0$  as well as  $f(x^\star) = 0$ 
  - Newton and secant divide by  $f' \Rightarrow$  possible division by values near zero
  - practical safeguard: check  $|f(x_i)|$  first and terminate when below tolerance (since  $f$  reaches 0 before  $f'$ )
3. **slower convergence:** standard Newton and secant become *linearly* convergent for multiple roots

# Modified Newton method

## Modified Newton method for known multiplicity

$$x_{i+1} = x_i - m \frac{f(x_i)}{f'(x_i)}$$

- $m$  is root multiplicity
- modify Newton to restore quadratic convergence
- requires prior knowledge of  $m$

**Modified Newton method:** applies Newton to  $u(x) = \frac{f(x)}{f'(x)}$

$$x_{i+1} = x_i - \frac{u(x_i)}{u'(x_i)} = x_i - \frac{f(x_i)f'(x_i)}{[f'(x_i)]^2 - f(x_i)f''(x_i)}$$

- $u(x) = \frac{f(x)}{f'(x)}$  has same roots as  $f(x)$
- regains fast (quadratic) convergence at multiple roots

## Example

use both the standard and modified Newton-Raphson to find the multiple root

$$f(x) = (x - 3)(x - 1)^2 = x^3 - 5x^2 + 7x - 3$$

- starting from  $x_0 = 0$
- we have

$$f'(x) = 3x^2 - 10x + 7, \quad f''(x) = 6x - 10$$

**(standard) Newton**

$$x_{i+1} = x_i - \frac{x_i^3 - 5x_i^2 + 7x_i - 3}{3x_i^2 - 10x_i + 7}$$

**modified Newton**

$$x_{i+1} = x_i - \frac{(x_i^3 - 5x_i^2 + 7x_i - 3)(3x_i^2 - 10x_i + 7)}{(3x_i^2 - 10x_i + 7)^2 - (x_i^3 - 5x_i^2 + 7x_i - 3)(6x_i - 10)}$$

## Example: results

### (standard) Newton

$i$	$x_i$	$ \varepsilon_t $ (%)
0	0.0000000	100
1	0.4285714	57
2	0.6857143	31
3	0.8328654	17
4	0.9133290	8.7
5	0.9557833	4.4
6	0.9776551	2.2

*behavior:* steady but linear approach to

$$x^{\star} = 1$$

### modified Newton

$i$	$x_i$	$ \varepsilon_t $ (%)
0	0.000000	100
1	1.105263	11
2	1.003082	0.31
3	1.000002	$2.4 \times 10^{-4}$

*behavior:* rapid (quadratic) approach to  
 $x^{\star} = 1$

## Example

to estimate root  $x = 3$ , we start at  $x_0 = 4$

**standard Newton**

$i$	$x_i$	$ \varepsilon_t $ (%)
0	4.000000	33
1	3.400000	13
2	3.100000	3.3
3	3.008696	0.29
4	3.000075	$2.5 \times 10^{-3}$
5	3.000000	$2 \times 10^{-7}$

**modified Newton**

$i$	$x_i$	$ \varepsilon_t $ (%)
0	4.000000	33
1	2.636364	12
2	2.820225	6.0
3	2.961728	1.3
4	2.998479	0.051
5	2.999998	$7.7 \times 10^{-5}$

- for a simple root, standard Newton is slightly more efficient
- and requires fewer computation (no  $f''$ )

## Modified secant method for multiple roots

- modified version of the secant method suited for multiple roots can also be developed
- apply secant method on  $u(x) = f(x)/f'(x)$
- the resulting formula is:

$$x_{i+1} = x_i - \frac{u(x_i) (x_{i-1} - x_i)}{u(x_{i-1}) - u(x_i)}$$



# Outline

- fixed point iteration
- Newton-Raphson
- secant method
- modified Newton-Raphson
- **MATLAB** `fzero` **function**

## MATLAB fzero function

fzero finds the real root of a single equation

### Basic syntax:

`fzero(function, x0)` or `fzero(function, [x0 x1])`

- `function`: function handle
- `x0`: initial guess or `x0`, `x1`: must bracket a root

**Example:**  $x^2 - 9 = 0$

- negative root:

`x = fzero(@(x) x^2-9,-4)`

- positive root:

`x = fzero(@(x) x^2-9,4)`

- using bracketing:

`x = fzero(@(x) x^2-9,[0 4])`

**Note:** if no sign change occurs in  $[x_0, x_1]$ , MATLAB returns an error

## More on MATLAB fzero

### Extended syntax:

```
[x,fx] = fzero(function,x0,options,p1,p2,...)
```

- `x` : computed root
- `fx` : function value at the root
- `options` : structure created using `optimset`
- `p1,p2,...` : extra parameters required by the function
- if parameters are passed but options are not used: pass `[]` in its place

### `optimset` syntax:

```
options = optimset('par1',val1,'par2',val2,...)
```

- we can list all available parameters via typing `optimset`
- commonly used with `fzero`:
  - `display = 'iter'` : shows iteration details
  - `tolx` : termination tolerance for  $x$

## Example

display result for finding root of  $f(x) = x^{10} - 1$

```
>> options = optimset('display','iter');  
>> [x,fx] = fzero(@(x) x^10 - 1,0.5,options)
```

Func-count	x	f(x)	Procedure
1	0.5	-0.999023	initial
2	0.485858	-0.999267	search
3	0.514142	-0.998709	search
4	0.48	-0.999351	search
5	0.52	-0.998554	search
6	0.471716	-0.999454	search

.  
.

Zero found in the interval: [-0.14, 1.14].

```
x =  
1  
fx =  
0
```

set tolerance

```
>> options = optimset ('tolx', 1e-3);  
>> [x,fx] = fzero(@(x) x^10 - 1,0.5,options)  
x =  
1.0009  
fx =  
0.0090
```

## References and further readings

- S. C. Chapra and R. P. Canale. *Numerical Methods for Engineers* (8th edition). McGraw Hill, 2021. (Ch.6)
- S. C. Chapra. *Applied Numerical Methods with MATLAB for Engineers and Scientists* (5th edition). McGraw Hill, 2023. (Ch.6)