

VECTOR FIELD:

$$F(x, y) = f(x, y)i + g(x, y)j \rightarrow 2 \text{ space}$$

$$F(x, y, z) = f(x, y, z)i + g(x, y, z)j + h(x, y, z)k \rightarrow 3 \text{ space}$$

Inverse square field.

$$F(r) = \frac{c}{\|r\|^3} \cdot r, \quad r \text{ is radius vector and } c \text{ is a constant}$$

$$\therefore F(x, y) = \frac{c}{(x^2 + y^2)^{3/2}} \cdot (xi + yj), \quad \begin{matrix} r_2 = xi + yj \\ r_3 = xi + yj + zk \end{matrix}$$

$$\therefore F(x, y, z) = \frac{c}{(x^2 + y^2 + z^2)^{3/2}} \cdot (xi + yj + zk)$$

Inverse square fields are conservative in any region that does not contain the origin.

E.g. $\phi(x, y) = -\frac{c}{(x^2 + y^2)^{1/2}} \rightarrow$ is a potential function

$$\nabla \phi(x, y) = \frac{\partial \phi}{\partial x} i + \frac{\partial \phi}{\partial y} j$$

$$= \frac{cx}{(x^2 + y^2)^{3/2}} i + \frac{cy}{(x^2 + y^2)^{3/2}} j$$

$$= \frac{c}{(x^2 + y^2)^{3/2}} (xi + yj)$$

$$= F(x, y)$$

$$\begin{aligned} &= \frac{c}{(x^2 + y^2)^{3/2}} (xi + yj) \\ &= \frac{c}{(x^2 + y^2)^{3/2}} (xi + yj) \end{aligned}$$

Ex. Show that divergence of inverse square field

$$F(x, y, z) = \frac{c}{(x^2 + y^2 + z^2)^{3/2}} (xi + yj + zk) \text{ is zero.}$$

Let $r = (x^2 + y^2 + z^2)^{1/2} \Rightarrow (x^2 + y^2 + z^2)^{1/2} = r \Rightarrow \boxed{r^3}$

$$F(x, y, z) = \frac{c}{r^3} (xi + yj + zk)$$

$$F(x, y, z) = \frac{cx}{r^3} i + \frac{cy}{r^3} j + \frac{cz}{r^3} k$$

$$\text{div } F = \nabla \cdot F = \left(\frac{\partial}{\partial x} i + \frac{\partial}{\partial y} j + \frac{\partial}{\partial z} k \right) \cdot \left(\frac{cx}{r^3} i + \frac{cy}{r^3} j + \frac{cz}{r^3} k \right)$$

$$\Rightarrow \left(\frac{\partial}{\partial x} i + \frac{\partial}{\partial y} j + \frac{\partial}{\partial z} k \right) \cdot \left[c \left(\frac{x}{r^3} i + \frac{y}{r^3} j + \frac{z}{r^3} k \right) \right]$$

$$\Rightarrow c \left[\frac{\partial}{\partial x} \left(\frac{x}{r^3} \right) + \frac{\partial}{\partial y} \left(\frac{y}{r^3} \right) + \frac{\partial}{\partial z} \left(\frac{z}{r^3} \right) \right] = \text{!}$$

$$r = (x^2 + y^2 + z^2)^{1/2}$$

$$\frac{\partial r}{\partial x} = \frac{1}{2} (x^2 + y^2 + z^2)^{-1/2} \cdot 2x$$

$$\frac{\partial r}{\partial y} = \frac{1}{2} (x^2 + y^2 + z^2)^{-1/2} \cdot 2y$$

$$\frac{\partial r}{\partial x} = \frac{1}{r} \cdot x$$

$$\frac{\partial r}{\partial y} = \frac{y}{r}$$

$$\frac{\partial r}{\partial x} = \frac{x}{r}$$

$$\frac{\partial r}{\partial z} = \frac{z}{r}$$

$$\frac{\partial r}{\partial z} = \frac{z}{r}$$

$$\text{---} \textcircled{!} \Rightarrow C \left[\frac{\partial}{\partial x} \left(\frac{x}{r^3} \right) + \frac{\partial}{\partial y} \left(\frac{y}{r^3} \right) + \frac{\partial}{\partial z} \left(\frac{z}{r^3} \right) \right]$$

$$\frac{\partial}{\partial x} \left(\frac{x}{r^3} \right) \because \frac{u}{v} = v \cdot du - u \cdot dv / v^2$$

$$= \frac{r^3 \cdot (1) \frac{\partial}{\partial x} - x \cdot 3r^2 \left(\frac{\partial r}{\partial x} \right)}{(r^3)^2} = \frac{r^3 - 3r^2 x \left(\frac{x}{r} \right)}{r^6}$$

$$= \frac{r^3 - 3rx^2}{r^6} \Rightarrow \frac{r^3}{r^6} - \frac{3rx^2}{r^6} \Rightarrow \frac{1}{r^3} - \frac{3x^2}{r^5}$$

$$\frac{\partial}{\partial x} \left(\frac{x}{r^3} \right) = \frac{1}{r^3} - \frac{3x^2}{r^5}$$

$$\frac{\partial}{\partial y} \left(\frac{y}{r^3} \right) = \frac{1}{r^3} - \frac{3y^2}{r^5}$$

$$\frac{\partial}{\partial z} \left(\frac{z}{r^3} \right) = \frac{1}{r^3} - \frac{3z^2}{r^5}$$

$$\text{div } F = C \left[\frac{1}{r^3} - \frac{3x^2}{r^5} + \frac{1}{r^3} - \frac{3y^2}{r^5} + \frac{1}{r^3} - \frac{3z^2}{r^5} \right]$$

$$\text{div } F = C \left[\frac{3}{r^3} - \frac{(3x^2 + 3y^2 + 3z^2)}{r^5} \right]$$

$$\text{div } F = C \left[\frac{3}{r^3} - \frac{3(x^2 + y^2 + z^2)}{r^5} \right] \because (x^2 + y^2 + z^2)^{1/2} = r$$

$$\text{div } F = C \left[\frac{3}{r^3} - \frac{3r^2}{r^5} \right] \Rightarrow C \left[\frac{3}{r^3} - \frac{3}{r^3} \right] \Rightarrow C(0)$$

$$\boxed{\text{div } F = 0} \text{ proved.}$$

Q. Find the divergence & curl of following vector field.

$$P(x, y, z) = xz^3i + 2y^4x^2j + 5z^2yk.$$

$$\therefore \text{div } F = \nabla \cdot F = \left(\frac{\partial}{\partial x} i + \frac{\partial}{\partial y} j + \frac{\partial}{\partial z} k \right) \cdot (xz^3i + 2y^4x^2j + 5z^2yk)$$

$$\text{div } F = \frac{\partial}{\partial x}(xz^3) + \frac{\partial}{\partial y}(2y^4x^2) + \frac{\partial}{\partial z}(5z^2y).$$

$$\text{div } F = z^3 + 8y^3x^2 + 10zy - \text{scalar} \checkmark$$

$$\text{Curl} = \nabla \times F = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{vmatrix} \quad \begin{array}{l} F_1 = xz^3 \\ F_2 = 2y^4x^2 \\ F_3 = 5z^2y \end{array}$$

$$\nabla \times F = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ xz^3 & 2y^4x^2 & 5z^2y \end{vmatrix}$$

$$\nabla \times F = i \left\{ \frac{\partial}{\partial y}(5z^2y) - \frac{\partial}{\partial z}(2y^4x^2) \right\} - j \left\{ \frac{\partial}{\partial x}(5z^2y) - \frac{\partial}{\partial z}(xz^3) \right\} + k \left\{ \frac{\partial}{\partial x}(2y^4x^2) - \frac{\partial}{\partial y}(xz^3) \right\}$$

$$\nabla \times F = i(5z^2 - 0) - j(0 - 3xz^2) + k(4xy^4 - 0)$$

$$\nabla \times F = 5z^2i + 3xz^3j + 4xy^4k - \text{vector} \checkmark$$

Q. Find the divergence & curl of the vector field.

$$P(x, y, z) = x^2yi + 2y^3zj + 3zk.$$

$$\therefore \text{div } F = \nabla \cdot F = \left(\frac{\partial}{\partial x} i + \frac{\partial}{\partial y} j + \frac{\partial}{\partial z} k \right) \cdot (x^2yi + 2y^3zj + 3zk)$$

$$\text{div } F = \frac{\partial}{\partial x}(x^2y) + \frac{\partial}{\partial y}(2y^3z) + \frac{\partial}{\partial z}(3z)$$

$$\text{div } F = 2xy + 6y^2z + 3 - \text{scalar} \checkmark$$

$$\text{curl} = \nabla \times F = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x^2y & 2y^3z & 3z \end{vmatrix}$$

$$\nabla \times F = i \left\{ \frac{\partial}{\partial y}(3z) - \frac{\partial}{\partial z}(2y^3z) \right\} - j \left\{ \frac{\partial}{\partial x}(3z) - \frac{\partial}{\partial z}(x^2y) \right\} + k \left\{ \frac{\partial}{\partial x}(2y^3z) - \frac{\partial}{\partial y}(x^2y) \right\}$$

$$\nabla \times F = i(0 - 2y^3) - j(0 - 0) + k(0 - x^2)$$

$$\nabla \times F = -2y^3i - x^2k - \text{vector} \checkmark$$

Vector field again

Q Confirm that " ϕ " is a potential function or not.

(i) $\phi(x, y) = 2y^2 + 3x^2y - xy^3$

$F(x, y) = 2xi + 6yj + 8zk$

Q

If $F = \nabla\phi$, it is a potential function else not.

$$\therefore \nabla\phi = \left(\frac{\partial}{\partial x}i + \frac{\partial}{\partial y}j\right)\phi(x, y)$$

$$= \left(\frac{\partial}{\partial x}i + \frac{\partial}{\partial y}j\right)(2y^2 + 3x^2y - xy^3)$$

$$= \frac{\partial}{\partial x}i(2y^2 + 3x^2y - xy^3) + \frac{\partial}{\partial y}j(2y^2 + 3x^2y - xy^3)$$

$$\nabla\phi = (0 + 6xy - y^3)i + (4y + 3x^2 - 3xy^2)j$$

Since $F \neq \nabla\phi$, ϕ is not a potential function

(ii) $\phi(x, y, z) = x \sin z + y \sin x + z \sin y$

Q $F(x, y, z) = (\sin z + y \cos x)i + (\sin x + z \cos y)j + (\sin y + x \cos z)k$

$$\therefore \nabla\phi = \left(\frac{\partial}{\partial x}i + \frac{\partial}{\partial y}j + \frac{\partial}{\partial z}k\right)\phi(x, y, z)$$

$$= \frac{\partial}{\partial x}(x \sin z + y \sin x + z \sin y)i + \frac{\partial}{\partial y}(x \sin z + y \sin x + z \sin y)j + \frac{\partial}{\partial z}(x \sin z + y \sin x + z \sin y)k$$

$$= (\sin z + y \cos x + 0)i + (0 + \sin x + z \cos y)j + (\cos z + 0 + \sin y)k$$

$$\nabla \phi = (\sin z + y \cos x)i + (\sin x + z \cos y)j + (\sin y + \cos z)k$$

Since $F = \nabla \phi$, it is a potential function ✓

Conservative or Non-Conservative fields.

$$F(x, y) = f(x, y)i + g(x, y)j$$

$$\text{if } \frac{\partial f}{\partial y} = \frac{\partial g}{\partial x} \quad (\text{conservative})$$

else non-conservative

it is valid for 2-space.

$$F(x, y, z) = f(x, y, z)i + g(x, y, z)j + h(x, y, z)k$$

$$\text{if } \nabla \times F = 0 \quad (\text{conservative})$$

else non-conservative.

for 3-space.

Q. Whether the field is conservative or not.

$$(i) F(x, y) = 2xy^3 i + (1 + 3x^2 y^2) j$$

$$f(x, y) = 2x^4 y^3$$

$$g(x, y) = 1 + 3x^2 y^2$$

$$\frac{\partial f}{\partial y} = 6xy^2$$

$$\frac{\partial g}{\partial x} = 6xy^2$$

$\frac{\partial f}{\partial y} = \frac{\partial g}{\partial x}$, it is a conservative vector field.