

# 1 | FORCING

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

The following lemma is from [?]

**Lemma 1.1.** *Let  $\lambda$  be an infinite cardinal,  $\mathcal{M}$  a transitive model with  $\text{On} \subseteq \mathcal{M}$ ,  $\mathbb{P} \in \mathcal{M}$  a forcing notion and  $g \subseteq \mathbb{P}$  an  $\mathcal{M}$ -generic filter. Then  $V \models {}^\lambda \mathcal{M} \subseteq \mathcal{M}$  implies that  $V[g] \models {}^\lambda \mathcal{M} \subseteq \mathcal{M}$ .*

PROOF. Work in  $V[g]$ . Let  $c := \langle c_\alpha \mid \alpha < \lambda \rangle$  be a  $\lambda$ -sequence such that  $c_\alpha \in \mathcal{M}[g]$  for every  $\alpha < \lambda$ . Fix for every  $\alpha < \lambda$  a  $\mathbb{P}$ -name  $\dot{c}_\alpha$  such that  $\dot{c}_\alpha^g = c_\alpha$ . Also let  $\dot{a}$  be a  $\mathbb{P}$ -name with  $\dot{a}^g = \langle \dot{c}_\alpha \mid \alpha < \lambda \rangle$  and choose  $p \in g$  such that

$$V \models \ulcorner p \Vdash \forall \alpha < \check{\lambda}: \dot{a}(\alpha) \in \mathcal{M}^{\mathbb{P}^\ulcorner} \urcorner.$$

Now, working in  $V$ , there is for each  $\alpha < \lambda$  a maximal antichain  $A_\alpha$  below  $p$  such that every  $q \in A_\alpha$  decides  $\dot{a}(\alpha)$ ; i.e.,  $q \Vdash \ulcorner \dot{a}(\alpha) = \check{x} \urcorner$  for some  $x \in \mathcal{M}$ .

Define now

$$\sigma := \{((\alpha, x), q) \mid \alpha \in \lambda \wedge q \in A_\alpha \wedge q \Vdash \dot{a}(\alpha) = \check{x}\}.$$

Then  $p \Vdash \sigma = \dot{a}^\frown$ . Note that  $|\sigma| \leq \lambda$ , since  $|A_\alpha| \leq \lambda$  for each  $\alpha < \lambda$ . Thus  $\sigma \in \mathcal{M}$ .

Now, going back to  $V[g]$  again, it holds that  $\langle \dot{c}_\alpha \mid \alpha < \lambda \rangle = \dot{a}^g = \sigma^g \in \mathcal{M}[g]$ . But we can compute  $c = \langle c_\alpha \mid \alpha < \lambda \rangle = \langle \dot{c}_\alpha^g \mid \alpha < \lambda \rangle$  from  $\langle \dot{c}_\alpha \mid \alpha < \lambda \rangle$  and  $g$ , so that  $c \in \mathcal{M}[g]$  by Replacement. ■