

# 1 | VIRTUAL LARGE CARDINALS

In this chapter we investigate the properties of virtual versions of well-known large cardinals, including measurables, strongs, supercompacts, woodins and vopěnkas. This entails firstly analysing the relationships between them, and secondly looking at more general properties in terms of their behaviour in canonical inner models as well as their indestructibility. This virtual perspective also allows us to analyse virtualised versions of large cardinals that are otherwise inconsistent with ZFC, such as the berkeley cardinals.

## 1.1 STRONGS & SUPERCOMPACTS

We start out with measurables, strongs and supercompacts. Their (non-virtual) definitions can be found in Appendix ??.

**Definition 1.1.** Let  $\theta$  be a regular uncountable cardinal. Then a cardinal  $\kappa < \theta$  is...

- **faintly  $\theta$ -measurable** if, in a forcing extension, there is a transitive class  $\mathcal{N}$  and an elementary embedding  $\pi: H_\theta^V \rightarrow \mathcal{N}$  with  $\text{crit } \pi = \kappa$ ;
- **faintly  $\theta$ -strong** if it's faintly  $\theta$ -measurable,  $H_\theta^V \subseteq \mathcal{N}$  and  $\pi(\kappa) > \theta$ ;
- **faintly  $\theta$ -supercompact** if it's faintly  $\theta$ -measurable,  ${}^{<\theta}\mathcal{N} \subseteq \mathcal{N}$  and  $\pi(\kappa) > \theta$ .

We further replace “faintly” by **virtually** when  $\mathcal{N} \subseteq V$ , we attach a “**pre**” if we don't assume that  $\pi(\kappa) > \theta$ , and we will leave out  $\theta$  when it holds for all regular  $\theta > \kappa$ . ◦

As a quick example of this terminology, a *faintly prestrong cardinal* is a cardinal  $\kappa$  such that for all regular  $\theta > \kappa$ ,  $\kappa$  is faintly  $\theta$ -measurable with  $H_\theta^V \subseteq \mathcal{N}$ .

We note that even small cardinals can be faintly measurable: we may for instance have a precipitous ideal on  $\omega_1$ ; see [?, Theorem 22.33]. The

## 1.1. STRONGS & SUPERCOMPACTS VIRTUAL LARGE CARDINALS

“virtually” adverb implies that the cardinals are in fact large cardinals in the usual sense, as Proposition 1.2 below shows.

**Proposition 1.2** (Virtualised folklore). *For any regular uncountable cardinal  $\theta$ , every virtually  $\theta$ -measurable cardinal is 1-iterable.*

PROOF. Let  $\kappa$  be virtually  $\theta$ -measurable, witnessed by a forcing  $\mathbb{P}$ , a transitive  $\mathcal{N} \subseteq V$  and an elementary  $\pi: H_\theta^V \rightarrow \mathcal{N}$  with  $\pi \in V^\mathbb{P}$ . If  $\kappa$  isn’t a strong limit then we have a surjection  $\pi(f): \mathcal{P}(\alpha) \rightarrow \pi(\kappa)$  with  $\text{ran } \pi(f) = \text{ran } f \subseteq \kappa$  for some  $\alpha < \kappa$ ,  $\nless \theta$ . Note that we used  $\mathcal{N} \subseteq V$  to ensure that  $\mathcal{P}(\alpha)^V = \mathcal{P}(\alpha)^\mathcal{N}$ . The same argument shows that  $\kappa$  is regular. By restricting the generic embedding and using that  $\mathcal{P}(\kappa)^V = \mathcal{P}(\kappa)^\mathcal{N}$  as  $\mathcal{N} \subseteq V$  and  $\mathcal{P}(\kappa)^V \subseteq \mathcal{N}$ , we get that  $\kappa$  is 1-iterable. ■

Along with the above definition of faintly supercompactness we can also virtualise Magidor’s characterisation of supercompact cardinals<sup>1</sup>, which was one of the original characterisations of the remarkable cardinals in [?].

**Definition 1.3.** Let  $\theta$  be a regular uncountable cardinal. Then a cardinal  $\kappa < \theta$  is **virtually  $\theta$ -Magidor-supercompact** if there are  $\bar{\kappa} < \bar{\theta} < \kappa$  and a generic elementary  $\pi: H_\theta^V \rightarrow H_{\bar{\theta}}^V$  such that  $\text{crit } \pi = \bar{\kappa}$  and  $\pi(\bar{\kappa}) = \kappa$ . ◦

In the virtual world these two versions of supercompacts remain equivalent, but they also turn out to be equivalent to the virtually strong:

**Theorem 1.4** (Gitman-Schindler). *For an uncountable cardinal  $\kappa$ , the following are equivalent.*<sup>2</sup>

- (i)  $\kappa$  is virtually strong;
- (ii)  $\kappa$  is virtually supercompact;
- (iii)  $\kappa$  is virtually Magidor-supercompact.

PROOF. (ii)  $\Rightarrow$  (i) is simply by definition.

<sup>1</sup>See Appendix ?? for the non-virtual version of this characterisation.

<sup>2</sup>A cardinal satisfying any/all of these conditions is usually called **remarkable**.

(i)  $\Rightarrow$  (iii): Fix  $\theta > \kappa$ . By (i) there exists a generic elementary embedding  $\pi: H_{(2^{<\theta})^+}^V \rightarrow \mathcal{M}$  with<sup>3</sup>  $\text{crit } \pi = \kappa$ ,  $\pi(\kappa) > \theta$ ,  $H_{(2^{<\theta})^+}^V \subseteq \mathcal{M}$  and  $\mathcal{M} \subseteq V$ . Since  $H_\theta^V, H_{\pi(\theta)}^\mathcal{M} \in \mathcal{M}$ , Countable Embedding Absoluteness ?? implies that  $\mathcal{M}$  has a generic elementary embedding  $\pi^*: H_\theta^V \rightarrow H_{\pi(\theta)}^\mathcal{M}$  with  $\text{crit } \pi^* = \kappa$  and  $\pi^*(\kappa) = \pi(\kappa) > \theta$ . Since  $H_\theta^V = H_\theta^\mathcal{M}$  as  $\mathcal{M} \subseteq V$  and  $H_\theta^V \subseteq \mathcal{M}$ , elementarity of  $\pi$  now implies that  $H_{(2^{<\theta})^+}^V$  has ordinals  $\bar{\kappa} < \bar{\theta} < \kappa$  and a generic elementary  $\sigma: H_\theta^V \rightarrow H_{\bar{\theta}}^V$  with  $\text{crit } \sigma = \bar{\kappa}$  and  $\sigma(\bar{\kappa}) = \kappa$ . This shows (iii).

(iii)  $\Rightarrow$  (ii): Fix  $\theta > \kappa$  and  $\delta := (2^{<\theta})^+$ . By (iii) there exist ordinals  $\bar{\kappa} < \bar{\delta} < \kappa$  and a generic elementary embedding  $\pi: H_{\bar{\delta}}^V \rightarrow H_\delta^V$  with  $\text{crit } \pi = \bar{\kappa}$  and  $\pi(\bar{\kappa}) = \kappa$ . We will argue that  $\bar{\kappa}$  is virtually  $\bar{\theta}$ -supercompact in  $H_{\bar{\delta}}^V$ , so that by elementarity  $\kappa$  is virtually  $\theta$ -supercompact in  $H_\delta^V$  and hence also in  $V$  by the choice of  $\delta$ . Consider the restriction

$$\sigma := \pi \upharpoonright H_{\bar{\theta}}^V: H_{\bar{\theta}}^V \rightarrow H_\theta^V.$$

Note that  $H_\theta^V$  is closed under  $<\bar{\theta}$ -sequences (and more) in  $V$ . Now define

$$X := \bar{\theta}+1 \cup \{x \in H_\theta^V \mid \exists y \in H_{\bar{\theta}}^V \exists p \in \text{Col}(\omega, H_{\bar{\theta}}^V): p \Vdash \dot{\sigma}(\dot{y}) = \dot{x}\} \in V.$$

Note that  $|X| = |H_{\bar{\theta}}^V| = 2^{<\bar{\theta}}$  and that  $\text{ran } \sigma \subseteq X$ . Now let  $\overline{\mathcal{M}} \prec H_\theta^V$  be such that  $X \subseteq \overline{\mathcal{M}}$  and  $\overline{\mathcal{M}}$  is closed under  $<\bar{\theta}$ -sequences. Note that we can find such an  $\overline{\mathcal{M}}$  of size  $(2^{<\bar{\theta}})^{<\bar{\theta}} = 2^{<\bar{\theta}}$ . Let  $\mathcal{M}$  be the transitive collapse of  $\overline{\mathcal{M}}$ , so that  $\mathcal{M}$  is still closed under  $<\bar{\theta}$ -sequences and we still have that  $|\mathcal{M}| = 2^{<\bar{\theta}} < \bar{\delta}$ , making  $\mathcal{M} \in H_{\bar{\delta}}^V$ .

Countable Embedding Absoluteness ?? then implies that  $H_{\bar{\delta}}^V$  has a generic elementary embedding  $\sigma^*: H_{\bar{\theta}}^V \rightarrow \mathcal{M}$  with  $\text{crit } \sigma^* = \bar{\kappa}$ , showing that  $\bar{\kappa}$  is virtually  $\bar{\theta}$ -supercompact in  $H_{\bar{\delta}}^V$ , which is what we wanted to show. ■

*Remark 1.5.* The above proof in fact shows something stronger: if  $\kappa$  is virtually  $(2^{<\theta})^+$ -strong then it is virtually  $\theta$ -supercompact, and if it's virtually

<sup>3</sup>The domain of  $\pi$  is  $H_{(2^{<\theta})^+}^V$  to ensure that  $H_\theta^V \in \text{dom } \pi$ .

### 1.1. STRONGS & SUPERCOMPACTS VIRTUAL LARGE CARDINALS

$(2^{<\theta})^+$ -Magidor-supercompact then it's virtually  $\theta$ -supercompact. It's open whether they are equivalent level-by-level (see Question ??).

A key difference between the normal large cardinals and the virtual kind is that we don't have a virtual version of the Kunen inconsistency<sup>4</sup>: it's perfectly valid to have a generic elementary embedding  $H_\theta^V \rightarrow H_\theta^V$  with  $\theta$  much larger than the critical point.

**Proposition 1.6** (Folklore). *If  $0^\sharp$  exists then there are inaccessible cardinals  $\kappa < \theta$  such that, in a generic extension of  $L$ , there is an elementary embedding  $\pi: L_\theta \rightarrow L_\theta$ . In other words,  $\pi$  witnesses a strong failure of the virtualised Kunen inconsistency.*

PROOF. From  $0^\sharp$  we get an elementary embedding  $j: L \rightarrow L$ . Let  $C \subseteq \text{On}$  be the proper class club of limit points of  $j$  above  $\text{crit } j$ , which then contains an inaccessible cardinal  $\theta$  as there are stationarily many such. Restrict  $j$  to  $\pi := j \upharpoonright L_\theta \rightarrow \mathcal{N}$  and note that  $\mathcal{N} = L_\theta$  by condensation and because  $\theta$  is a limit point of  $j$ . Let  $\kappa := \text{crit } \pi$ . Now an application of Countable Embedding Absoluteness ?? shows that a generic extension of  $L$  contains an elementary embedding  $\tilde{\pi}: L_\theta \rightarrow L_\theta$  with  $\text{crit } \tilde{\pi} = \kappa$ . ■

This becomes important when dealing with the “pre”-versions of the large cardinals. We next move to a virtualisation of the  $\alpha$ -superstrong cardinals.

**Definition 1.7** (N.). Let  $\theta$  be a regular uncountable cardinal and  $\alpha$  an ordinal. Then a cardinal  $\kappa < \theta$  is **faintly  $(\theta, \alpha)$ -superstrong** if it's faintly  $\theta$ -measurable,  $H_\theta^V \subseteq \mathcal{N}$  and  $\pi^\alpha(\kappa) \leq \theta^5$ . We replace “faintly” by **virtually** when  $\mathcal{N} \subseteq V$ , we say that  $\kappa$  is **faintly  $\alpha$ -superstrong** if it's faintly  $(\theta, \alpha)$ -superstrong for *some*  $\theta$ , and lastly  $\kappa$  is simply **faintly superstrong** if it is faintly 1-superstrong.<sup>6</sup> ◦

<sup>4</sup>See Appendix ?? for the Kunen inconsistency.

<sup>5</sup>Here we set  $\pi^\alpha(\kappa) := \sup_{\xi < \alpha} \pi^\xi(\kappa)$  when  $\alpha$  is a limit ordinal.

<sup>6</sup>Note that the conventions stated here are different from the ones in Definition 1.1.

As in the non-virtual case, the virtually superstrongs supercede the virtually strongs in consistency strength. Note that this then also implies that the superstrongs are stronger than the virtually supercompacts, which is *not* the case outside the virtual world.

**Proposition 1.8** (N.). *If  $\kappa$  is faintly superstrong then  $H_\kappa$  has a proper class of virtually strong cardinals.*

PROOF. Fix a regular  $\theta > \kappa$  and a generic embedding  $\pi: H_\theta^V \rightarrow \mathcal{N}$  with  $\text{crit } \pi = \kappa$ ,  $H_\theta^V \subseteq \mathcal{N}$  and  $\pi(\kappa) < \theta$ . Then  $\pi(\kappa)$  is a  $V$ -cardinal, so that  $H_{\pi(\kappa)}^V$  thinks that  $\kappa$  is virtually strong. This implies that  $H_\kappa^V$  thinks there is a proper class of virtually strong cardinals, using that  $H_\kappa^V \prec H_{\pi(\kappa)}^V$ . ■

The following theorem then shows that the only thing stopping prestrongness from being equivalent to strongness is the existence of “Kunen inconsistencies”.

**Theorem 1.9** (N.). *Let  $\theta$  be an uncountable cardinal. Then a cardinal  $\kappa < \theta$  is virtually  $\theta$ -prestrong iff either*

- (i)  $\kappa$  is virtually  $\theta$ -strong; or
- (ii)  $\kappa$  is virtually  $(\theta, \omega)$ -superstrong.

PROOF.  $(\Leftarrow)$  is trivial, so we show  $(\Rightarrow)$ . Let  $\kappa$  be virtually  $\theta$ -prestrong. Assume (i) fails, meaning that there’s a generic extension  $V^\mathbb{P}$  and an elementary embedding  $\pi \in V^\mathbb{P}$  such that  $\pi: H_\theta^V \rightarrow \mathcal{N}$  for some transitive  $\mathcal{N}$  with  $H_\theta^V \subseteq \mathcal{N}$ ,  $\mathcal{N} \subseteq V$ ,  $\text{crit } \pi = \kappa$  and  $\pi(\kappa) \leq \theta$ . Assume  $\pi^n(\kappa)$  is defined for all  $n < \omega$  and define  $\lambda := \sup_{n < \omega} \pi^n(\kappa)$ . If  $\lambda \leq \theta$  then  $\kappa$  is virtually  $(\theta, \omega)$ -superstrong by definition, so assume that there’s some least  $n < \omega$  such that  $\pi^{n+1}(\kappa) > \theta$ .

This means that  $\kappa$  is virtually  $\nu$ -strong for every regular  $\nu \in (\kappa, \pi^n(\kappa))$ , which is a  $\Delta_0$ -statement in  $\{H_\nu^V\}$  and hence downwards absolute to  $H_{\pi^n(\kappa)}^V$ . This means that  $\kappa$  is virtually strong in  $H_{\pi^n(\kappa)}^V$  and also that  $\pi^n(\kappa)$  is virtually strong in  $H_{\pi^{n+1}(\kappa)}^V$  by elementarity, and so in particular virtually  $\theta$ -

### 1.1. STRONGS & SUPERSTRONGS VIRTUAL LARGE CARDINALS

strong in  $\mathcal{N}$ . This means that there's some generic elementary embedding

$$\sigma: H_\theta^\mathcal{N} \rightarrow \mathcal{M}$$

with  $H_\theta^\mathcal{N} \subseteq \mathcal{M}$ ,  $\mathcal{M} \subseteq \mathcal{N}$ ,  $\text{crit } \sigma = \pi^n(\kappa)$  and  $\sigma(\pi^n(\kappa)) > \theta$ . We can now restrict  $\sigma$  to its critical point  $\pi^n(\kappa)$  to get that

$$H_{\pi^n(\kappa)}^V = H_{\pi^n(\kappa)}^\mathcal{N} \prec H_{\sigma(\pi^n(\kappa))}^\mathcal{M},$$

using that  $H_\theta^V = H_\theta^\mathcal{N}$  holds as  $\pi$  is a virtual embedding. Since  $\kappa$  is virtually strong in  $H_{\pi^n(\kappa)}^V$  this means that  $\kappa$  is also virtually strong in  $H_{\sigma(\pi^n(\kappa))}^\mathcal{M}$ . In particular,  $\kappa$  is virtually  $\theta$ -strong in  $\mathcal{M}$ , and as  $H_\theta^\mathcal{M} = H_\theta^\mathcal{N} = H_\theta^V$ , this means that  $\kappa$  is virtually  $\theta$ -strong in  $V$ , contradicting (i). ■

We then get the following consistency result.

**Corollary 1.10** (N.). *For any uncountable regular  $\theta$ , the existence of a virtually  $\theta$ -strong cardinal is equiconsistent with the existence of a faintly  $\theta$ -measurable cardinal.*

PROOF. The above Proposition 1.8 and Theorem 1.9 show that virtually  $\theta$ -prestrongs are equiconsistent with virtually  $\theta$ -strongs. Now note that Countable Embedding Absoluteness ?? and condensation in  $L$  imply that every faintly  $\theta$ -measurable cardinal is virtually  $\theta$ -prestrong in  $L$ . ■

Recall that a cardinal  $\kappa$  is **virtually rank-into-rank** if there exists a cardinal  $\theta > \kappa$  and a generic elementary embedding  $\pi: H_\theta^V \rightarrow H_\theta^V$  with  $\text{crit } \pi = \kappa$ . We firstly note that the virtually  $\omega$ -superstrongs coincide with the virtually rank-into-ranks.

**Proposition 1.11** (N.). *A regular uncountable cardinal  $\kappa$  is virtually  $\omega$ -superstrong iff it is virtually rank-into-rank.*

PROOF. If  $\kappa$  is virtually  $\omega$ -superstrong, witnessed by a generic embedding  $\pi: H_\theta^V \rightarrow \mathcal{N}$ , then  $\lambda := \sup_{n < \omega} \pi^n(\kappa)$  is well-defined. By restricting  $\pi$  to  $\pi \upharpoonright H_\lambda^V: H_\lambda^V \rightarrow H_\lambda^V$  we get a witness to  $\kappa$  being virtually  $\lambda$ -rank-into-rank.

Conversely, if  $\kappa$  is  $\theta$ -rank-into-rank, witnessed by a generic embedding  $\pi: H_\theta^V \rightarrow H_\theta^V$ , then one readily checks that  $\pi$  also witnesses that  $\kappa$  is virtually  $\omega$ -superstrong.  $\blacksquare$

## 1.2 WOODINS & VOPĚNKAS

In this section we will analyse the virtualisations of the woodin and vopěnka cardinals, which can be seen as “boldface” variants of strongs and supercompacts.

**Definition 1.12.** Let  $\theta$  be a regular uncountable cardinal. Then a cardinal  $\kappa < \theta$  is **faintly**  $(\theta, A)$ -**strong** for a set  $A \subseteq H_\theta^V$  if there exists a generic elementary embedding

$$\pi: (H_\theta^V, \in, A) \rightarrow (\mathcal{M}, \in, B)$$

with  $\mathcal{M}$  transitive, such that  $\text{crit } \pi = \kappa$ ,  $\pi(\kappa) > \theta$ ,  $H_\theta^V \subseteq \mathcal{M}$  and  $B \cap H_\theta^V = A$ . We say that  $\kappa$  is **faintly**  $(\theta, A)$ -**supercompact** if we further have that  ${}^{<\theta}\mathcal{M} \cap V \subseteq \mathcal{M}$  and say that  $\kappa$  is **faintly**  $(\theta, A)$ -**extendible** if  $\mathcal{M} = H_\mu^V$  for some  $V$ -cardinal  $\mu$ . We will leave out  $\theta$  if it holds for all regular  $\theta > \kappa$ .  $\circ$

**Definition 1.13.** A cardinal  $\delta$  is **faintly woodin** if given any  $A \subseteq H_\delta^V$  there exists a faintly  $(<\delta, A)$ -strong cardinal  $\kappa < \delta$ .  $\circ$

As with the previous definitions, for both of the above two definitions we substitute “faintly” for **virtually** when  $\mathcal{M} \subseteq V$ , and substitute “strong”, “supercompact” and “woodin” for **prestrong**, **presupercompact** and **prewoodin** when we don’t require that  $\pi(\kappa) > \theta$ .

We note in the following proposition that, in analogy with the real woodin cardinals, virtually woodin cardinals are mahlo. This contrasts the virtually prewoodins since [?], together with Theorem 1.20 below, shows that they can be singular.

**Proposition 1.14** (Virtualised folklore). *Virtually woodin cardinals are mahlo.*

PROOF. Let  $\delta$  be virtually woodin. Note that  $\delta$  is a limit of weakly compact cardinals by Proposition 1.2, making  $\delta$  a strong limit. As for regularity, assume that we have a cofinal increasing function  $f: \alpha \rightarrow \delta$  with  $f(0) > \alpha$  and  $\alpha < \delta$ , and note that  $f$  cannot have any closure points. Fix a virtually  $(<\delta, f)$ -strong cardinal  $\kappa < \delta$ ; we claim that  $\kappa$  is a closure point for  $f$ , which will yield our desired contradiction.

Let  $\gamma < \kappa$  and choose a regular  $\theta \in (f(\gamma), \delta)$ . We then have a generic embedding  $\pi: (H_\theta^V, \in, f \cap H_\theta^V) \rightarrow (\mathcal{N}, \in, f^+)$  with  $H_\theta^V \subseteq \mathcal{N}$ ,  $\mathcal{N} \subseteq V$ ,  $\text{crit } \pi = \kappa$ ,  $\pi(\kappa) > \theta$  and  $f^+$  is a function such that  $f^+ \cap H_\theta^V = f \cap H_\theta^V$ . But then  $f^+(\gamma) = f(\gamma) < \pi(\kappa)$  by our choice of  $\theta$ , so elementarity implies that  $f(\gamma) < \kappa$ , making  $\kappa$  a closure point for  $f$ ,  $\nmid$ . This shows that  $\delta$  is inaccessible.

As for mahloness, let  $C \subseteq \delta$  be a club and  $\kappa < \delta$  a virtually  $(<\delta, C)$ -strong cardinal. Let  $\theta \in (\min C, \delta)$  and let  $\pi: H_\theta^V \rightarrow \mathcal{N}$  be the associated generic elementary embedding. Then for every  $\gamma < \kappa$  there exists an element of  $C$  below  $\pi(\kappa)$ , namely  $\min C$ , so by elementarity  $\kappa$  is a limit of elements of  $C$ , making it an element of  $C$ . As  $\kappa$  is regular, this shows that  $\delta$  is mahlo. ■

The well-known equivalence of the “function definition” and “ $A$ -strong” definition of woodin cardinals<sup>7</sup> holds if we restrict ourselves to *virtually* woodins, and the analogue of the equivalence between virtually strongs and virtually supercompacts allows us to strengthen this:

**Proposition 1.15** (Dimopoulos-Gitman-N.). *For an uncountable cardinal  $\delta$ , the following are equivalent.*

- (i)  $\delta$  is virtually woodin.
- (ii) for every  $A \subseteq H_\delta^V$  there exists a virtually  $(<\delta, A)$ -supercompact  $\kappa < \delta$ .
- (iii) for every  $A \subseteq H_\delta^V$  there exists a virtually  $(<\delta, A)$ -extendible  $\kappa < \delta$ .

<sup>7</sup>See Appendix ?? for this characterisation of (non-virtual) woodin cardinals.






Figure 1.1: Proof strategy of Proposition 1.15, dotted lines are trivial implications.

- (iv) for every function  $f: \delta \rightarrow \delta$  there are regular cardinals  $\kappa < \theta < \delta$ , where  $\kappa$  is a closure point for  $f$ , and a generic elementary  $\pi: H_\theta^V \rightarrow \mathcal{M}$  such that  $\text{crit } \pi = \kappa$ ,  $H_\theta^V \subseteq \mathcal{M}$ ,  $\mathcal{M} \subseteq V$  and  $\theta = \pi(f \restriction \kappa)(\kappa)$ .
- (v) for every function  $f: \delta \rightarrow \delta$  there are regular cardinals  $\kappa < \theta < \delta$ , where  $\kappa$  is a closure point for  $f$ , and a generic elementary  $\pi: H_\theta^V \rightarrow \mathcal{M}$  such that  $\text{crit } \pi = \kappa$ ,  ${}^{<\pi(f)(\kappa)}\mathcal{M} \subseteq \mathcal{M}$ ,  $\mathcal{M} \subseteq V$  and  $\theta = \pi(f \restriction \kappa)(\kappa)$ .
- (vi) for every function  $f: \delta \rightarrow \delta$  there are regular cardinals  $\bar{\theta} < \kappa < \theta < \delta$ , where  $\kappa$  is a closure point for  $f$ , and a generic elementary embedding  $\pi: H_{\bar{\theta}}^V \rightarrow H_\theta^V$  with  $\pi(\text{crit } \pi) = \kappa$ ,  $f(\text{crit } \pi) = \bar{\theta}$  and  $f \restriction \kappa \in \text{ran } \pi$ .

PROOF. Firstly note that (iii)  $\Rightarrow$  (ii)  $\Rightarrow$  (i) and (v)  $\Rightarrow$  (iv) are simply by definition.

(i)  $\Rightarrow$  (iv) Assume  $\delta$  is virtually woodin, and fix a function  $f: \delta \rightarrow \delta$ . Let  $\kappa < \delta$  be virtually  $(<\delta, f)$ -strong and let  $\theta := \sup_{\alpha < \kappa} f(\alpha) + 1$ . Then there's a generic elementary embedding  $\pi: (H_\theta^V, \in, f \restriction H_\theta^V) \rightarrow (\mathcal{M}, \in, f^+)$  where  $f^+ \restriction \kappa = f \restriction \kappa$ ,  $\mathcal{M} \subseteq V$  and  $\pi(\kappa) > \theta$ . We firstly want to show that  $\kappa$  is a closure point for  $f$ , so let  $\alpha < \kappa$ . Then

$$f(\alpha) = f^+(\alpha) = \pi(f)(\alpha) = \pi(f)(\pi(\alpha)) = \pi(f(\alpha)),$$

so  $\pi$  fixes  $f(\alpha)$  for every  $\alpha < \kappa$ . Now, if  $\kappa$  wasn't a closure point for  $f$  then, letting  $\alpha < \kappa$  be the least such that  $f(\alpha) \geq \kappa$ ,

$$\theta > f(\alpha) = \pi(f(\alpha)) > \theta,$$

a contradiction. Note that we used that  $\pi(\kappa) > \theta$  here, so this argument wouldn't work if we had only assumed  $\delta$  to be virtually prewoodin. Lastly,  $\theta$ -strongness implies that  $H_\theta^V \subseteq \mathcal{M}$ , and  $\mathcal{M} \subseteq V$  holds by assumption.

$(iv) \Rightarrow (vi)$  Assume  $(iv)$  holds, let  $f: \delta \rightarrow \delta$  be given and define  $g: \delta \rightarrow \delta$  as  $g(\alpha) := (2^{<f(\alpha)})^+$ . By  $(iv)$  there's a  $\kappa < \delta$  which is a closure point of  $g$  and there's a regular  $\theta \in (\kappa, \delta)$  and a generic elementary  $\pi: H_\theta^V \rightarrow \mathcal{M}$  with  $\text{crit } \pi = \kappa$ ,  $H_\theta^V \subseteq \mathcal{M}$ ,  $\mathcal{M} \subseteq V$  and  $\theta = \pi(f \upharpoonright \kappa)(\kappa)$ . We want to find a regular  $\bar{\theta} < \kappa$  and another elementary embedding  $\sigma: H_{\bar{\theta}}^V \rightarrow H_\theta^V$  with  $\sigma(\text{crit } \sigma) = \kappa$ ,  $f(\text{crit } \sigma) = \bar{\theta}$  and  $f \upharpoonright \kappa \in \text{ran } \sigma$ .

Note that  $\mathcal{M} \subseteq V$  and  $H_\theta^V \subseteq \mathcal{M}$  implies that  $H_\theta^V = H_\theta^\mathcal{M}$ , so that both  $H_\theta^V$  and  $H_{\pi(\theta)}^\mathcal{M}$  are elements of  $\mathcal{M}$  (we introduced  $g$  to ensure that  $\pi(\theta)$  makes sense). An application of Countable Embedding Absoluteness ?? then yields that  $\mathcal{M}$  has a generic elementary embedding  $\pi^*: H_\theta^\mathcal{M} \rightarrow H_{\pi(\theta)}^\mathcal{M}$  such that  $\text{crit } \pi^* = \kappa$ ,  $\pi^*(\kappa) = \pi(\kappa)$  and  $\pi(f \upharpoonright \kappa) \in \text{ran } \pi^*$ .

By elementarity of  $\pi$ ,  $H_\theta^V$  has an ordinal  $\bar{\theta} < \kappa$  and a generic elementary embedding  $\sigma: H_{\bar{\theta}}^V \rightarrow H_\theta^V$  with  $\sigma(\text{crit } \sigma) = \kappa$ ,  $f \upharpoonright \kappa \in \text{ran } \sigma$  and  $\bar{\theta} = f(\text{crit } \sigma)$ , which is what we wanted to show.

$(vi) \Rightarrow (v)$  Assume  $(vi)$  holds and let  $f: \delta \rightarrow \delta$  be given. Define  $g: \delta \rightarrow \delta$  as  $g(\alpha) := (2^{<f(\alpha)})^+$ , so that by  $(vi)$  there exist regular  $\bar{\kappa} < \bar{\theta} < \kappa < \theta$  such that  $\kappa$  is a closure point for  $g$  and there exists a generic elementary embedding  $\pi: H_{\bar{\theta}}^V \rightarrow H_\theta^V$  with  $\text{crit } \pi = \bar{\kappa}$ ,  $\pi(\bar{\kappa}) = \kappa$ ,  $g(\bar{\kappa}) = \bar{\theta}$  and  $g \upharpoonright \kappa \in \text{ran } \pi$ .

Now, following the  $(iii) \Rightarrow (ii)$  direction in the proof of Theorem 1.4 we get a transitive  $\mathcal{M} \in H_{g(\bar{\kappa})}^V$  closed under  $<f(\bar{\kappa})$ -sequences, and  $H_{g(\bar{\kappa})}^V$  has a generic elementary embedding  $\sigma: H_{f(\bar{\kappa})}^V \rightarrow \mathcal{M}$  with  $\text{crit } \sigma = \bar{\kappa}$  and  $\sigma(\bar{\kappa}) = \kappa > f(\bar{\kappa})$ . In other words,  $\bar{\kappa}$  is virtually  $f(\bar{\kappa})$ -supercompact in  $H_{\bar{\theta}}^V$ . Elementarity of  $\pi$  then implies that  $\kappa$  is virtually  $\pi(f)(\bar{\kappa})$ -supercompact in  $H_\theta^V$ , which is what we wanted to show.

$(vi) \Rightarrow (iii)$  Let  $C$  be the club of all  $\alpha$  such that  $(H_\alpha^V, \in, A \cap H_\alpha^V) \prec (H_\delta^V, \in, A)$ . Let  $f: \delta \rightarrow \delta$  be given as  $f(\alpha) = \langle \alpha_0, \alpha_1 \rangle$  with  $\langle -, - \rangle$  being the Gödel pairing function, where  $\alpha_0$  is the first limit of elements of  $C$  above  $\alpha$  and the  $\alpha_1$ 's are chosen such that  $\{\alpha_1 \mid \alpha < \beta\}$  encodes  $A \cap \beta$ . This definition makes sense since  $\delta$  is inaccessible by Proposition 1.2.

Let  $\kappa < \delta$  be a closure point of  $f$  such that there are regular cardinals  $\bar{\theta} < \kappa$ ,  $\theta > \kappa$  and a generic elementary embedding  $\pi: H_{\bar{\theta}}^V \rightarrow H_{\theta}^V$  such that  $\pi(\text{crit } \pi) = \kappa$ ,  $f(\text{crit } \pi) = \bar{\theta}$ , and  $f \restriction \kappa \in \text{ran } \pi$ . We claim that  $\bar{\kappa} := \text{crit } \pi$  is virtually  $(< \delta, A)$ -extendible. To see this, it suffices by the definition of  $C$  to show that

$$(H_{\kappa}^V, \in, A \cap H_{\kappa}^V) \models \ulcorner \bar{\kappa} \text{ is virtually } (A \cap H_{\kappa})\text{-extendible} \urcorner, \quad (1)$$

since  $\kappa \in C$  because it is a closure point of  $f$ . Let  $\beta := \min(C - \bar{\kappa}) < \bar{\theta}$  and note that  $\beta$  exists as  $f(\bar{\kappa}) = \bar{\theta}$  so the definition of  $f$  says that  $\bar{\theta}$  is a limit of elements of  $C$  above  $\bar{\kappa}$ . It then holds that  $(H_{\kappa}^V, \in, A \cap H_{\kappa}^V) \prec (H_{\beta}^V, \in, A \cap H_{\beta}^V)$  as both  $\bar{\kappa}$  and  $\beta$  are elements of  $C$ . Since  $f$  encodes  $A$  in the manner previously described and  $\pi^{-1}(f) \restriction \bar{\kappa} = f \restriction \bar{\kappa}$ , we get that  $\pi(A \cap H_{\kappa}^V) = A \cap H_{\kappa}^V$  and thus

$$(H_{\kappa}^V, \in, A \cap H_{\kappa}^V) \prec (H_{\pi(\beta)}^V, \in, A^*) \quad (2)$$

for  $A^* := \pi(A \cap H_{\beta}^V)$ . Now, as  $(H_{\gamma}^V, \in, A \cap H_{\gamma}^V)$  and  $(H_{\pi(\gamma)}^V, \in, A^* \cap H_{\pi(\gamma)}^V)$  are elements of  $H_{\pi(\beta)}^V$  for every  $\gamma < \kappa$ , Countable Embedding Absoluteness ?? implies that  $H_{\pi(\beta)}^V$  sees that  $\bar{\kappa}$  is virtually  $(< \kappa, A^*)$ -extendible, which by (2) then implies (1), which is what we wanted to show. ■

*Remark 1.16.* The above proof shows that the  $\mathcal{M} \subseteq V$  assumptions can be replaced by “sufficient” agreement between  $\mathcal{M}$  and  $V$ : for (i)-(iii) this means that  $H_{\theta}^{\mathcal{M}} = H_{\theta}^V$  whenever  $\mathcal{M}$  is the codomain of a virtual  $(\theta, A)$ -strong/supercompact/extendible embedding, and in (iv)-(v) this means that  $H_{\pi(f)(\kappa)}^{\mathcal{M}} = H_{\pi(f)(\kappa)}^V$ . The same thing holds in the “lightface” setting of Theorem 1.4.

We will now step away from the woodins for a little bit, and introduce the vopěnkas. In anticipation of the next section we will work with the

class-sized version here, but all the following results work equally well for inaccessible virtually vopěnka cardinals<sup>8</sup>.

**Definition 1.17** (GBC). The **Generic Vopěnka Principle** (gVP) states that for any class  $C$  consisting of structures in a common language, there are distinct  $\mathcal{M}, \mathcal{N} \in C$  and a generic elementary embedding  $\pi: \mathcal{M} \rightarrow \mathcal{N}$ . ◦

We will be using a standard variation of gVP involving the following *natural sequences*.

**Definition 1.18** (GBC). Say that a class function  $f: \text{On} \rightarrow \text{On}$  is an **indexing function** if it satisfies that  $f(\alpha) > \alpha$  and  $f(\alpha) \leq f(\beta)$  for all  $\alpha < \beta$ . ◦

**Definition 1.19** (GBC). Say that an On-sequence  $\langle \mathcal{M}_\alpha \mid \alpha < \text{On} \rangle$  is **natural** if there exists an indexing function  $f: \text{On} \rightarrow \text{On}$  and unary relations  $R_\alpha \subseteq V_{f(\alpha)}$  such that  $\mathcal{M}_\alpha = (V_{f(\alpha)}, \in, \{\alpha\}, R_\alpha)$  for every  $\alpha$ . Denote this indexing function by  $f^{\vec{\mathcal{M}}}$  and the unary relations as  $R_\alpha^{\vec{\mathcal{M}}}$ . ◦

The following Theorem 1.20 is then the main theorem of this section. Firstly it shows that inaccessible cardinals are virtually vopěnka iff they are virtually prewoodin, but also that adding the “virtually” adverb doesn’t do anything in this context, in contrast to Theorem 1.46.

**Theorem 1.20** (GBC, Dimopoulos-Gitman-N.). *The following are equivalent:*

- (i) *gVP holds;*
- (ii) *For any natural On-sequence  $\vec{\mathcal{M}}$  there exists a generic elementary embedding  $\pi: \mathcal{M}_\alpha \rightarrow \mathcal{M}_\beta$  for some  $\alpha < \beta$ ;*
- (iii) *On is virtually prewoodin;*
- (iv) *On is faintly prewoodin.*

<sup>8</sup>Note however that we have to require inaccessibility here: see [?] for an analysis of the singular virtually vopěnka cardinals.

Redundant; mention this properly above

PROOF. (i)  $\Rightarrow$  (ii) and (iii)  $\Rightarrow$  (iv) are trivial.

(iv)  $\Rightarrow$  (i): Assume On is faintly prewoodin and fix some On-sequence  $\vec{\mathcal{M}} := \langle \mathcal{M}_\alpha \mid \alpha < \text{On} \rangle$  of structures in a common language. Let  $\kappa$  be  $(<\text{On}, \vec{\mathcal{M}})$ -prestrong and fix some regular  $\theta > \kappa$  satisfying that  $\mathcal{M}_\alpha \in H_\theta^V$  for every  $\alpha < \theta$ , and fix a generic elementary embedding

$$\pi: (H_\theta^V, \in, \vec{\mathcal{M}}) \rightarrow (\mathcal{N}, \in, \mathcal{M}^*)$$

with  $H_\theta^V \subseteq \mathcal{N}$  and  $\vec{\mathcal{M}} \cap H_\theta^V = \mathcal{M}^* \cap H_\theta^V$ . Set  $\kappa := \text{crit } \pi$ .

We have that  $\pi \restriction \mathcal{M}_\kappa: \mathcal{M}_\kappa \rightarrow \mathcal{M}_{\pi(\kappa)}^*$ , but we need to reflect this embedding down below  $\theta$  as we don't know whether  $\mathcal{M}_{\pi(\kappa)}^*$  is on the  $\vec{\mathcal{M}}$  sequence. Working in the generic extension, we have

$$\mathcal{N} \models \exists \bar{\kappa} < \pi(\kappa) \exists \dot{\sigma} \in V^{\text{Col}(\omega, \mathcal{M}_{\bar{\kappa}}^*)}: \ulcorner \dot{\sigma}: \mathcal{M}_{\bar{\kappa}}^* \rightarrow \mathcal{M}_{\pi(\kappa)}^* \text{ is elementary} \urcorner.$$

Here  $\kappa$  realises  $\bar{\kappa}$  and  $\pi \restriction \mathcal{M}_\kappa$  realises  $\sigma$ . Note that  $\mathcal{M}_\kappa^* = \mathcal{M}_\kappa$  since we ensured that  $\mathcal{M}_\kappa \in H_\theta^V$  and we are assuming that  $\vec{\mathcal{M}} \cap H_\theta^V = \mathcal{M}^* \cap H_\theta^V$ , so the domain of  $\sigma (= \pi \restriction \mathcal{M}_\kappa)$  is  $\mathcal{M}_\kappa^*$  — also note that  $\sigma$  exists in a  $\text{Col}(\omega, \mathcal{M}_\kappa)$  extension of  $\mathcal{N}$  by an application of Countable Embedding Absoluteness ???. Now elementarity of  $\pi$  implies that

$$H_\theta^V \models \exists \bar{\kappa} < \kappa \exists \dot{\sigma} \in V^{\text{Col}(\omega, \mathcal{M}_{\bar{\kappa}})}: \ulcorner \dot{\sigma}: \mathcal{M}_{\bar{\kappa}} \rightarrow \mathcal{M}_\kappa \text{ is elementary} \urcorner,$$

which is upwards absolute to  $V$ , from which we can conclude that  $\sigma: \mathcal{M}_{\bar{\kappa}} \rightarrow \mathcal{M}_\kappa$  witnesses that gVP holds.

(ii)  $\Rightarrow$  (iii): Assume (ii) holds and assume that On is not virtually prewoodin, which means that there exists some class  $A$  such that there are no virtually  $A$ -prestrong cardinals. This allows us to define a function  $f: \text{On} \rightarrow \text{On}$  as  $f(\alpha)$  being the least regular  $\eta > \alpha$  such that  $\alpha$  is not virtually  $(\eta, A)$ -prestrong.

We also define  $g: \text{On} \rightarrow \text{On}$  as taking  $\alpha$  to the least strong limit cardinal above  $\alpha$  which is a closure point for  $f$ . Note that  $g$  is an indexing function, so we can let  $\vec{\mathcal{M}}$  be the natural sequence induced by  $g$  and  $R_\alpha := A \cap H_{g(\alpha)}^V$ .

(ii) supplies us with  $\alpha < \beta$  and a generic elementary embedding<sup>9</sup>

$$\pi: (H_{g(\alpha)}^V, \in, A \cap H_{g(\alpha)}^V) \rightarrow (H_{g(\beta)}^V, \in, A \cap H_{g(\beta)}^V).$$

Since  $g(\alpha)$  is a closure point for  $f$  it holds that  $f(\text{crit } \pi) < g(\alpha)$ , so fixing a regular  $\theta \in (f(\text{crit } \pi), g(\alpha))$  we get that  $\text{crit } \pi$  is virtually  $(\theta, A)$ -prestrong, contradicting the definition of  $f$ . Hence  $\text{On}$  is virtually prewoodin. ■

### 1.2.1 Weak Vopěnka

Scrap this section?

We now move to a *weak* variant of  $\text{gVP}$ , introduced in a category-theoretic context in [?]. It starts with the following equivalent characterisation of  $\text{gVP}$ , which is the virtual analogue of the characterisation shown in [?].

**Lemma 1.21** (GBC, Virtualised Adámek-Rosický).  *$\text{gVP}$  is equivalent to there not existing an  $\text{On}$ -sequence of first-order structures  $\langle \mathcal{M}_\alpha \mid \alpha < \text{On} \rangle$  satisfying that<sup>10</sup>*

- (i)  $\text{gVP}$
- (ii) *There is not a natural  $\text{On}$ -sequence  $\langle \mathcal{M}_\alpha \mid \alpha < \text{On} \rangle$  satisfying that*
  - *there is a generic homomorphism  $\mathcal{M}_\alpha \rightarrow \mathcal{M}_\beta$  for every  $\alpha \leq \beta$ , which is unique in all generic extensions;*
  - *there is no generic homomorphism  $\mathcal{M}_\beta \rightarrow \mathcal{M}_\alpha$  for any  $\alpha < \beta$ .*
- (iii) *There is not a natural  $\text{On}$ -sequence  $\langle \mathcal{M}_\alpha \mid \alpha < \text{On} \rangle$  satisfying that*
  - *there is a homomorphism  $\mathcal{M}_\alpha \rightarrow \mathcal{M}_\beta$  in  $V$  for every  $\alpha \leq \beta$ , which is unique in all generic extensions;*
  - *there is no generic homomorphism  $\mathcal{M}_\beta \rightarrow \mathcal{M}_\alpha$  for any  $\alpha < \beta$ .*

PROOF. Note that the only difference between (ii) and (iii) is that the homomorphism exists in  $V$ , making (ii)  $\Rightarrow$  (iii) trivial.

(iii)  $\Rightarrow$  (i): Assume that  $\text{gVP}$  fails, meaning by Theorem 1.20 that we have a natural  $\text{On}$ -sequence  $\vec{\mathcal{M}}_\alpha$  such that, in every generic extension, there's

<sup>9</sup>Note that  $V_{g(\alpha)} = H_{g(\alpha)}^V$  since  $g(\alpha)$  is a strong limit cardinal.

<sup>10</sup>This is equivalent to saying that  $\text{On}$ , viewed as a category, can't be fully embedded into the category  $\mathbf{Gra}$  of graphs, which is how it's stated in [?].

no homomorphism between any two distinct  $\mathcal{M}_\alpha$ 's. Define an On-sequence  $\langle \mathcal{N}_\kappa \mid \kappa \in \text{Card} \rangle$  as

$$\mathcal{N}_\kappa := \prod_{\xi \leq \kappa} \mathcal{M}_\xi = \{(x, \xi) \mid \xi \leq \kappa \wedge \xi \in \text{Card} \wedge x \in \mathcal{M}_\xi\},^{11}$$

with a unary relation  $R^*$  given as  $R^*(x, \xi)$  iff  $\mathcal{M}_\xi \models R(x)$  and a binary relation  $\sim^*$  given as  $(x, \xi) \sim^* (x', \xi')$  iff  $\xi = \xi'$ . Whenever we have a homomorphism  $f: \mathcal{N}_\kappa \rightarrow \mathcal{N}_\lambda$  we then get an induced homomorphism  $\tilde{f}: \mathcal{M}_0 \rightarrow \mathcal{M}_\xi$ , given as  $\tilde{f}(x) := f(x, 0)$ , where  $\xi \leq \kappa$  is given by preservation of  $\sim^*$ .

For any two cardinals  $\kappa < \lambda$  we have a homomorphism  $j_{\kappa\lambda}: \mathcal{N}_\kappa \rightarrow \mathcal{N}_\lambda$  in  $V$ , given as  $j_{\kappa\lambda}(x, \xi) := (x, \xi)$ . This embedding must also be the *unique* such embedding in all generic extensions, as otherwise we get a generic homomorphism between two distinct  $\mathcal{M}_\alpha$ 's. Furthermore, there can't be any homomorphism  $\mathcal{N}_\lambda \rightarrow \mathcal{N}_\kappa$  as that would also imply the existence of a generic homomorphism between two distinct  $\mathcal{M}_\alpha$ 's.

(i)  $\Rightarrow$  (ii): Assume that we have an On-sequence  $\vec{\mathcal{M}}_\alpha$  as in the theorem, with generic homomorphisms  $j_{\alpha\beta}: \mathcal{M}_\alpha \rightarrow \mathcal{M}_\beta$  that are unique in all generic extensions for every  $\alpha \leq \beta$ , with no generic homomorphisms going the other way.

We first note that we can for every  $\alpha \leq \beta$  choose the  $j_{\alpha\beta}$  in a  $\text{Col}(\omega, \mathcal{M}_\alpha)$ -extension, by a proof similar to the proof of Lemma ?? and using the uniqueness of  $j_{\alpha\beta}$ . Next, fix a proper class  $C \subseteq \text{On}$  such that  $\alpha \in C$  implies that

$$\sup_{\xi \in C \cap \alpha} |\mathcal{M}_\xi|^V < |\mathcal{M}_\alpha|^V.$$

and note that this implies that  $V[g] \models |\mathcal{M}_\xi| < |\mathcal{M}_\alpha|$  for every  $V$ -generic  $g \subseteq \text{Col}(\omega, \mathcal{M}_\xi)$ . This means that for every  $\alpha \in C$  we may choose some  $\eta_\alpha \in \mathcal{M}_\alpha$  which is *not* in the range of any  $j_{\xi\alpha}$  for  $\xi < \alpha$ . But now define first-order structures  $\langle \mathcal{N}_\alpha \mid \alpha \in C \rangle$  as  $\mathcal{N}_\alpha := (\mathcal{M}_\alpha, \eta_\alpha)$ . Then, by our assumption on the  $\mathcal{M}_\alpha$ 's and construction of the  $\mathcal{N}_\alpha$ 's, there can be no generic homomorphism between any two distinct  $\mathcal{N}_\alpha$ , showing that gVP fails. ■

Note that the proof of the above lemma shows that we without loss of generality may assume that the generic homomorphism in (i) exists in  $V$ , which we record here:

**Lemma 1.22** (GBC, Virtualised Adámek-Rosický). *gVP is equivalent to there not existing an On-sequence of first-order structures  $\langle \mathcal{M}_\alpha \mid \alpha < \text{On} \rangle$  satisfying that<sup>12</sup>*

- (i) *there is a homomorphism  $\mathcal{M}_\alpha \rightarrow \mathcal{M}_\beta$  in  $V$  for every  $\alpha \leq \beta$ , which is unique in all generic extensions;*
- (ii) *there is no generic homomorphism  $\mathcal{M}_\beta \rightarrow \mathcal{M}_\alpha$  for any  $\alpha < \beta$ .*

■

The *weak* version of gVP is then simply “flipping the arrows around” in the above characterisation of gVP.

**Definition 1.23** (GBC). **Generic Weak Vopěnka’s Principle** (gWVP) states that there does *not* exist an On-sequence of first-order structures  $\langle \mathcal{M}_\alpha \mid \alpha < \text{On} \rangle$  such that

- there is a generic homomorphism  $\mathcal{M}_\beta \rightarrow \mathcal{M}_\alpha$  for every  $\alpha \leq \beta$ , which is unique in all generic extensions;
- there is *no* generic homomorphism  $\mathcal{M}_\alpha \rightarrow \mathcal{M}_\beta$  for any  $\alpha < \beta$ .

○

Denoting the corresponding non-generic principle by WVP [?] showed the following.

**Theorem 1.24** (Wilson). *WVP is equivalent to On being a Woodin cardinal.*

Given our 1.20 we may then suspect that in the virtual world these two are equivalent, which indeed turns out to be the case. We will be roughly following the argument in [?], but we have to diverge from it at several points in which they’re using the fact that they’re working with class-sized elementary embeddings.

---

<sup>12</sup>This is equivalent to saying that On, viewed as a category, can’t be fully embedded into the category **Gra** of graphs, which is how it’s stated in [?].



Indeed, in that paper they establish a correspondence between elementary embeddings and certain homomorphisms, a correspondence we won't achieve here. Proving that the elementary embeddings we *do* get are non-trivial seems to furthermore require extra assumptions on our structures. Let's begin.

Define for every strong limit cardinal  $\lambda$  and  $\Sigma_1$ -formula  $\varphi$  the relations

$$\begin{aligned} R^\varphi &:= \{x \in V \mid (V, \in) \models \varphi[x]\} \\ R_\lambda^\varphi &:= \{x \subseteq H_\lambda^V \mid \exists y \in R^\varphi: y \cap H_\lambda^V = x\} \end{aligned}$$

and given any class  $A$  define the structure

$$\mathcal{P}_{\lambda,A} := (H_{\lambda+}^V, R_\lambda^\varphi, \{\lambda\}, A \cap H_\lambda^V)_{\varphi \in \Sigma_1}.$$

Say that a homomorphism  $h: \mathcal{P}_{\lambda,A} \rightarrow \mathcal{P}_{\eta,A}$  is **trivial** if  $h(x) \cap H_\eta^V = x \cap H_\eta^V$  for every  $x \in H_{\lambda+}^V$ . Note that  $h$  can only be trivial if  $\eta \leq \lambda$  since  $h(\lambda) = \eta$ .

**Lemma 1.25** (GBC, Gitman-N.). *Let  $\lambda$  be a singular strong limit cardinal,  $\eta$  a strong limit cardinal and  $A \subseteq V$  a class. If there exists a non-trivial generic homomorphism  $h: \mathcal{P}_{\lambda,A} \rightarrow \mathcal{P}_{\eta,A}$  then there's a non-trivial generic elementary embedding*

$$\pi: (H_{\lambda+}^V, \in, A \cap H_\lambda^V) \rightarrow (\mathcal{M}, \in, B)$$

for some transitive  $\mathcal{M}$  such that, letting  $\nu := \min\{\lambda, \eta\}$ , it holds that  $H_\nu^V \subseteq \mathcal{M}$ ,  $A \cap H_\nu^V = B \cap H_\nu^V$  and  $\text{crit } \pi < \nu$ .

PROOF. Assume that we have a non-trivial homomorphism  $h: \mathcal{P}_{\lambda,A} \rightarrow \mathcal{P}_{\eta,A}$  in a forcing extension  $V[g]$ , define in  $V[g]$  the set

$$\mathcal{M}^* := \{\langle b, f \rangle \mid b \in [H_\nu]^{<\omega} \wedge f \in H_{\lambda+}^V \wedge f: H_\lambda^V \rightarrow H_\lambda^V\},$$

and define the relation  $\in^*$  on  $\mathcal{M}^*$  as

$$\langle b_0, f_0 \rangle \in^* \langle b_1, f_1 \rangle \quad \text{iff} \quad b_0 b_1 \in h(\{xy \in [H_\lambda^V]^{<\omega} \mid f_0(x) \in f_1(y)\}).$$

*Claim 1.26.*  $\in^*$  is wellfounded.

PROOF OF CLAIM. Assume not and let  $\cdots \in^* \langle b_1, f_1 \rangle \in^* \langle b_0, f_0 \rangle$  be an  $\in^*$ -decreasing chain, which by definition means that, for every  $n < \omega$ ,

$$b_{n+1}b_n \in h(\{xy \in [H_\lambda^V]^{<\omega} \mid f_{n+1}(x) \in f_n(y)\}). \quad (1)$$

Define the relation  $R(v_0, v_1, v_2)$  on  $H_\lambda^V$  as

$$R(X, f, g) \text{ iff } X = \{xy \in [H_\lambda^V]^{<\omega} \mid f(x) \in g(y)\}.$$

This relation is equal to  $R_\lambda^\varphi$  for some  $\varphi$ , so  $h$  moves  $\langle X, f, g \rangle \in R_\lambda^\varphi$  to

$$\langle h(X), h(f), h(g) \rangle \in R_\eta^\varphi,$$

meaning that

$$h(\{xy \in [H_\lambda^V]^{<\omega} \mid f_{n+1}(x) \in f_n(y)\}) = \{xy \in [H_\eta^V]^{<\omega} \mid f_{n+1}^*(x) \in f_n^*(y)\}$$

for some  $f_n^*$  such that  $f_n^* \cap H_\eta^V = h(f_n)$  for all  $n < \omega$ . But now (1) implies that

$$b_{n+1}b_n \in \{xy \in [H_\eta^V]^{<\omega} \mid f_{n+1}^*(x) \in f_n^*(y)\}$$

and so  $h(f_{n+1})(x) = f_{n+1}^*(x) \in f_n^*(y) = h(f_n)(y)$ , giving an  $\in$ -decreasing sequence in  $V[g]$  using transitivity of  $H_\eta^V$ , a contradiction!

Hence  $\in^*$  is wellfounded.  $\dashv$

$\mathcal{M}^*$  is a set, so  $\in^*$  is trivially set-like. This means that we can take the transitive collapse  $(\mathcal{M}, \in) \cong (\mathcal{M}^*, \in^*)$ , and we note that  $\mathcal{M} = \{[b, f] \mid \langle b, f \rangle \in \mathcal{M}^*\}$ , where  $[b, f] := \{[\bar{b}, \bar{f}] \mid \langle \bar{b}, \bar{f} \rangle \in^* \langle b, f \rangle\}$ .

We now get a version of Łoś' Theorem whose proof is straight-forward, using that  $h$  preserves all  $\Sigma_1$ -relations and that  $H_\lambda^V \models \text{ZFC}^-$ .

*Claim 1.27.* For every formula  $\varphi(v_1, \dots, v_n)$  and every  $[b_1, f_1], \dots, [b_n, f_n] \in \mathcal{M}$  the following are equivalent:

- (i)  $(\mathcal{M}, \in) \models \varphi[[b_1, f_1], \dots, [b_n, f_n]]$ ;
- (ii)  $b_1 \cdots b_n \in h(\{a_1 \cdots a_n \mid \mathcal{P}_{\lambda, A} \models \varphi[f_1(a_1), \dots, f_n(a_n)]\})$ .

PROOF OF CLAIM. The proof is straightforward, using that  $h$  preserves  $\Sigma_1$ -relations. We prove this by induction on  $\varphi$ . If  $\varphi$  is  $v_i \in v_j$  then we have that

$$\begin{aligned}
 & (\mathcal{M}, \in) \models \varphi[[b_1, f_1], \dots, [b_n, f_n]] \\
 & \Leftrightarrow [b_i, f_i] \in [b_j, f_j] \\
 & \Leftrightarrow \langle b_i, f_i \rangle \in^* \langle b_j, f_j \rangle \\
 & \Leftrightarrow b_i b_j \in h(\{a_i a_j \in [H_\lambda^V]^{<\omega} \mid f_i(a_i) \in f_j(a_j)\}) \\
 & \Leftrightarrow b_1 \cdots b_n \in h(\{a_1 \cdots a_n \mid f_i(a_i) \in f_j(a_j)\}) \\
 & \Leftrightarrow b_1 \cdots b_n \in h(\{a_1 \cdots a_n \mid \mathcal{P}_{\lambda, A} \models \varphi[f_1(a_1), \dots, f_n(a_n)]\}).
 \end{aligned}$$

If  $\varphi$  is  $\psi \wedge \chi$  then

$$\begin{aligned}
 & (\mathcal{M}, \in) \models \varphi[[b_1, f_1], \dots, [b_n, f_n]] \\
 & \Leftrightarrow (\mathcal{M}, \in) \models \psi[[b_1, f_1], \dots, [b_n, f_n]] \wedge \chi[[b_1, f_1], \dots, [b_n, f_n]] \\
 & \Leftrightarrow b_1 \cdots b_n \in h(\{a_1 \cdots a_n \mid \mathcal{P}_{\lambda, A} \models \psi[f_1(a_1), \dots, f_n(a_n)]\}) \cap \\
 & \quad h(\{a_1 \cdots a_n \mid \mathcal{P}_{\lambda, A} \models \chi[f_1(a_1), \dots, f_n(a_n)]\}) \\
 & \Leftrightarrow b_1 \cdots b_n \in h(\{a_1 \cdots a_n \mid \mathcal{P}_{\lambda, A} \models \varphi[f_1(a_1), \dots, f_n(a_n)]\}).
 \end{aligned}$$

If  $\varphi$  is  $\neg\psi$  then

$$\begin{aligned}
 & (\mathcal{M}, \in) \models \varphi[[b_1, f_1], \dots, [b_n, f_n]] \\
 & \Leftrightarrow (\mathcal{M}, \in) \models \neg\psi[[b_1, f_1], \dots, [b_n, f_n]] \\
 & \Leftrightarrow (\mathcal{M}, \in) \not\models \psi[[b_1, f_1], \dots, [b_n, f_n]] \\
 & \Leftrightarrow b_1 \cdots b_n \notin h(\{a_1 \cdots a_n \mid \mathcal{P}_{\lambda, A} \models \psi[f_1(a_1), \dots, f_n(a_n)]\}) \\
 & \Leftrightarrow b_1 \cdots b_n \in h(\{a_1 \cdots a_n \mid \mathcal{P}_{\lambda, A} \models \varphi[f_1(a_1), \dots, f_n(a_n)]\}).
 \end{aligned}$$

Finally, if  $\varphi$  is  $\exists x\psi$  then

$$\begin{aligned} & (\mathcal{M}, \in) \models \varphi[[b_1, f_1], \dots, [b_n, f_n]] \\ \Leftrightarrow & (\mathcal{M}, \in) \models \exists x\psi[x, [b_1, f_1], \dots, [b_n, f_n]] \\ \Leftrightarrow & \exists \langle b, f \rangle \in \mathcal{M}^*: (\mathcal{M}, \in) \models \psi[[b, f], [b_1, f_1], \dots, [b_n, f_n]] \\ \Leftrightarrow & \exists \langle b, f \rangle \in \mathcal{M}^*: bb_1 \cdots b_n \in h(\{aa_1 \cdots a_n \mid \mathcal{P}_{\lambda, A} \models \psi[f(a), f_1(a_1), \dots, f_n(a_n)]\}) \\ \Leftrightarrow & b_1 \cdots b_n \in h(\{a_1 \cdots a_n \mid \mathcal{P}_{\lambda, A} \models \varphi[f_1(a_1), \dots, f_n(a_n)]\}). \end{aligned}$$

This finishes the proof.  $\dashv$

Next up, we have the following standard lemma, which implies that  $H_\eta^V \subseteq \mathcal{M}$ :

*Claim 1.28.* For all  $y \in H_\eta^V$  we have  $y = [\langle y \rangle, \text{pr}]$ , where  $\text{pr}(\langle x \rangle) := x$ .

**PROOF OF CLAIM.** We prove this by  $\in$ -induction on  $y \in H_\eta^V$ , so suppose that  $y' = [\langle y' \rangle, \text{pr}]$  for every  $y' \in y$ , which implies that  $y \subseteq \mathcal{M}$  by transitivity of  $\mathcal{M}$ . We then get that, for every  $[b, f] \in \mathcal{M}$ ,

$$\begin{aligned} [b, f] \in [\langle y \rangle, \text{pr}] & \Leftrightarrow b\langle y \rangle \in h(\{a\langle x \rangle \mid f(a) \in \text{pr}(\langle x \rangle)\}) \\ & \Leftrightarrow \exists y' \in y: b\langle y' \rangle \in h(\{a\langle x \rangle \mid f(a) = x\}) \\ & \Leftrightarrow \exists y' \in y: [b, f] = [\langle y' \rangle, \text{pr}] = y' \\ & \Leftrightarrow [b, f] \in y, \end{aligned}$$

showing that  $y = [\langle y \rangle, \text{pr}]$ .  $\dashv$

Now define

$$B := \{[b, f] \in \mathcal{M} \mid b \in h(\{x \in H_\lambda^V \mid f(x) \in A\})\}.$$

and, in  $V[g]$ , let  $\pi: (H_\lambda^V, \in, A \cap H_\lambda^V) \rightarrow (\mathcal{M}, \in, B)$  be given as  $\pi(x) := [\langle x \rangle, c_x]$ .

*Claim 1.29.*  $\pi$  is elementary.

PROOF OF CLAIM. For  $x_1, \dots, x_n \in H_\lambda^V$  it holds that

$$\begin{aligned} & (\mathcal{M}, \in, B) \models \varphi[\pi(x_1), \dots, \pi(x_n)] \\ \Leftrightarrow & (\mathcal{M}, \in) \models \varphi[\pi(x_1), \dots, \pi(x_n)] \\ \Leftrightarrow & \langle \rangle \in h(\{\langle \rangle \mid \mathcal{P}_{\lambda, A} \models \varphi[x_1, \dots, x_n]\}) \\ \Leftrightarrow & (H_{\lambda^+}^V, \in, A \cap H_\lambda^V) \models \varphi[x_1, \dots, x_n] \end{aligned}$$

and we also get that, for every  $x \in H_\lambda^V$ ,

$$x \in A \Leftrightarrow \langle \rangle \in h(\{a \in H_\lambda^V \mid x \in A\}) \Leftrightarrow \pi(x) \in B,$$

which shows elementarity.  $\dashv$

We next need to show that  $B \cap H_\nu^V = A \cap H_\nu^V$ , so let  $x \in H_\nu^V$ . Note that  $x = [\langle x \rangle, \text{pr}]$  by Claim 1.28, which means that

$$x \in B \Leftrightarrow \langle x \rangle \in h(\{\langle y \rangle \in H_\lambda^V \mid y \in A\}) \Leftrightarrow x \in A.$$

The last thing we need to show is that  $\text{crit } \pi < \nu$ . We start with an analogous result about  $h$ .

*Claim 1.30.* There exists some  $b \in H_\nu^V$  such that  $h(b) \neq b$ .

PROOF OF CLAIM. Assume the claim fails. We now have two cases.

**Case 1:**  $\lambda \geq \eta$

By non-triviality of  $h$  there's an  $x \in H_{\lambda^+}^V$  such that  $h(x) \cap H_\eta^V \neq x \cap H_\eta^V$ , which means that there exists an  $a \in H_\eta^V$  such that  $a \in h(x) \Leftrightarrow a \notin x$ .

If  $a \in x$  then  $\{a\} = h(\{a\}) \subseteq h(x)$ ,<sup>13</sup> making  $a \in h(x)$ ,  $\nmid$ , so assume instead that  $a \in h(x)$ . Since  $\eta$  is a strong limit cardinal we may fix a

<sup>13</sup>Note that as  $h$  preserves  $\Sigma_1$  formulas it also preserves singletons and boolean operations.

cardinal  $\theta < \eta$  such that  $a \in H_\theta^V$  and  $H_\theta^V \in H_\eta^V$ . We then have that<sup>14</sup>

$$\{a\} \subseteq h(x) \cap H_\theta^V = h(x) \cap h(H_\theta^V) = h(x \cap H_\theta^V) = x \cap H_\theta^V,$$

so that  $a \in x$ ,  $\nless$ .

**Case 2:**  $\lambda < \eta$

In this case we are assuming that  $h \restriction H_\lambda^V = \text{id}$ , but  $h(\lambda) = \eta > \lambda$ . Since  $\lambda$  is singular we can fix some  $\gamma < \lambda$  and a cofinal function  $f: \gamma \rightarrow \lambda$ . Define the relation

$$R = \{(\alpha, \beta, \bar{\alpha}, \bar{\beta}, g) \mid \ulcorner g \text{ is a cofinal function } g: \alpha \rightarrow \beta^\top \wedge g(\bar{\alpha}) = \bar{\beta}\}.$$

Then  $R(\gamma, \lambda, \alpha, f(\alpha), f)$  holds by assumption for every  $\alpha < \gamma$ , so that  $R$  holds for some  $(\gamma^*, \lambda^*, \alpha^*, f(\alpha)^*, f^*)$  such that

$$\begin{aligned} (\gamma^*, \lambda^*, \alpha^*, f(\alpha)^*, f^*) \cap H_\eta^V &= (h(\gamma), h(\lambda), h(\alpha), h(f(\alpha)), h(f)) \\ &= (\gamma, \eta, \alpha, f(\alpha), h(f)), \end{aligned}$$

using our assumption that  $h$  fixes every  $b \in H_\lambda^V$ . Since  $\gamma$ ,  $\alpha$  and  $f(\alpha)$  are transitive and bounded in  $H_\lambda^V$  it holds that  $h(\gamma) = \gamma^*$ ,  $h(\alpha) = \alpha^*$  and  $h(f(\alpha)) = f(\alpha)^*$ . Also, since  $\text{dom}(f^*) = \gamma = \text{dom}(f)$  we must in fact have that  $f^* = h(f)$ . But this means that  $h(f): \gamma \rightarrow \eta$  is cofinal and  $\text{ran}(h(f)) \subseteq \lambda$ , a contradiction!  $\dashv$

To use the above Claim 1.30 to conclude anything about  $\pi$  we'll make use of the following standard lemma.

*Claim 1.31.* For any  $x \in H_\lambda^V$  it holds that  $h(x) \cap H_\eta^V = \pi(x) \cap H_\eta^V$ .

<sup>14</sup>Note that we're using  $\lambda \geq \eta$  here to ensure that  $H_\theta^V \in \text{dom } h$ .

PROOF OF CLAIM. For any  $n < \omega$  and  $\langle a_1, \dots, a_n \rangle \in [H_\eta^V]^n$  we have that

$$\begin{aligned} & \langle a_1, \dots, a_n \rangle \in \pi(x) \\ \Leftrightarrow & (\mathcal{M}, \in) \models \langle a_1, \dots, a_n \rangle \in \pi(x) \\ \Leftrightarrow & (\mathcal{M}, \in) \models \langle [\langle a_1 \rangle, \text{pr}], \dots, [\langle a_n \rangle, \text{pr}] \rangle \in [\langle \rangle, c_x] \\ \Leftrightarrow & \langle a_1, \dots, a_n \rangle \in h(\{ \langle x_1, \dots, x_n \rangle \mid \mathcal{P}_{\lambda, A} \models \langle x_1, \dots, x_n \rangle \in x \}) \\ \Leftrightarrow & \langle a_1, \dots, a_n \rangle \in h(x), \end{aligned}$$

showing that  $h(x) \cap H_\eta^V = \pi(x) \cap H_\eta^V$ . ⊢

Now use Claim 1.30 to fix a  $b \in H_\nu^V$  which is moved by  $h$ . Claim 1.31 then implies that

$$\pi(b) \cap H_\eta^V = h(b) \cap H_\eta^V = h(b) \neq b = b \cap H_\eta^V,$$

showing that  $\pi(b) \neq b$  and hence  $\text{crit } \pi < \nu$ . This finishes the proof of the lemma. ■

**Theorem 1.32** (GBC, Gitman-N.). *gVP is equivalent to gWVP.*

PROOF. ( $\Rightarrow$ ): Assume gVP holds and gWVP fails, and let  $\langle \mathcal{M}_\alpha \mid \alpha < \text{On} \rangle$  be an On-sequence of first-order structures such that for every  $\alpha \leq \beta$  there exists a generic homomorphism

$$j_{\beta\alpha}: \mathcal{M}_\beta \rightarrow \mathcal{M}_\alpha$$

in some  $V[g]$  which is unique in all generic extensions, with no generic homomorphisms going the other way. Here we may assume, as in the proof of Lemma 1.21, that  $g \subseteq \text{Col}(\omega, \mathcal{M}_\beta)$ . We can then find a proper class  $C \subseteq \text{On}$  such that  $|\mathcal{M}_\alpha|^V < |\mathcal{M}_\beta|^V$  for every  $\alpha < \beta$  in  $C$ . By gVP there are then

$\alpha < \beta$  in  $C$  and a generic homomorphism

$$\pi: \mathcal{M}_\alpha \rightarrow \mathcal{M}_\beta.$$

in some  $V[h]$ , where again we may assume that  $h \subseteq \text{Col}(\omega, \mathcal{M}_\alpha)$ . But then  $\pi \circ j_{\beta\alpha} = \text{id}$  by uniqueness of  $j_{\beta\beta} = \text{id}$ , which means that  $j_{\beta\alpha}$  is injective in  $V[g \times h]$  and hence also in  $V[g]$ . But then  $|\mathcal{M}_\beta|^{V[g]} \leq |\mathcal{M}_\alpha|^{V[g]}$ , which implies that  $|\mathcal{M}_\beta|^V \leq |\mathcal{M}_\alpha|^V$  by the  $|\mathcal{M}_\beta|^{+V}$ -cc of  $\text{Col}(\omega, \mathcal{M}_\beta)$ , contradicting the definition of  $C$ .

( $\Leftarrow$ ): Assume that  $\text{gVP}$  fails, which by Theorem 1.20 is equivalent to  $\text{On}$  not being faintly prewoodin. This means that there exists a class  $A$  such that there are no faintly  $A$ -prestrong cardinals. We can therefore assign to any cardinal  $\kappa$  the least cardinal  $f(\kappa) > \kappa$  such that  $\kappa$  is not faintly  $(f(\kappa), A)$ -prestrong.

Also define a function  $g: \text{On} \rightarrow \text{Card}$  as taking an ordinal  $\alpha$  to the least singular strong limit cardinal above  $\alpha$  closed under  $f$ . Then we're assuming that there's no non-trivial generic elementary embedding

$$\pi: (H_{g(\alpha)}^V, \in, A \cap H_{g(\alpha)}^V) \rightarrow (\mathcal{M}, \in, B)$$

with  $H_{g(\alpha)}^V \subseteq \mathcal{M}$  and  $B \cap H_{g(\alpha)}^V = A \cap H_{g(\alpha)}^V$ . Assume towards a contradiction that for some  $\alpha, \beta$  there is a non-trivial generic homomorphism  $h: \mathcal{P}_{g(\alpha), A} \rightarrow \mathcal{P}_{g(\beta), A}$ . Lemma 1.25 then gives us a non-trivial generic elementary embedding

$$\pi: (H_{g(\alpha)}^V, \in, A \cap H_{g(\alpha)}^V) \rightarrow (\mathcal{M}, \in, B)$$

for some transitive  $\mathcal{M}$  such that  $H_\nu^V \subseteq \mathcal{M}$  with  $\nu := \min\{g(\alpha), g(\beta)\}$  and  $A \cap H_\nu^V = B \cap H_\nu^V$ , a contradiction! Therefore every generic homomorphism  $h: \mathcal{P}_{g(\alpha), A} \rightarrow \mathcal{P}_{g(\beta), A}$  is trivial. Since there is a unique trivial homomorphism when  $\alpha \geq \beta$  and no trivial homomorphism when  $\alpha < \beta$  since  $g(\alpha)$  is sent to  $g(\beta)$ , the sequence of structures

$$\langle \mathcal{P}_{g(\alpha), A} \mid \alpha \in \text{On} \rangle$$



is a counterexample to  $\mathbf{gWVP}$ , which is what we wanted to show.  $\blacksquare$

### 1.3 BERKELEYS

Berkeley cardinals was introduced by Woodin at University of California, Berkeley around 1992, and was introduced as a large cardinal candidate that would be inconsistent with  $\mathbf{ZF}$ . They trivially imply the Kunen inconsistency and are therefore at least inconsistent with  $\mathbf{ZFC}$ , but that's as far as it currently goes. In the virtual setting the virtually berkeley cardinals, like all the other virtual large cardinals, are simply downwards absolute to  $L$ . See Appendix ?? for a quick overview of what is currently known about the non-virtual Berkeley cardinals.

It turns out that virtually berkeley cardinals are natural objects, as the main theorem of this section shows that these large cardinals are precisely what separates virtually prewoodins from the virtually woodins, as well as separating virtually vopěnka cardinals from mahlo cardinals.

**Definition 1.33.** Say that a cardinal  $\delta$  is **virtually proto-berkeley** if for every transitive set  $\mathcal{M}$  such that  $\delta \subseteq \mathcal{M}$  there exists a generic elementary embedding  $\pi: \mathcal{M} \rightarrow \mathcal{M}$  with  $\text{crit } \pi < \delta$ .

If  $\text{crit } \pi$  can be chosen arbitrarily large below  $\delta$  then  $\delta$  is **virtually berkeley**, and if  $\text{crit } \pi$  can be chosen as an element of any club  $C \subseteq \delta$  we say  $\delta$  is **virtually club berkeley**.  $\circ$

Virtually (proto-)berkeley cardinals turn out to be equivalent to their “bold-face” versions, the proof of which is a straight-forward virtualisation of Lemma 2.1.12 and Corollary 2.1.13 in [?].

**Proposition 1.34** (Virtualised Cutolo). *If  $\delta$  is virtually proto-berkeley then for every transitive set  $\mathcal{M}$  such that  $\delta \subseteq \mathcal{M}$  and every subset  $A \subseteq \mathcal{M}$  there exists a generic elementary embedding  $\pi: (\mathcal{M}, \in, A) \rightarrow (\mathcal{M}, \in, A)$  with  $\text{crit } \pi < \delta$ . If  $\delta$  is virtually berkeley then we can furthermore ensure that  $\text{crit } \pi$  is arbitrarily large below  $\delta$ .*

PROOF. Let  $\mathcal{M}$  be transitive with  $\delta \subseteq \mathcal{M}$  and  $A \subseteq \mathcal{M}$ . Let

$$\mathcal{N} := \mathcal{M} \cup \{\{\langle A, x \rangle \mid x \in \mathcal{M}\}\}$$

and note that  $\mathcal{N}$  is transitive. Further, both  $A$  and  $\mathcal{M}$  are definable in  $\mathcal{N}$  without parameters:  $a$  is the first element in the pairs belonging to the set of highest rank, and  $\mathcal{M}$  is what remains if we remove the set with the highest rank. But this means that a generic elementary embedding  $\pi: \mathcal{N} \rightarrow \mathcal{N}$  fixes both  $\mathcal{M}$  and  $a$ , giving us a generic elementary  $\sigma: (\mathcal{M}, \in, A) \rightarrow (\mathcal{M}, \in, A)$  with  $\text{crit } \sigma = \text{crit } \pi$ , yielding the wanted conclusion. ■

The following is a straight-forward virtualisation of the usual definition of the vopěnka filter (see e.g. [?]).

**Definition 1.35** (GBC). Define the **virtually vopěnka filter**  $F$  on  $\text{On}$  as  $X \in F$  iff there's a natural  $\text{On}$ -sequence  $\vec{\mathcal{M}}$  such that  $\text{crit } \pi \in X$  for any  $\alpha < \beta$  and any generic elementary  $\pi: \mathcal{M}_\alpha \rightarrow \mathcal{M}_\beta$ . ◦

Theorem 1.20 shows that  $\emptyset$  is in the virtually vopěnka filter iff  $\text{gVP}$  fails, in analogy with the non-virtual case. Normality also holds in the virtual context, as the following proof shows.

**Lemma 1.36** (GBC, Virtualised folklore). *The virtually vopěnka filter is a normal filter.*

PROOF. Let  $F$  be the virtually vopěnka filter. We first show that  $F$  is actually a filter. If  $X \in F$  and  $Y \supseteq X$  then  $Y \in F$  simply by definition of  $F$ . If  $X, Y \in F$ , witnessed by natural sequences  $\vec{\mathcal{M}}$  and  $\vec{\mathcal{N}}$ , then  $X \cap Y \in F$  as well, witnessed by the natural sequence  $\vec{\mathcal{P}}$  induced by the indexing function  $f^{\vec{\mathcal{P}}} := \max(f^{\vec{\mathcal{M}}}, f^{\vec{\mathcal{N}}})$  and unary relations  $R_\alpha^{\vec{\mathcal{P}}} := \text{Code}(\langle R_\alpha^{\vec{\mathcal{M}}}, R_\alpha^{\vec{\mathcal{N}}} \rangle)$ . Indeed, if  $\pi: \mathcal{P}_\alpha \rightarrow \mathcal{P}_\beta$  is a generic elementary embedding with critical point  $\mu$  then  $\mu$  is also the critical point of both  $\pi \restriction \mathcal{M}_\alpha: \mathcal{M}_\alpha \rightarrow \mathcal{M}_\beta$  and  $\pi \restriction \mathcal{N}_\alpha: \mathcal{N}_\alpha \rightarrow \mathcal{N}_\beta$ .

For normality, let  $X \in F^+$  be  $F$ -positive, where we recall that this means that  $X \cap C \neq \emptyset$  for every  $C \in F$ , and let  $f: X \rightarrow \mathbf{On}$  be regressive. We want to show that  $f$  is constant on an  $F$ -positive set.

Assume this fails, meaning that there are natural sequences  $\vec{\mathcal{M}}^\gamma$  for  $\gamma$  such that for any generic elementary  $\pi: \mathcal{M}_\alpha^\gamma \rightarrow \mathcal{M}_\beta^\gamma$  satisfies that  $f(\text{crit } \pi) \neq \gamma$ . Define a new natural sequence  $\vec{\mathcal{N}}$  as induced by the indexing function  $g: \mathbf{On} \rightarrow \mathbf{On}$  given as  $g(\alpha) := \sup_{\gamma < \alpha} \text{rk } \mathcal{M}_\alpha^\gamma + \omega$  and unary relations  $R_\alpha^{\vec{\mathcal{N}}}$  given as

$$R_\alpha^{\vec{\mathcal{N}}} := \text{Code}(\langle \langle \mathcal{M}_\alpha^\gamma \mid \gamma < \alpha \rangle, f \restriction \alpha \rangle).$$

Now since  $X$  is  $F$ -positive there exists a generic elementary embedding  $\pi: \mathcal{N}_\alpha \rightarrow \mathcal{N}_\beta$  with  $\text{crit } \pi \in X$ . As  $f(\text{crit } \pi) < \text{crit } \pi$  we get that  $\pi(f(\text{crit } \pi)) = f(\text{crit } \pi)$ , so that we have a generic elementary embedding

$$\pi \restriction \mathcal{M}_\alpha^{f(\text{crit } \pi)}: \mathcal{M}_\alpha^{f(\text{crit } \pi)} \rightarrow \mathcal{M}_\beta^{f(\text{crit } \pi)},$$

but this contradicts the definition of  $\vec{\mathcal{M}}^{f(\text{crit } \pi)}$ ! Thus  $F$  is normal.  $\blacksquare$

The reason why we are being careful in showing all these analogous properties for the virtual vopěnka filter is that not all properties carry over. Indeed, note that uniformity of filters is non-trivial as we're working with proper classes<sup>15</sup>, and we will see in Theorem 1.40 shows that uniformity of this filter is equivalent to there being no virtually berkeley cardinals — the following lemma is the first implication.

**Lemma 1.37** (GBC, N.). *Assume  $gVP$  and that there are no virtually berkeley cardinals. Then the virtually vopěnka filter  $F$  on  $\mathbf{On}$  contains every class club  $C$ .*

**PROOF.** The crucial extra property we get by assuming that there aren't any virtually berkeleys is that  $F$  becomes uniform, i.e. contains every tail  $(\delta, \mathbf{On}) \subseteq \mathbf{On}$ . Indeed, assume that  $\delta$  is the least cardinal such that  $(\delta, \mathbf{On}) \notin F$ .

<sup>15</sup>This boils down to the fact that the class club filter is not provably normal in GBC, see [?]

$F$ . Let  $M$  be a transitive set with  $\delta \subseteq M$  and  $\gamma < \delta$  a cardinal. As  $(\gamma, \text{On}) \in F$  by minimality of  $\delta$ , we may fix a natural sequence  $\vec{\mathcal{N}}$  witnessing this. Let  $\vec{\mathcal{M}}$  be the natural sequence induced by the indexing function  $f: \text{On} \rightarrow \text{On}$  given by

$$f(\alpha) := \max(\alpha + 1, \delta + 1)$$

and unary relations  $R_\alpha := \langle M, \mathcal{N}_\alpha \rangle$ . If  $\pi: \mathcal{M}_\alpha \rightarrow \mathcal{M}_\beta$  is a generic elementary embedding with  $\text{crit } \pi \leq \delta$ , which exists as  $(\delta, \text{On}) \notin F$ , then  $\pi(R_\alpha) = R_\beta$  implies that  $\pi \restriction \mathcal{M}: \mathcal{M} \rightarrow \mathcal{M}$  with  $\text{crit } \pi \leq \delta$ . We also get that  $\text{crit } \pi > \gamma$ , as

$$\pi \restriction \mathcal{N}_{\text{crit } \pi}: \mathcal{N}_{\text{crit } \pi} \rightarrow \mathcal{N}_{\pi(\text{crit } \pi)}$$

is an embedding between two structures in  $\vec{\mathcal{N}}$  and hence  $\text{crit } \pi > \gamma$  as  $(\gamma, \text{On}) \in F$ . This means that  $\delta$  is virtually berkeley, a contradiction. Thus  $\text{crit } \pi > \delta$ , implying that  $(\delta, \text{On}) \in F$ .

Note that the class  $C_0 \subseteq \text{On}$  of limit ordinals is in  $F$ , since it's the diagonal intersection of the tails  $(\alpha + 1, \text{On})$ . Now let  $C \subseteq \text{On}$  be a class club, and let  $C = \{a_\alpha \mid \alpha < \text{On}\}$  be its increasing enumeration. Then  $C \supseteq C_0 \cap \Delta_{\alpha < \text{On}}(a_\alpha, \text{On})$ , implying that  $C \in F$ . ■

**Theorem 1.38** (GBC, N.). *If there are no virtually berkeley cardinals then  $\text{On}$  is virtually prewoodin iff  $\text{On}$  is virtually woodin.*

PROOF. Assume  $\text{On}$  is virtually prewoodin, so gVP holds by Theorem 1.20 and we can let  $F$  be the virtually vopěnka filter. The assumption that there aren't any virtually berkeley cardinals implies that for any class  $A$  we not only get a virtually  $A$ -prestrong cardinal, but we get stationarily many such. Indeed, assume this fails — we will follow the proof of Theorem 1.20.

Failure means that there is some class  $A$  and some class club  $C$  such that there are no virtually  $A$ -prestrong cardinals in  $C$ . Since there are no virtually berkeley cardinals, Lemma 1.37 implies that  $C \in F$ , so there exists some natural sequence  $\vec{\mathcal{N}}$  such that whenever  $\pi: \mathcal{N}_\alpha \rightarrow \mathcal{N}_\beta$  is an elementary

embedding between two distinct structures of  $\vec{\mathcal{N}}$  it holds that  $\text{crit } \pi \in C$ . Define  $f: \text{On} \rightarrow \text{On}$  as sending  $\alpha$  to the least cardinal  $\eta > \alpha$  such that  $\alpha$  is not virtually  $(\eta, A)$ -prestrong if  $\alpha \in C$ , and set  $f(\alpha) := \alpha$  if  $\alpha \notin C$ . Also define  $g: \text{On} \rightarrow \text{On}$  as  $g(\alpha)$  being the least strong limit cardinal in  $C$  above  $\alpha$  which is a closure point for  $f$ .

Now let  $\vec{\mathcal{M}}$  be the natural sequence induced by  $g$  and  $R_\alpha := \text{Code}(\langle A \cap H_{g(\alpha)}^V, \mathcal{N}_\alpha \rangle)$  and apply **gVP** to get  $\alpha < \beta$  and a generic elementary embedding  $\pi: \mathcal{M}_\alpha \rightarrow \mathcal{M}_\beta$ , which restricts to

$$\pi \upharpoonright (H_{g(\alpha)}^V, \in, A \cap H_{g(\alpha)}^V): (H_{g(\alpha)}^V, \in, A \cap H_{g(\alpha)}^V) \rightarrow (H_{g(\beta)}^V, \in, A \cap H_{g(\beta)}^V),$$

making  $\text{crit } \pi$  virtually  $(g(\alpha), A)$ -prestrong and thus  $\text{crit } \pi \notin C$ . But as we also get the embedding  $\pi \upharpoonright \mathcal{N}_\alpha: \mathcal{N}_\alpha \rightarrow \mathcal{N}_\beta$ , we have that  $\text{crit } \pi \in C$  by definition of  $\vec{\mathcal{N}}$ ,  $\nless$ .

Now fix any class  $A$  and some large  $n < \omega$  and define the class

$$C := \{\kappa \in \text{Card} \mid (H_\kappa^V, \in, A \cap H_\kappa^V) \prec_{\Sigma_n} (V, \in, A)\}.$$

This is a club and we can therefore find a virtually  $A$ -prestrong cardinal  $\kappa \in C$ . Assume that  $\kappa$  is not virtually  $A$ -strong and let  $\theta$  be least such that it isn't virtually  $(\theta, A)$ -strong. Fix a generic elementary embedding

$$\pi: (H_\theta^V, \in, A \cap H_\theta^V) \rightarrow (M, \in, B)$$

with  $\text{crit } \pi = \kappa$ ,  $H_\theta^V \subseteq M$ ,  $M \subseteq V$ ,  $A \cap H_\theta^V = B \cap H_\theta^V$  and  $\pi(\kappa) < \theta$ .

Now  $\pi(\kappa)$  is inaccessible, and  $(H_{\pi(\kappa)}^V, \in, A \cap H_{\pi(\kappa)}^V) = (H_{\pi(\kappa)}^M, \in, B \cap H_{\pi(\kappa)}^M)$  believes that  $\kappa$  is virtually  $(A \cap H_{\pi(\kappa)}^V)$ -strong as in the proof of Theorem 1.9, meaning that  $(H_\kappa^V, \in, A \cap H_\kappa^V)$  believes that there is a proper class of virtually  $(A \cap H_\kappa^V)$ -strong cardinals. But  $\kappa \in C$ , which means that

$$(V, \in, A) \models \ulcorner \text{There exists a proper class of virtually } A\text{-strong cardinals} \urcorner,$$

implying that  $\text{On}$  is virtually woodin. ■

**Theorem 1.39** (GBC, N.). *If there exists a virtually berkeley cardinal  $\delta$  then  $\mathbf{gVP}$  holds and  $\mathbf{On}$  is not mahlo.*

PROOF. If  $\mathbf{On}$  was Mahlo then there would in particular exist an inaccessible cardinal  $\kappa > \delta$ , but then  $H_\kappa^V \models \ulcorner \text{there exists a virtually berkeley cardinal} \urcorner$ , contradicting the incompleteness theorem.

To show  $\mathbf{gVP}$  we show that  $\mathbf{On}$  is virtually prewoodin, which is equivalent by Theorem 1.20. Fix therefore a class  $A$  — we have to show that there exists a virtually  $A$ -prestrong cardinal. For every cardinal  $\theta \geq \delta$  there exists a generic elementary embedding

$$\pi_\theta: (H_\theta^V, \in, A \cap H_\theta^V) \rightarrow (H_\theta^V, \in, A \cap H_\theta^V)$$

with  $\text{crit } \pi < \delta$ . By the pigeonhole principle we thus get some  $\kappa < \delta$  which is the critical point of proper class many  $\pi_\theta$ , showing that  $\kappa$  is virtually  $A$ -prestrong, making  $\mathbf{On}$  virtually prewoodin. ■

**Theorem 1.40** (GBC, N.). *The following are equivalent:*

- (i)  $\mathbf{gVP}$  implies that  $\mathbf{On}$  is mahlo;
- (ii)  $\mathbf{On}$  is virtually prewoodin iff  $\mathbf{On}$  is virtually woodin;
- (iii) There are no virtually berkeley cardinals.

PROOF. (iii)  $\Rightarrow$  (ii) is Theorem 1.38, and the contraposited version of (i)  $\Rightarrow$  (iii) is Theorem 1.39. For (ii)  $\Rightarrow$  (i) note that  $\mathbf{gVP}$  implies that  $\mathbf{On}$  is virtually prewoodin by Theorem 1.20, which by (ii) means that it's virtually woodin and the usual proof shows that virtually woodins are mahlo<sup>16</sup>, showing (i). ■

This also immediately implies the following equiconsistency, as virtually berkeley cardinals have strictly larger consistency strength than virtually woodin cardinals.

---

<sup>16</sup>See e.g. Exercise 26.10 in [?].

**Corollary 1.41** (N.). *The existence of an inaccessible virtually prewoodin cardinal is equiconsistent with the existence of an inaccessible virtually woodin cardinal.* ■

## 1.4 BEHAVIOUR IN CANONICAL INNER MODELS

Most of the cardinals turn out to be downwards absolute to most inner models, including  $L$ :

**Proposition 1.42.** *For any regular uncountable cardinal  $\theta$ , faintly  $\theta$ -measurable cardinals are downwards absolute to any transitive class  $\mathcal{U} \subseteq V$  satisfying  $ZF^- + DC$ .*

PROOF. Let  $\kappa$  be faintly  $\theta$ -measurable, witnessed by a forcing poset  $\mathbb{P}$  and a  $V$ -generic  $g \subseteq \mathbb{P}$  such that, in  $V[g]$ , there's a transitive  $\mathcal{M}$  and an elementary embedding  $\pi: H_\theta^V \rightarrow \mathcal{M}$  with  $\text{crit } \pi = \kappa$ . Fix a transitive class  $\mathcal{U} \subseteq V$  which satisfies  $ZF^- + DC$ . Restricting the embedding to  $\pi \restriction H_\theta^\mathcal{U}: H_\theta^\mathcal{U} \rightarrow \mathcal{N}$  we can now apply the Countable Absoluteness Lemma ?? to  $\pi \restriction H_\theta^\mathcal{U}$  to get that there exists an embedding  $\pi^*: H_\theta^\mathcal{U} \rightarrow \mathcal{N}^*$  in a generic extension of  $\mathcal{U}$ , making  $\kappa$  faintly  $\theta$ -measurable in  $\mathcal{U}$ . ■

**Proposition 1.43** (N.). *Let  $\theta$  be a regular uncountable cardinal.*

- (i)  $L \models \ulcorner \text{faintly } \theta\text{-measurables are equivalent to virtually } \theta\text{-prestrongs} \urcorner$ .
- (ii) Assume that  $L[\mu]$  exists. It then holds that  $L[\mu] \models \ulcorner \text{faintly } \theta\text{-measurables are equivalent to virtually } \theta\text{-measurables} \urcorner$ .
- (iii) Assume there is no inner model with a woodin. It then holds that  $K \models \ulcorner \text{faintly } \theta\text{-measurables are equivalent to virtually } \theta\text{-measurables} \urcorner$ .

PROOF. For (i) simply note that if  $\pi: L_\theta \rightarrow \mathcal{N}$  is a generic elementary embedding with  $\mathcal{N}$  transitive, then by condensation we have that  $\mathcal{N} = L_\gamma$  for some  $\gamma \geq \theta$ , so that  $\pi$  also witnesses the virtual  $\theta$ -prestrongness of  $\text{crit } \pi$ .

(ii): Assume that  $V = L[\mu]$  for notational simplicity and let  $\kappa$  be faintly  $\theta$ -measurable, witnessed by a generic elementary embedding  $\pi: L_\theta[\mu] \rightarrow \mathcal{N}$  existing in some generic extension  $V[g]$ . By condensation we get that  $\mathcal{N} =$

$L_\gamma[\bar{\mu}]$  for some  $\gamma \geq \theta$  and  $\bar{\mu} \in V[g]$ , but we're not guaranteed that  $\bar{\mu} \in V$  here. Let  $\lambda$  be the unique measurable cardinal of  $V = L[\mu]$ .

Note that  $\bar{\mu}$  is a measure on  $\pi(\lambda) \geq \lambda$ . If  $\pi(\lambda) = \lambda$  then  $L[\mu] = L[\bar{\mu}]$  by [?, Theorem 20.10] and we trivially get that  $\mathcal{N} \subseteq V$ . Assume thus that  $\pi(\lambda) > \lambda$ , which implies that  $L[\bar{\mu}]$  is an internal iterate of  $L[\mu]$  by [?, Theorem 20.12]. In particular it then holds that  $L[\bar{\mu}] \subseteq L[\mu]$ , so again we get that  $\mathcal{N} \subseteq V$ .

(iii): Assume that  $V = K = L[\mathcal{E}]$  and fix a faintly  $\theta$ -measurable cardinal  $\kappa$ , witnessed by a generic embedding  $\pi: L_\theta[\mathcal{E}] \rightarrow \mathcal{N} = L_\gamma[\bar{\mathcal{E}}]$  in some generic extension  $V[g]$ . Now coiterate  $L[\mathcal{E}]$  with  $L[\bar{\mathcal{E}}]$ , and denote the last models by  $\mathcal{P}$  and  $\mathcal{Q}$ . Since  $K = K^{V[g]}$  and as  $K$  is universal we get that  $\mathcal{Q} \leq \mathcal{P}$ . Then the  $L[\bar{\mathcal{E}}]$ -to- $\mathcal{Q}$  branch did not drop, giving us an iteration embedding  $i: L[\bar{\mathcal{E}}] \rightarrow \mathcal{Q}$ .

Note that  $\text{crit } i \geq \kappa$  as  $\bar{\mathcal{E}}$  is simply the pointwise image of  $\mathcal{E}$  under  $\pi$ , so nothing below  $\kappa$  is touched and is therefore not used in the comparison either. This means that  $\text{crit}(i \circ \pi) = \kappa$ , so that  $(i \circ \pi): L_\theta[\mathcal{E}] \rightarrow \mathcal{Q}$  witnesses that  $\kappa$  is virtually  $\theta$ -measurable, since  $\mathcal{Q} \leq \mathcal{P}$  implies that  $\mathcal{Q} \subseteq K$ . ■

Note that the proofs of (ii) and (iii) above do not show that  $\kappa$  is virtually  $\theta$ -prestrong, as it might still be the case that  $\bar{\mu} \neq \mu$  or  $\bar{\mathcal{E}} \neq \mathcal{E}$ , so we cannot conclude that  $L_\theta[\mu] \subseteq L_\theta[\bar{\mu}]$  or  $L_\theta[\mathcal{E}] \subseteq L_\theta[\bar{\mathcal{E}}]$ . It might still hold however; see Questions ?? and ??.

## 1.5 SEPARATION RESULTS

Having proving many positive results about the relations between the virtual large cardinals in the previous sections, this section is dedicated to the negatives. More precisely, we will aim to *separate* many of the defined notions (potentially under suitable large cardinal assumptions).

Our first separation result is that the virtuals form a level-by-level hierarchy.



**Theorem 1.44** (N.). *Let  $\alpha < \kappa$  and assume that  $\kappa$  is faintly  $\kappa^{+\alpha+2}$ -measurable. Then*

$$L_\kappa \models \ulcorner \text{There's a proper class of } \lambda \text{ which are virtually } \lambda^{+\alpha+1}\text{-strong} \urcorner.$$

PROOF. Write  $\theta := \kappa^{+\alpha+1}$ . Then by Theorem 1.9 we get that either  $\kappa$  is faintly  $\theta^+$ -strong in  $L$  or otherwise, in particular,  $L_\kappa$  thinks that there's a proper class of remarkables. In the second case we also get that  $L_\kappa$  thinks that there's a proper class of  $\lambda$  such that  $\lambda$  is virtually  $\lambda^{+\alpha+1}$ -strong and we'd be done, so assume the first case. Then  $L_\kappa \prec_2 L_{\theta^+}$ , so define for each  $\xi < \kappa$  the sentence  $\psi_\xi$  as

$$\psi_\xi := \exists \lambda < \xi : \ulcorner \lambda \text{ is virtually } \lambda^{+\alpha+1}\text{-strong} \urcorner.$$

Then  $\psi_\xi$  is  $\Sigma_2(\{\alpha, \xi\})$  since being virtually  $\beta$ -strong is a  $\Delta_2(\{\beta\})$ -statement. As  $L_{\theta^+} \models \psi_\xi$  for all  $\xi < \kappa$  we also get that  $L_\kappa \models \psi_\xi$  for all  $\xi < \kappa$ , which is what we wanted to show. ■

As we are only assuming  $\kappa$  to be *faintly* measurable in the above, this also shows that the faintly  $\kappa^{+\alpha+1}$ -measurable cardinals  $\kappa$  form a strict hierarchy whenever  $\alpha < \kappa$ .

Next, we show that the “virtually” adverb *does* yield cardinals different from the faintly ones. This is trivial in general as successor cardinals can be faintly measurable and are never virtually measurable, but the separation still holds true if we rule out this successor case. For a slightly more fine-grained distinction let's define an intermediate large cardinal between the faintly and virtual.

**Definition 1.45.** Let  $\kappa < \theta$  be infinite regular cardinals. Say that  $\kappa$  is **faintly  $\theta$ -power- $\Phi$**  for  $\Phi \in \{\text{measurable, prestrong, strong}\}$  if it is faintly  $\theta$ - $\Phi$ , witnessed by an embedding  $\pi: H_\theta^V \rightarrow \mathcal{N}$ , and  $\mathcal{P}^V(\kappa) = \mathcal{P}^\mathcal{N}(\kappa)$ . ◦

Note that the proof of Lemma 1.2 shows that faintly power-measurables are also 1-iterable and so in particular weakly compact. Our separation result is then the following.

**Theorem 1.46** (Gitman-N.). *For  $\Phi \in \{\text{measurable, prestrong, strong}\}$ , if  $\kappa$  is virtually  $\Phi$  then there exist forcing extensions  $V[g]$  and  $V[h]$  such that*

- (i) *In  $V[g]$ ,  $\kappa$  is inaccessible and faintly  $\Phi$  but not faintly power- $\Phi$ ; and*
- (ii) *In  $V[h]$ ,  $\kappa$  is faintly power- $\Phi$  but not virtually  $\Phi$ .*

PROOF. We start with (i). Let  $\mathbb{P}_\kappa$  be the Easton support iteration that adds a Cohen subset to every regular  $\lambda < \kappa$ , and let  $g \subseteq \mathbb{P}_\kappa$  be  $V$ -generic. Note that  $\kappa$  remains inaccessible in  $V[g]$ . Fix a regular  $\theta > \kappa$  and let  $\mathbb{Q}_\theta$  be a forcing witnessing that  $\kappa$  is virtually  $\theta$ -measurable.

Since  $\kappa$  is *virtually* measurable we may without loss of generality assume that  $\mathbb{Q}_\theta = \text{Col}(\omega, \theta)$  by applying Countable Embedding Absoluteness ???. Fixing a  $V[g]$ -generic  $h \subseteq \mathbb{Q}_\theta$  we get a transitive  $\mathcal{N} \subseteq V$  and in  $V[h]$  an elementary embedding

$$\pi: H_\theta^V \rightarrow \mathcal{N}$$

with  $\text{crit } \pi = \kappa$ . Let's now work in  $V[g][h] = V[h][g] = V[g \times h]$ , in which we still have access to  $\pi$ . The lifting criterion<sup>17</sup> is trivial for  $\mathbb{P}_\kappa$ , so we get an  $\mathcal{N}$ -generic  $\tilde{g} \subseteq \pi(\mathbb{P}_\kappa)$  and an elementary

$$\pi^+: H_\theta^{V[g]} \rightarrow \mathcal{N}[\tilde{g}]$$

with  $\pi \subseteq \pi^+$ . Note here that without loss of generality  $\pi(\kappa)$  is countable as otherwise we replace  $\mathcal{N}$  by a countable hull, so we can indeed construct such a  $\tilde{g}$ . By elementarity of  $\pi$  it holds that

$$\pi(\mathbb{P}_\kappa) = \mathbb{P}_\kappa * \prod_{\lambda \in [\kappa, \pi(\kappa))} \text{Add}(\lambda, 1), \quad (1)$$

so that  $\mathcal{N}[\tilde{g}] \not\subseteq V$  as it in particular contains a new subset of  $\kappa$ . If  $\Phi = \text{measurable}$  then we're done at this point. For  $\Phi = \text{prestrong}$  we simply note that  $g \in \mathcal{N}[\tilde{g}]$  by (1) so that  $H_\theta^{V[g]} \subseteq \mathcal{N}[\tilde{g}]$  as well, and since  $\pi^+$  lifts  $\pi$  it holds that  $\pi^+(\kappa) = \pi(\kappa) > \theta$  in the  $\Phi = \text{strong}$  case.

---

<sup>17</sup>See Appendix ?? for the definition and characterisations of this criterion.

As for (ii), we simply change  $\mathbb{P}_\kappa$  to only add Cohen subsets to *successor* cardinals  $\lambda < \kappa$ , which means that  $\pi(\mathbb{P}_\kappa)$  doesn't add any subsets of  $\kappa$  and  $\kappa$  thus remains faintly power- $\Phi$ . By choosing  $\theta > \kappa^+$  it *does* add a subset to  $\kappa^+$  however, showing that  $\kappa$  is not virtually  $\Phi$ . ■

Note however, that in contrast to the above separation result, Theorem 1.20 showed that the faintly-virtually distinction vanishes when we're dealing with woodin cardinals.

Our next separation result is concerning the virtually prestrong and virtually strong cardinals.

**Corollary 1.47** (N.). *There exists a virtually rank-into-rank cardinal iff there is an uncountable cardinal  $\theta$  and a virtually  $\theta$ -prestrong cardinal which is not virtually  $\theta$ -strong.*

PROOF. ( $\Leftarrow$ ) is directly from the above Proposition 1.11 and Theorem 1.9.

( $\Rightarrow$ ): Here we have to show that if there exists a virtually rank-into-rank cardinal then there exists a  $\theta > \kappa$  and a virtually  $\theta$ -prestrong cardinal which is not virtually  $\theta$ -strong. Let  $(\kappa, \theta)$  be the lexicographically least pair such that  $\kappa$  is virtually  $\theta$ -rank-into-rank, which trivially makes  $\kappa$  virtually  $\theta$ -prestrong. If  $\kappa$  was also virtually  $\theta$ -strong then it would be  $\Sigma_2$ -reflecting, so that the statement that there exists a virtually rank-into-rank cardinal would reflect down to  $H_\kappa^V$ , contradicting the minimality of  $\kappa$ . ■

Figure 1.2 summarises the separation results along with the results from Section 1.4.

## 1.6 INDESTRUCTIBILITY

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla

Explain the diagram, possibly also in caption




Figure 1.2: Direct implications between virtuals.

ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.