

# Model Theory

## An introduction

Dan Saattrup Nielsen

October 25, 2013

- Introduction
- Basic concepts
- Awesome theorems

- Introduction
- Basic concepts
- *Awesome theorems*

- Introduction
- Basic concepts
- Awesome theorems

- Introduction
- Basic concepts
- Awesome theorems

# So.. what *is* model theory?

- Model theory is about *models* and *theories*
- Classical model theory = algebra + logic
- Syntax and semantics

# So.. what *is* model theory?

- Model theory is about *models* and *theories*
- Classical model theory = algebra + logic
- Syntax and semantics

# So.. what *is* model theory?

- Model theory is about *models* and *theories*
- Classical model theory = algebra + logic
- Syntax and semantics



# So.. what *is* model theory?

- Model theory is about *models* and *theories*
- Classical model theory = algebra + logic
- Syntax and semantics

# Contents

- Introduction
- Basic concepts
- Awesome theorems

## Definition

A *language*  $\mathcal{L}$  is a set of relation symbols  $(R_i)_{i \in I}$ , function symbols  $(f_j)_{j \in J}$  and constant symbols  $(c_k)_{k \in K}$ .

## Definition

A *language*  $\mathcal{L}$  is a set of relation symbols  $(R_i)_{i \in I}$ , function symbols  $(f_j)_{j \in J}$  and constant symbols  $(c_k)_{k \in K}$ .

## Definition

A *language*  $\mathcal{L}$  is a set of relation symbols  $(R_i)_{i \in I}$ , function symbols  $(f_j)_{j \in J}$  and constant symbols  $(c_k)_{k \in K}$ .

## Definition

A *language*  $\mathcal{L}$  is a set of relation symbols  $(R_i)_{i \in I}$ , function symbols  $(f_j)_{j \in J}$  and constant symbols  $(c_k)_{k \in K}$ .

## Definition

Let  $\mathcal{L}$  be a language. Then an  $\mathcal{L}$ -structure  $\mathfrak{M}$  is a set  $M$  along with

- an  $n$ -ary relation  $R_i^{\mathfrak{M}} \subseteq M^n$  for each  $n$ -ary relation symbol  $R_i \in \mathcal{L}$ ,
- an  $n$ -ary function  $f_j^{\mathfrak{M}} : M^n \rightarrow M$  for each  $n$ -ary function symbol  $f_j \in \mathcal{L}$
- a distinguished element  $c_k^{\mathfrak{M}} \in M$  for each constant symbol  $c_k \in \mathcal{L}$ .

## Definition

Let  $\mathcal{L}$  be a language. Then an  $\mathcal{L}$ -structure  $\mathfrak{M}$  is a set  $M$  along with

- an  $n$ -ary relation  $R_i^{\mathfrak{M}} \subseteq M^n$  for each  $n$ -ary relation symbol  $R_i \in \mathcal{L}$ ,
- an  $n$ -ary function  $f_j^{\mathfrak{M}} : M^n \rightarrow M$  for each  $n$ -ary function symbol  $f_j \in \mathcal{L}$
- a distinguished element  $c_k^{\mathfrak{M}} \in M$  for each constant symbol  $c_k \in \mathcal{L}$ .



## Definition

Let  $\mathcal{L}$  be a language. Then an  $\mathcal{L}$ -structure  $\mathfrak{M}$  is a set  $M$  along with

- an  $n$ -ary relation  $R_i^{\mathfrak{M}} \subseteq M^n$  for each  $n$ -ary relation symbol  $R_i \in \mathcal{L}$ ,
- an  $n$ -ary function  $f_j^{\mathfrak{M}} : M^n \rightarrow M$  for each  $n$ -ary function symbol  $f_j \in \mathcal{L}$
- a distinguished element  $c_k^{\mathfrak{M}} \in M$  for each constant symbol  $c_k \in \mathcal{L}$ .

## Definition

Let  $\mathcal{L}$  be a language. Then an  $\mathcal{L}$ -structure  $\mathfrak{M}$  is a set  $M$  along with

- an  $n$ -ary relation  $R_i^{\mathfrak{M}} \subseteq M^n$  for each  $n$ -ary relation symbol  $R_i \in \mathcal{L}$ ,
- an  $n$ -ary function  $f_j^{\mathfrak{M}} : M^n \rightarrow M$  for each  $n$ -ary function symbol  $f_j \in \mathcal{L}$
- a distinguished element  $c_k^{\mathfrak{M}} \in M$  for each constant symbol  $c_k \in \mathcal{L}$ .

## Definition

Let  $\mathcal{L}$  be a language. Then an  $\mathcal{L}$ -structure  $\mathfrak{M}$  is a set  $M$  along with

- an  $n$ -ary relation  $R_i^{\mathfrak{M}} \subseteq M^n$  for each  $n$ -ary relation symbol  $R_i \in \mathcal{L}$ ,
- an  $n$ -ary function  $f_j^{\mathfrak{M}} : M^n \rightarrow M$  for each  $n$ -ary function symbol  $f_j \in \mathcal{L}$
- a distinguished element  $c_k^{\mathfrak{M}} \in M$  for each constant symbol  $c_k \in \mathcal{L}$ .

Let  $\mathcal{L}$  be a language.

## Definition

An  $\mathcal{L}$ -theory  $\mathcal{T}$  is just a set of  $\mathcal{L}$ -sentences.  $\mathcal{T}$  is called *satisfiable* if it has a model.

Let  $\mathcal{L}$  be a language.

## Definition

An  $\mathcal{L}$ -theory  $\mathcal{T}$  is just a set of  $\mathcal{L}$ -sentences.  $\mathcal{T}$  is called *satisfiable* if it has a model.

Let  $\mathcal{L}$  be a language.

## Definition

An  $\mathcal{L}$ -theory  $\mathcal{T}$  is just a set of  $\mathcal{L}$ -sentences.  $\mathcal{T}$  is called *satisfiable* if it has a model.

- Introduction
- Basic concepts
- Awesome theorems

# Completeness (first kind)

## Definition

A *dense linear ordering*  $(M, <)$  is an ordered set on which the ordering  $<$  is

- irreflexive
- transitive
- total
- dense.



# Completeness (first kind)

## Definition

A *dense linear ordering*  $(M, <)$  is an ordered set on which the ordering  $<$  is

- irreflexive
- transitive
- total
- dense.

# Completeness (first kind)

## Definition

A *dense linear ordering*  $(M, <)$  is an ordered set on which the ordering  $<$  is

- irreflexive
- transitive
- total
- dense.

# Completeness (first kind)

## Definition

A *dense linear ordering*  $(M, <)$  is an ordered set on which the ordering  $<$  is

- irreflexive
- transitive
- total
- dense.

# Completeness (first kind)

## Definition

A *dense linear ordering*  $(M, <)$  is an ordered set on which the ordering  $<$  is

- irreflexive
- transitive
- total
- dense.

## Definition

A *dense linear ordering*  $(M, <)$  is an ordered set on which the ordering  $<$  is

- irreflexive
- transitive
- total
- dense.

# Completeness (first kind)

Let  $\mathcal{L}_{strict} := \{<\}$  be the language of strict orderings.

## Proposition

Let  $\mathfrak{M}, \mathfrak{N}$  be two DLO's without endpoints and let  $\sigma$  be an  $\mathcal{L}_{strict}$ -sentence. Then

$$\mathfrak{M} \models \sigma \Leftrightarrow \mathfrak{N} \models \sigma.$$

## Proposition

Let  $\mathfrak{M}$  be a DLO without endpoints. There exists no  $\mathcal{L}_{strict}$ -sentence  $\sigma$  such that  $\mathfrak{M} \models \sigma$  iff the ordering on  $\mathfrak{M}$  is complete.

# Completeness (first kind)

Let  $\mathcal{L}_{strict} := \{<\}$  be the language of strict orderings.

## Proposition

Let  $\mathfrak{M}, \mathfrak{N}$  be two DLO's without endpoints and let  $\sigma$  be an  $\mathcal{L}_{strict}$ -sentence. Then

$$\mathfrak{M} \models \sigma \Leftrightarrow \mathfrak{N} \models \sigma.$$

## Proposition

Let  $\mathfrak{M}$  be a DLO without endpoints. There exists no  $\mathcal{L}_{strict}$ -sentence  $\sigma$  such that  $\mathfrak{M} \models \sigma$  iff the ordering on  $\mathfrak{M}$  is complete.

# Completeness (first kind)

Let  $\mathcal{L}_{strict} := \{<\}$  be the language of strict orderings.

## Proposition

Let  $\mathfrak{M}, \mathfrak{N}$  be two DLO's without endpoints and let  $\sigma$  be an  $\mathcal{L}_{strict}$ -sentence. Then

$$\mathfrak{M} \models \sigma \Leftrightarrow \mathfrak{N} \models \sigma.$$

## Proposition

Let  $\mathfrak{M}$  be a DLO without endpoints. There exists no  $\mathcal{L}_{strict}$ -sentence  $\sigma$  such that  $\mathfrak{M} \models \sigma$  iff the ordering on  $\mathfrak{M}$  is complete.



# Completeness (first kind)

Let  $\mathcal{L}_{strict} := \{<\}$  be the language of strict orderings.

## Proposition

Let  $\mathfrak{M}, \mathfrak{N}$  be two DLO's without endpoints and let  $\sigma$  be an  $\mathcal{L}_{strict}$ -sentence. Then

$$\mathfrak{M} \models \sigma \Leftrightarrow \mathfrak{N} \models \sigma.$$

## Proposition

Let  $\mathfrak{M}$  be a DLO without endpoints. There exists no  $\mathcal{L}_{strict}$ -sentence  $\sigma$  such that  $\mathfrak{M} \models \sigma$  iff the ordering on  $\mathfrak{M}$  is complete.

# Completeness (first kind)

Let  $\mathcal{L}_{strict} := \{<\}$  be the language of strict orderings.

## Proposition

Let  $\mathfrak{M}, \mathfrak{N}$  be two DLO's without endpoints and let  $\sigma$  be an  $\mathcal{L}_{strict}$ -sentence. Then

$$\mathfrak{M} \models \sigma \Leftrightarrow \mathfrak{N} \models \sigma.$$

## Proposition

Let  $\mathfrak{M}$  be a DLO without endpoints. There exists no  $\mathcal{L}_{strict}$ -sentence  $\sigma$  such that  $\mathfrak{M} \models \sigma$  iff the ordering on  $\mathfrak{M}$  is complete.

# Completeness (first kind)

Let  $\mathcal{L}_{strict} := \{<\}$  be the language of strict orderings.

## Proposition

Let  $\mathfrak{M}, \mathfrak{N}$  be two DLO's without endpoints and let  $\sigma$  be an  $\mathcal{L}_{strict}$ -sentence. Then

$$\mathfrak{M} \models \sigma \Leftrightarrow \mathfrak{N} \models \sigma.$$

## Proposition

Let  $\mathfrak{M}$  be a DLO without endpoints. There exists no  $\mathcal{L}_{strict}$ -sentence  $\sigma$  such that  $\mathfrak{M} \models \sigma$  iff the ordering on  $\mathfrak{M}$  is complete.

# Completeness (first kind)

Let  $\mathcal{L}_{strict} := \{<\}$  be the language of strict orderings.

## Proposition

Let  $\mathfrak{M}, \mathfrak{N}$  be two DLO's without endpoints and let  $\sigma$  be an  $\mathcal{L}_{strict}$ -sentence. Then

$$\mathfrak{M} \models \sigma \Leftrightarrow \mathfrak{N} \models \sigma.$$

## Proposition

Let  $\mathfrak{M}$  be a DLO without endpoints. There exists no  $\mathcal{L}_{strict}$ -sentence  $\sigma$  such that  $\mathfrak{M} \models \sigma$  iff the ordering on  $\mathfrak{M}$  is complete.

# Completeness (second kind)

Let  $\mathcal{L}$  be a language.

## Definition

A theory  $\mathcal{T}$  is *complete* if for every  $\mathcal{L}$ -sentence  $\sigma$ , either  $\mathcal{T} \models \sigma$  or  $\mathcal{T} \models \neg\sigma$ .

## Theorem (Vaught's test)

Let  $\mathcal{T}$  be a satisfiable  $\mathcal{L}$ -theory with no finite models and every model of cardinality  $\kappa$  is isomorphic, for some cardinal  $\kappa \geq |\mathcal{L}|$ . Then  $\mathcal{T}$  is complete.

## Corollary

$\text{Vec}_\infty$  is complete in  $\mathcal{L}_{\text{vec}}$ , the language of vector spaces.

# Completeness (second kind)

Let  $\mathcal{L}$  be a language.

## Definition

A theory  $\mathcal{T}$  is *complete* if for every  $\mathcal{L}$ -sentence  $\sigma$ , either  $\mathcal{T} \models \sigma$  or  $\mathcal{T} \models \neg\sigma$ .

## Theorem (Vaught's test)

Let  $\mathcal{T}$  be a satisfiable  $\mathcal{L}$ -theory with no finite models and every model of cardinality  $\kappa$  is isomorphic, for some cardinal  $\kappa \geq |\mathcal{L}|$ . Then  $\mathcal{T}$  is complete.

## Corollary

$\text{Vec}_\infty$  is complete in  $\mathcal{L}_{\text{vec}}$ , the language of vector spaces.

# Completeness (second kind)

Let  $\mathcal{L}$  be a language.

## Definition

A theory  $\mathcal{T}$  is *complete* if for every  $\mathcal{L}$ -sentence  $\sigma$ , either  $\mathcal{T} \models \sigma$  or  $\mathcal{T} \models \neg\sigma$ .

## Theorem (Vaught's test)

Let  $\mathcal{T}$  be a satisfiable  $\mathcal{L}$ -theory with no finite models and every model of cardinality  $\kappa$  is isomorphic, for some cardinal  $\kappa \geq |\mathcal{L}|$ . Then  $\mathcal{T}$  is complete.

## Corollary

$\text{Vec}_\infty$  is complete in  $\mathcal{L}_{\text{vec}}$ , the language of vector spaces.

# Completeness (second kind)

Let  $\mathcal{L}$  be a language.

## Definition

A theory  $\mathcal{T}$  is *complete* if for every  $\mathcal{L}$ -sentence  $\sigma$ , either  $\mathcal{T} \models \sigma$  or  $\mathcal{T} \models \neg\sigma$ .

## Theorem (Vaught's test)

Let  $\mathcal{T}$  be a satisfiable  $\mathcal{L}$ -theory with no finite models and every model of cardinality  $\kappa$  is isomorphic, for some cardinal  $\kappa \geq |\mathcal{L}|$ . Then  $\mathcal{T}$  is complete.

## Corollary

$\text{Vec}_\infty$  is complete in  $\mathcal{L}_{\text{vec}}$ , the language of vector spaces.



# Completeness (second kind)

Let  $\mathcal{L}$  be a language.

## Definition

A theory  $\mathcal{T}$  is *complete* if for every  $\mathcal{L}$ -sentence  $\sigma$ , either  $\mathcal{T} \models \sigma$  or  $\mathcal{T} \models \neg\sigma$ .

## Theorem (Vaught's test)

Let  $\mathcal{T}$  be a satisfiable  $\mathcal{L}$ -theory with no finite models and every model of cardinality  $\kappa$  is isomorphic, for some cardinal  $\kappa \geq |\mathcal{L}|$ .  
Then  $\mathcal{T}$  is complete.

## Corollary

$\text{Vec}_\infty$  is complete in  $\mathcal{L}_{\text{vec}}$ , the language of vector spaces.

# Completeness (second kind)

Let  $\mathcal{L}$  be a language.

## Definition

A theory  $\mathcal{T}$  is *complete* if for every  $\mathcal{L}$ -sentence  $\sigma$ , either  $\mathcal{T} \models \sigma$  or  $\mathcal{T} \models \neg\sigma$ .

## Theorem (Vaught's test)

Let  $\mathcal{T}$  be a satisfiable  $\mathcal{L}$ -theory with no finite models and every model of cardinality  $\kappa$  is isomorphic, for some cardinal  $\kappa \geq |\mathcal{L}|$ . Then  $\mathcal{T}$  is complete.

## Corollary

$\text{Vec}_\infty$  is complete in  $\mathcal{L}_{\text{vec}}$ , the language of vector spaces.

# Completeness (second kind)

Let  $\mathcal{L}$  be a language.

## Definition

A theory  $\mathcal{T}$  is *complete* if for every  $\mathcal{L}$ -sentence  $\sigma$ , either  $\mathcal{T} \models \sigma$  or  $\mathcal{T} \models \neg\sigma$ .

## Theorem (Vaught's test)

Let  $\mathcal{T}$  be a satisfiable  $\mathcal{L}$ -theory with no finite models and every model of cardinality  $\kappa$  is isomorphic, for some cardinal  $\kappa \geq |\mathcal{L}|$ . Then  $\mathcal{T}$  is complete.

## Corollary

$\text{Vec}_\infty$  is complete in  $\mathcal{L}_{\text{vec}}$ , the language of vector spaces.

# Completeness (third kind)

Let  $\mathcal{L}$  be a language and  $\mathcal{T}$  an  $\mathcal{L}$ -theory.

Theorem (Gödel's Completeness Theorem)

Let  $\sigma$  be an  $\mathcal{L}$ -sentence. Then

$$\mathcal{T} \models \sigma \Leftrightarrow \mathcal{T} \vdash \sigma.$$

Theorem (Compactness Theorem)

$\mathcal{T}$  is satisfiable iff every finite  $\Delta \subseteq \mathcal{T}$  is satisfiable.

Corollary

If  $\mathcal{T} \models \sigma$  then  $\Delta \models \sigma$  for some finite  $\Delta \subseteq \mathcal{T}$ .

# Completeness (third kind)

Let  $\mathcal{L}$  be a language and  $\mathcal{T}$  an  $\mathcal{L}$ -theory.

## Theorem (Gödel's Completeness Theorem)

Let  $\sigma$  be an  $\mathcal{L}$ -sentence. Then

$$\mathcal{T} \models \sigma \Leftrightarrow \mathcal{T} \vdash \sigma.$$

## Theorem (Compactness Theorem)

$\mathcal{T}$  is satisfiable iff every finite  $\Delta \subseteq \mathcal{T}$  is satisfiable.

## Corollary

If  $\mathcal{T} \models \sigma$  then  $\Delta \models \sigma$  for some finite  $\Delta \subseteq \mathcal{T}$ .

# Completeness (third kind)

Let  $\mathcal{L}$  be a language and  $\mathcal{T}$  an  $\mathcal{L}$ -theory.

## Theorem (Gödel's Completeness Theorem)

Let  $\sigma$  be an  $\mathcal{L}$ -sentence. Then

$$\mathcal{T} \models \sigma \Leftrightarrow \mathcal{T} \vdash \sigma.$$

## Theorem (Compactness Theorem)

$\mathcal{T}$  is satisfiable iff every finite  $\Delta \subseteq \mathcal{T}$  is satisfiable.

## Corollary

If  $\mathcal{T} \models \sigma$  then  $\Delta \models \sigma$  for some finite  $\Delta \subseteq \mathcal{T}$ .

# Completeness (third kind)

Let  $\mathcal{L}$  be a language and  $\mathcal{T}$  an  $\mathcal{L}$ -theory.

Theorem (Gödel's Completeness Theorem)

Let  $\sigma$  be an  $\mathcal{L}$ -sentence. Then

$$\mathcal{T} \models \sigma \Leftrightarrow \mathcal{T} \vdash \sigma.$$

Theorem (Compactness Theorem)

$\mathcal{T}$  is satisfiable iff every finite  $\Delta \subseteq \mathcal{T}$  is satisfiable.

Corollary

If  $\mathcal{T} \models \sigma$  then  $\Delta \models \sigma$  for some finite  $\Delta \subseteq \mathcal{T}$ .

# Completeness (third kind)

Let  $\mathcal{L}$  be a language and  $\mathcal{T}$  an  $\mathcal{L}$ -theory.

## Theorem (Gödel's Completeness Theorem)

Let  $\sigma$  be an  $\mathcal{L}$ -sentence. Then

$$\mathcal{T} \models \sigma \Leftrightarrow \mathcal{T} \vdash \sigma.$$

## Theorem (Compactness Theorem)

$\mathcal{T}$  is satisfiable iff every finite  $\Delta \subseteq \mathcal{T}$  is satisfiable.

## Corollary

If  $\mathcal{T} \models \sigma$  then  $\Delta \models \sigma$  for some finite  $\Delta \subseteq \mathcal{T}$ .



# Completeness (third kind)

## Lemma

Let  $\sigma$  be a sentence in the language of rings. Then  $\text{ACF}_0 \models \sigma$  if  $\text{ACF}_p \models \sigma$  for all primes  $p$ . In particular  $\mathbb{C} \models \sigma$ .

## Theorem (Ax)

Every injective polynomial  $p : \mathbb{C}^n \rightarrow \mathbb{C}^n$  is surjective.

# Completeness (third kind)

## Lemma

Let  $\sigma$  be a sentence in the language of rings. Then  $\text{ACF}_0 \models \sigma$  if  $\text{ACF}_p \models \sigma$  for all primes  $p$ . In particular  $\mathbb{C} \models \sigma$ .

## Theorem (Ax)

Every injective polynomial  $p : \mathbb{C}^n \rightarrow \mathbb{C}^n$  is surjective.

# Completeness (third kind)

## Lemma

Let  $\sigma$  be a sentence in the language of rings. Then  $\text{ACF}_0 \models \sigma$  if  $\text{ACF}_p \models \sigma$  for all primes  $p$ . In particular  $\mathbb{C} \models \sigma$ .

## Theorem (Ax)

Every injective polynomial  $p : \mathbb{C}^n \rightarrow \mathbb{C}^n$  is surjective.

# Completeness (third kind)

## Lemma

Let  $\sigma$  be a sentence in the language of rings. Then  $\text{ACF}_0 \models \sigma$  if  $\text{ACF}_p \models \sigma$  for all primes  $p$ . In particular  $\mathbb{C} \models \sigma$ .

## Theorem (Ax)

Every injective polynomial  $p : \mathbb{C}^n \rightarrow \mathbb{C}^n$  is surjective.