

# Games and Ramsey-like cardinals

SET THEORY TODAY CONFERENCE, VIENNA

Dan Saattrup Nielsen  
University of Bristol

September 14, 2018

Let's play a game  $\mathcal{G}_\alpha^\theta(\kappa)$

I

II

Let's play a game  $\mathcal{G}_\alpha^\theta(\kappa)$

I  $\mathcal{M}_0$   
II

①  $\mathcal{M}_\eta \prec H_\theta$  is a  $\kappa$ -sized model of  $\text{ZFC}^-$  containing  $\kappa+1$

Let's play a game  $\mathcal{G}_\alpha^\theta(\kappa)$

I  $\mathcal{M}_0$   
II  $\mu_0$

- ①  $\mathcal{M}_\eta \prec H_\theta$  is a  $\kappa$ -sized model of  $\text{ZFC}^-$  containing  $\kappa+1$
- ②  $\mu_\eta$  is an  $\mathcal{M}_\eta$ -normal  $\mathcal{M}_\eta$ -measure on  $\kappa$  such that  $\text{Ult}(\mathcal{M}_\eta, \mu_\eta)$  is wellfounded

Let's play a game  $\mathcal{G}_\alpha^\theta(\kappa)$

$$\begin{array}{ll} \text{I} & \mathcal{M}_0 \quad \mathcal{M}_1 \\ \text{II} & \mu_0 \end{array}$$

- ①  $\mathcal{M}_\eta \prec H_\theta$  is a  $\kappa$ -sized model of  $\text{ZFC}^-$  containing  $\kappa+1$
- ②  $\mu_\eta$  is an  $\mathcal{M}_\eta$ -normal  $\mathcal{M}_\eta$ -measure on  $\kappa$  such that  $\text{Ult}(\mathcal{M}_\eta, \mu_\eta)$  is wellfounded
- ③ The  $\mathcal{M}_\eta$ 's and  $\mu_\eta$ 's are  $\subseteq$ -increasing

Let's play a game  $\mathcal{G}_\alpha^\theta(\kappa)$

$$\begin{array}{lll} \text{I} & \mathcal{M}_0 & \mathcal{M}_1 \\ \text{II} & & \mu_0 \quad \mu_1 \end{array}$$

- ①  $\mathcal{M}_\eta \prec H_\theta$  is a  $\kappa$ -sized model of  $\text{ZFC}^-$  containing  $\kappa+1$
- ②  $\mu_\eta$  is an  $\mathcal{M}_\eta$ -normal  $\mathcal{M}_\eta$ -measure on  $\kappa$  such that  $\text{Ult}(\mathcal{M}_\eta, \mu_\eta)$  is wellfounded
- ③ The  $\mathcal{M}_\eta$ 's and  $\mu_\eta$ 's are  $\subseteq$ -increasing
- ④ We take **unions** at limit rounds

Let's play a game  $\mathcal{G}_\alpha^\theta(\kappa)$

$$\begin{array}{ccccccc} \text{I} & \mathcal{M}_0 & & \mathcal{M}_1 & & \cdots & \\ \text{II} & & \mu_0 & & \mu_1 & & \cdots \end{array}$$

- ①  $\mathcal{M}_\eta \prec H_\theta$  is a  $\kappa$ -sized model of  $\text{ZFC}^-$  containing  $\kappa+1$
- ②  $\mu_\eta$  is an  $\mathcal{M}_\eta$ -normal  $\mathcal{M}_\eta$ -measure on  $\kappa$  such that  $\text{Ult}(\mathcal{M}_\eta, \mu_\eta)$  is wellfounded
- ③ The  $\mathcal{M}_\eta$ 's and  $\mu_\eta$ 's are  $\subseteq$ -increasing
- ④ We take unions at limit rounds

Let's play a game  $\mathcal{G}_\alpha^\theta(\kappa)$

$$\begin{array}{ccccccc} \text{I} & \mathcal{M}_0 & & \mathcal{M}_1 & & \cdots & & \mathcal{M}_\alpha \\ \text{II} & & \mu_0 & & \mu_1 & & \cdots & \end{array}$$

- ①  $\mathcal{M}_\eta \prec H_\theta$  is a  $\kappa$ -sized model of  $\text{ZFC}^-$  containing  $\kappa+1$
- ②  $\mu_\eta$  is an  $\mathcal{M}_\eta$ -normal  $\mathcal{M}_\eta$ -measure on  $\kappa$  such that  $\text{Ult}(\mathcal{M}_\eta, \mu_\eta)$  is wellfounded
- ③ The  $\mathcal{M}_\eta$ 's and  $\mu_\eta$ 's are  $\subseteq$ -increasing
- ④ We take unions at limit rounds
- ⑤ The game lasts for  $\alpha+1$  rounds



Let's play a game  $\mathcal{G}_\alpha^\theta(\kappa)$

$$\begin{array}{ccccccc} \text{I} & \mathcal{M}_0 & & \mathcal{M}_1 & & \cdots & & \mathcal{M}_\alpha \\ \text{II} & & \mu_0 & & \mu_1 & & \cdots & & \mu_\alpha \end{array}$$

- ①  $\mathcal{M}_\eta \prec H_\theta$  is a  $\kappa$ -sized model of  $\text{ZFC}^-$  containing  $\kappa+1$
- ②  $\mu_\eta$  is an  $\mathcal{M}_\eta$ -normal  $\mathcal{M}_\eta$ -measure on  $\kappa$  such that  $\text{Ult}(\mathcal{M}_\eta, \mu_\eta)$  is wellfounded
- ③ The  $\mathcal{M}_\eta$ 's and  $\mu_\eta$ 's are  $\subseteq$ -increasing
- ④ We take unions at limit rounds
- ⑤ The game lasts for  $\alpha+1$  rounds
- ⑥ **Player II wins** iff they can continue playing all rounds

Let's play a game  $\mathcal{G}_\alpha^\theta(\kappa)$

$$\begin{array}{ccccccc} \text{I} & \mathcal{M}_0 & & \mathcal{M}_1 & & \cdots & & \mathcal{M}_\alpha \\ \text{II} & & \mu_0 & & \mu_1 & & \cdots & & \mu_\alpha \end{array}$$

- ①  $\mathcal{M}_\eta \prec H_\theta$  is a  $\kappa$ -sized model of  $\text{ZFC}^-$  containing  $\kappa+1$
- ②  $\mu_\eta$  is an  $\mathcal{M}_\eta$ -normal  $\mathcal{M}_\eta$ -measure on  $\kappa$  such that  $\text{Ult}(\mathcal{M}_\eta, \mu_\eta)$  is wellfounded
- ③ The  $\mathcal{M}_\eta$ 's and  $\mu_\eta$ 's are  $\subseteq$ -increasing
- ④ We take unions at limit rounds
- ⑤ The game lasts for  $\alpha+1$  rounds
- ⑥ Player II wins iff they can continue playing all rounds
- ⑦ **IMPORTANT REMARK:** The  $\mathcal{M}_\eta$ 's are **not** necessarily transitive!

# Yet another large cardinal notion?!

Definition (Holy-Schlicht '18)

# Yet another large cardinal notion?!

## Definition (Holy-Schlicht '18)

For  $\alpha \leq \kappa$ , a cardinal  $\kappa$  is **strategic  $\alpha$ -Ramsey** if, for all regular  $\theta > \kappa$ , player II has a strategy in  $\mathcal{G}_\alpha^\theta(\kappa)$  which is always winning.

# Yet another large cardinal notion?!

## Definition

For  $\alpha \leq \kappa$ , a cardinal  $\kappa$  is **weakly strategic  $\alpha$ -Ramsey** if, for all regular  $\theta > \kappa$ , player II has a strategy in  $\mathcal{G}_\alpha^\theta(\kappa)$  which is always winning **the first  $\alpha$  rounds**.

# What are the **finite versions** good for?

## Definition

Let  $\kappa$  be a cardinal. Then

# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow (\kappa)^2$

# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow (\kappa)^2$ , i.e. that for every colouring  $c : [\kappa]^2 \rightarrow \{0, 1\}$  of 2-element subsets of  $\kappa$  into two colours there's a **homogeneous**  $X \subseteq \kappa$  of size  $\kappa$ ; that is, all pairs of  $X$  have the same colour:  $|c"[X]^2| = 1$ .



# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow (\kappa)^2$ , i.e. that for every colouring  $c : [\kappa]^2 \rightarrow \{0, 1\}$  of 2-element subsets of  $\kappa$  into two colours there's a **homogeneous**  $X \subseteq \kappa$  of size  $\kappa$ ; that is, all pairs of  $X$  have the same colour:  $|c"[X]^2| = 1$ .

# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow (\kappa)^2$ , i.e. that for every colouring  $c : [\kappa]^2 \rightarrow \{0, 1\}$  of 2-element subsets of  $\kappa$  into two colours there's a **homogeneous**  $X \subseteq \kappa$  of size  $\kappa$ ; that is, all pairs of  $X$  have the same colour:  $|c"[X]^2| = 1$ .

# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow (\kappa)^2$ , i.e. that for every colouring  $c : [\kappa]^2 \rightarrow \{0, 1\}$  of 2-element subsets of  $\kappa$  into two colours there's a **homogeneous**  $X \subseteq \kappa$  of size  $\kappa$ ; that is, all pairs of  $X$  have the same colour:  $|c"[X]^2| = 1$ .

# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow ([\kappa]^\kappa)^2$ , i.e. that for every colouring  $c : [\kappa]^2 \rightarrow \{0, 1\}$  of 2-element subsets of  $\kappa$  into two colours there's a **homogeneous**  $X \in [\kappa]^\kappa$ ; that is, all pairs of  $X$  have the same colour:  $|c''[X]^2| = 1$ .

# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow ([\kappa]^\kappa)^2$

# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow ([\kappa]^\kappa)^2$
- $\kappa$  is **ineffable** if  $\kappa \rightarrow (\{\text{stationary subsets of } \kappa\})^2$

# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow ([\kappa]^\kappa)^2$
- $\kappa$  is **ineffable** if  $\kappa \rightarrow (\{\text{stationary subsets of } \kappa\})^2$
- $\kappa$  is **completely ineffable** if there's a non-empty upwards-closed set  $R \subseteq \mathcal{P}(\kappa)$  consisting of stationary subsets of  $\kappa$ , such that  $A \rightarrow (R)^2$  for every  $A \in R$

# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow ([\kappa]^\kappa)^2$
- $\kappa$  is **ineffable** if  $\kappa \rightarrow (\{\text{stationary subsets of } \kappa\})^2$
- $\kappa$  is **completely ineffable** if there's a non-empty upwards-closed set  $R \subseteq \mathcal{P}(\kappa)$  consisting of stationary subsets of  $\kappa$ , such that  $A \rightarrow (R)^2$  for every  $A \in R$

## Theorem (Abramson-Harrington-Kleinberg-Zwicker '77)

Let  $\kappa = \kappa^{<\kappa}$  be a cardinal. Then



# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow ([\kappa]^\kappa)^2$
- $\kappa$  is **ineffable** if  $\kappa \rightarrow (\{\text{stationary subsets of } \kappa\})^2$
- $\kappa$  is **completely ineffable** if there's a non-empty upwards-closed set  $R \subseteq \mathcal{P}(\kappa)$  consisting of stationary subsets of  $\kappa$ , such that  $A \rightarrow (R)^2$  for every  $A \in R$

## Theorem (Abramson-Harrington-Kleinberg-Zwicker '77)

Let  $\kappa = \kappa^{<\kappa}$  be a cardinal. Then

- $\kappa$  is strategic 0-Ramsey  $\Leftrightarrow \kappa$  is weakly compact

# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow ([\kappa]^\kappa)^2$
- $\kappa$  is **ineffable** if  $\kappa \rightarrow (\{\text{stationary subsets of } \kappa\})^2$
- $\kappa$  is **completely ineffable** if there's a non-empty upwards-closed set  $R \subseteq \mathcal{P}(\kappa)$  consisting of stationary subsets of  $\kappa$ , such that  $A \rightarrow (R)^2$  for every  $A \in R$

## Theorem (Abramson-Harrington-Kleinberg-Zwicker '77)

Let  $\kappa = \kappa^{<\kappa}$  be a cardinal. Then

- $\kappa$  is strategic 0-Ramsey  $\Leftrightarrow \kappa$  is weakly compact
- $\kappa$  is strategic 1-Ramsey  $\Rightarrow \kappa$  is ineffable ( $\Rightarrow \kappa$  is strategic 0-Ramsey)

# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow ([\kappa]^\kappa)^2$
- $\kappa$  is **ineffable** if  $\kappa \rightarrow (\{\text{stationary subsets of } \kappa\})^2$
- $\kappa$  is **completely ineffable** if there's a non-empty upwards-closed set  $R \subseteq \mathcal{P}(\kappa)$  consisting of stationary subsets of  $\kappa$ , such that  $A \rightarrow (R)^2$  for every  $A \in R$

## Theorem (N.)

A cardinal  $\kappa$  is weakly strategic  $\omega$ -Ramsey iff  $\kappa$  is completely ineffable.

# What are the finite versions good for?

## Definition

Let  $\kappa$  be a cardinal. Then

- $\kappa$  is **weakly compact** if  $\kappa \rightarrow ([\kappa]^\kappa)^2$
- $\kappa$  is **ineffable** if  $\kappa \rightarrow (\{\text{stationary subsets of } \kappa\})^2$
- $\kappa$  is **completely ineffable** if there's a non-empty upwards-closed set  $R \subseteq \mathcal{P}(\kappa)$  consisting of stationary subsets of  $\kappa$ , such that  $A \rightarrow (R)^2$  for every  $A \in R$

## Theorem (N.)

A cardinal  $\kappa$  is weakly strategic  $\omega$ -Ramsey iff  $\kappa$  is completely ineffable.

The proof uses the previous theorem as the base case. In the “ $\Leftarrow$ ” direction, care must be taken at successor stages to ensure that the measures cohere.

What are the **countable versions** good for?

Definition (Gitman-Schindler '16; original definition from Schindler '00)

A cardinal  $\kappa$  is **remarkable** if

# What are the countable versions good for?


Definition (Gitman-Schindler '16; original definition from Schindler '00)

A cardinal  $\kappa$  is **remarkable** if to every regular  $\lambda > \kappa$  there is a regular  $\nu > \lambda$  and a transitive  $M$  closed under  $\lambda$ -sequences such that, in a generic extension, there's an elementary embedding  $\pi : H_\nu^V \rightarrow M$  with  $\text{crit } \pi = \kappa$  and  $\pi(\kappa) > \lambda$ .

# What are the countable versions good for?

Definition (Gitman-Schindler '16; original definition from Schindler '00)

A cardinal  $\kappa$  is **remarkable** if to every regular  $\lambda > \kappa$  there is a regular  $\nu > \lambda$  and a transitive  $M$  closed under  $\lambda$ -sequences such that, in a generic extension, there's an elementary embedding  $\pi : H_\nu^V \rightarrow M$  with  $\text{crit } \pi = \kappa$  and  $\pi(\kappa) > \lambda$ .



# What are the countable versions good for?

## Definition (Gitman-Schindler '16; original definition from Schindler '00)

A cardinal  $\kappa$  is **remarkable** if to every regular  $\lambda > \kappa$  there is a regular  $\nu > \lambda$  and a transitive  $M$  closed under  $\lambda$ -sequences such that, in a generic extension, there's an elementary embedding  $\pi : H_\nu^V \rightarrow M$  with  $\text{crit } \pi = \kappa$  and  $\pi(\kappa) > \lambda$ .

## Proposition (Schindler '00)

Remarkable cardinals are downwards absolute to  $L$ .



# What are the countable versions good for?

## Definition (Gitman-Schindler '16; original definition from Schindler '00)

A cardinal  $\kappa$  is **remarkable** if to every regular  $\lambda > \kappa$  there is a regular  $\nu > \lambda$  and a transitive  $M$  closed under  $\lambda$ -sequences such that, in a generic extension, there's an elementary embedding  $\pi : H_\nu^V \rightarrow M$  with  $\text{crit } \pi = \kappa$  and  $\pi(\kappa) > \lambda$ .

## Theorem(s)

The existence of a remarkable cardinal is equiconsistent with..

# What are the countable versions good for?

## Definition (Gitman-Schindler '16; original definition from Schindler '00)

A cardinal  $\kappa$  is **remarkable** if to every regular  $\lambda > \kappa$  there is a regular  $\nu > \lambda$  and a transitive  $M$  closed under  $\lambda$ -sequences such that, in a generic extension, there's an elementary embedding  $\pi : H_\nu^V \rightarrow M$  with  $\text{crit } \pi = \kappa$  and  $\pi(\kappa) > \lambda$ .

## Theorem(s)

The existence of a remarkable cardinal is equiconsistent with..

- 1 (Schindler '00) proper forcing cannot change the theory of  $L(\mathbb{R})$

# What are the countable versions good for?

## Definition (Gitman-Schindler '16; original definition from Schindler '00)

A cardinal  $\kappa$  is **remarkable** if to every regular  $\lambda > \kappa$  there is a regular  $\nu > \lambda$  and a transitive  $M$  closed under  $\lambda$ -sequences such that, in a generic extension, there's an elementary embedding  $\pi : H_\nu^V \rightarrow M$  with  $\text{crit } \pi = \kappa$  and  $\pi(\kappa) > \lambda$ .

## Theorem(s)

The existence of a remarkable cardinal is equiconsistent with..

- 1 (Schindler '00) proper forcing cannot change the theory of  $L(\mathbb{R})$
- 2 (Schindler '04) semi-proper forcing cannot change the theory of  $L(\mathbb{R})$

# What are the countable versions good for?

## Definition (Gitman-Schindler '16; original definition from Schindler '00)

A cardinal  $\kappa$  is **remarkable** if to every regular  $\lambda > \kappa$  there is a regular  $\nu > \lambda$  and a transitive  $M$  closed under  $\lambda$ -sequences such that, in a generic extension, there's an elementary embedding  $\pi : H_\nu^V \rightarrow M$  with  $\text{crit } \pi = \kappa$  and  $\pi(\kappa) > \lambda$ .

## Theorem(s)

The existence of a remarkable cardinal is equiconsistent with..

- 1 (Schindler '00) proper forcing cannot change the theory of  $L(\mathbb{R})$
- 2 (Schindler '04) semi-proper forcing cannot change the theory of  $L(\mathbb{R})$
- 3 (Cheng-Schindler '15)  $3^{\text{rd}}$  order number theory + Harrington's Principle ("there is a real  $x$  such that every  $x$ -admissible ordinal is an  $L$ -cardinal")

# What are the countable versions good for?

Definition (Gitman-Schindler '16; original definition from Schindler '00)

A cardinal  $\kappa$  is **remarkable** if to every regular  $\lambda > \kappa$  there is a regular  $\nu > \lambda$  and a transitive  $M$  closed under  $\lambda$ -sequences such that, in a generic extension, there's an elementary embedding  $\pi : H_\nu^V \rightarrow M$  with  $\text{crit } \pi = \kappa$  and  $\pi(\kappa) > \lambda$ .

Theorem (Schindler-N.)

Every remarkable cardinal is strategic  $\omega$ -Ramsey, and if  $\kappa$  is strategic  $\omega$ -Ramsey then either  $\kappa$  is remarkable in  $L$  or

$L_\kappa \models$  There is a proper class of strategic  $\omega$ -Ramseys.

# What are the countable versions good for?

## Definition (Gitman-Schindler '16; original definition from Schindler '00)

A cardinal  $\kappa$  is **remarkable** if to every regular  $\lambda > \kappa$  there is a regular  $\nu > \lambda$  and a transitive  $M$  closed under  $\lambda$ -sequences such that, in a generic extension, there's an elementary embedding  $\pi : H_\nu^V \rightarrow M$  with  $\text{crit } \pi = \kappa$  and  $\pi(\kappa) > \lambda$ .

## Theorem (Schindler-N.)

Every remarkable cardinal is strategic  $\omega$ -Ramsey, and if  $\kappa$  is strategic  $\omega$ -Ramsey then either  $\kappa$  is remarkable in  $L$  or

$$L_\kappa \models \text{There is a proper class of strategic } \omega\text{-Ramseys.}$$

Consequently, strategic  $\omega$ -Ramseys are equiconsistent with remarkables.

# What are the countable versions good for?

## Definition (Gitman-Schindler '16; original definition from Schindler '00)

A cardinal  $\kappa$  is **remarkable** if to every regular  $\lambda > \kappa$  there is a regular  $\nu > \lambda$  and a transitive  $M$  closed under  $\lambda$ -sequences such that, in a generic extension, there's an elementary embedding  $\pi : H_\nu^V \rightarrow M$  with  $\text{crit } \pi = \kappa$  and  $\pi(\kappa) > \lambda$ .

## Theorem (Schindler-N.)

Every remarkable cardinal is strategic  $\omega$ -Ramsey, and if  $\kappa$  is strategic  $\omega$ -Ramsey then either  $\kappa$  is remarkable in  $L$  or

$$L_\kappa \models \text{There is a proper class of strategic } \omega\text{-Ramseys.}$$

The proof goes via the notion of a so-called *virtually measurable cardinal*.

What are the **uncountable versions** good for?



# What are the uncountable versions good for?

## Proposition

Every measurable cardinal  $\kappa$  is strategic  $\kappa$ -Ramsey.

# What are the uncountable versions good for?

## Proposition

Every measurable cardinal  $\kappa$  is strategic  $\kappa$ -Ramsey.

**Proof:** In  $\mathcal{G}_\kappa^\theta(\kappa)$  simply play the measure on  $\kappa$ . □

# What are the uncountable versions good for?

## Proposition

Every measurable cardinal  $\kappa$  is strategic  $\kappa$ -Ramsey.

# What are the uncountable versions good for?

## Proposition

Every measurable cardinal  $\kappa$  is strategic  $\kappa$ -Ramsey.

## Theorem (Welch '18)

Every strategic  $\omega_1$ -Ramsey cardinal  $\kappa$  is measurable in the core model  $K$  below the sharp of a strong cardinal,  $0^\sharp$ .

# What are the uncountable versions good for?

## Proposition

Every measurable cardinal  $\kappa$  is strategic  $\kappa$ -Ramsey.

## Theorem (Welch '18)

Every strategic  $\omega_1$ -Ramsey cardinal  $\kappa$  is measurable in the core model  $K$  below the sharp of a strong cardinal,  $0^\sharp$ .

Proof sketch.

# What are the uncountable versions good for?

## Proposition

Every measurable cardinal  $\kappa$  is strategic  $\kappa$ -Ramsey.

## Theorem (Welch '18)

Every strategic  $\omega_1$ -Ramsey cardinal  $\kappa$  is measurable in the core model  $K$  below the sharp of a strong cardinal,  $0^\sharp$ .

**Proof sketch.** Jump to  $V^{\text{Col}(\omega_1, \kappa^{+K})}$ , let  $\eta_\alpha \rightarrow \kappa^{+K}$  and let player I in  $\mathcal{G}_{\omega_1}^\theta(\kappa)^V$  play models

$$\mathcal{M}_\alpha := K|_{\eta_\alpha}$$

Player II follows their winning strategy.

# What are the uncountable versions good for?

## Proposition

Every measurable cardinal  $\kappa$  is strategic  $\kappa$ -Ramsey.

## Theorem (Welch '18)

Every strategic  $\omega_1$ -Ramsey cardinal  $\kappa$  is measurable in the core model  $K$  below the sharp of a strong cardinal,  $0^\sharp$ .

**Proof sketch.** Jump to  $V^{\text{Col}(\omega_1, \kappa^{+K})}$ , let  $\eta_\alpha \rightarrow \kappa^{+K}$  and let player I in  $\mathcal{G}_{\omega_1}^\theta(\kappa)^V$  play models

$$\mathcal{M}_\alpha := \text{Hull}^{H_\theta^V}(K|\eta_\alpha).$$

Player II follows their winning strategy.

# What are the uncountable versions good for?

## Proposition

Every measurable cardinal  $\kappa$  is strategic  $\kappa$ -Ramsey.

## Theorem (Welch '18)

Every strategic  $\omega_1$ -Ramsey cardinal  $\kappa$  is measurable in the core model  $K$  below the sharp of a strong cardinal,  $0^\sharp$ .

**Proof sketch.** Jump to  $V^{\text{Col}(\omega_1, \kappa^{+K})}$ , let  $\eta_\alpha \rightarrow \kappa^{+K}$  and let player I in  $\mathcal{G}_{\omega_1}^\theta(\kappa)^V$  play models

$$\mathcal{M}_\alpha := \text{Hull}^{H_\theta^V}(K|\eta_\alpha).$$

Player II follows their winning strategy.

The final measure  $\mu_{\omega_1}$  is then a  $K$ -measure and we can wlog assume that  $\mu_{\omega_1}$  is countably complete and weakly amenable.



# What are the uncountable versions good for?

## Proposition

Every measurable cardinal  $\kappa$  is strategic  $\kappa$ -Ramsey.

## Theorem (Welch '18)

Every strategic  $\omega_1$ -Ramsey cardinal  $\kappa$  is measurable in the core model  $K$  below the sharp of a strong cardinal,  $0^\sharp$ .

**Proof sketch.** Jump to  $V^{\text{Col}(\omega_1, \kappa^{+K})}$ , let  $\eta_\alpha \rightarrow \kappa^{+K}$  and let player I in  $\mathcal{G}_{\omega_1}^\theta(\kappa)^V$  play models

$$\mathcal{M}_\alpha := \text{Hull}^{H_\theta^V}(K|\eta_\alpha).$$

Player II follows their winning strategy.

The final measure  $\mu_{\omega_1}$  is then a  $K$ -measure and we can wlog assume that  $\mu_{\omega_1}$  is countably complete and weakly amenable.

Then, since we're below  $0^\sharp$ , inner model theory magic (“the beaver argument”) implies that  $\mu_{\omega_1} \in K$  and we're done. □

# What are the uncountable versions good for?

## Proposition

Every measurable cardinal  $\kappa$  is strategic  $\kappa$ -Ramsey.

## Theorem (Welch '18)

Every strategic  $\omega_1$ -Ramsey cardinal  $\kappa$  is measurable in the core model  $K$  below the sharp of a strong cardinal,  $0^\sharp$ .

Strategic  $\omega_1$ -Ramseys are therefore equiconsistent with measurables.

# What are the uncountable versions good for?

## Proposition

Every measurable cardinal  $\kappa$  is strategic  $\kappa$ -Ramsey.

## Theorem (Schindler '18)

Every strategic  $\omega_1$ -Ramsey cardinal  $\kappa$  is measurable in the core model  $K$  below a Woodin cardinal.

Strategic  $\omega_1$ -Ramseys are therefore equiconsistent with measurables.

# What are the uncountable versions good for?

## Proposition

Every measurable cardinal  $\kappa$  is strategic  $\kappa$ -Ramsey.

## Theorem (Schindler '18)

Every strategic  $\omega_1$ -Ramsey cardinal  $\kappa$  is measurable in the core model  $K$  below a Woodin cardinal.

The proof is similar, but different inner model theory magic is used.

So what happens between  $\omega$  and  $\omega_1$ ?

### Definition

A cardinal  $\kappa$  is **Ramsey** if  $\kappa \rightarrow (\kappa)^{<\omega}$ .

# So what happens between $\omega$ and $\omega_1$ ?

## Definition

A cardinal  $\kappa$  is **Ramsey** if  $\kappa \rightarrow (\kappa)^{<\omega}$ .

## Facts

- 1 If there exists a Ramsey cardinal then  $V \neq L$ .

# So what happens between $\omega$ and $\omega_1$ ?

## Definition

A cardinal  $\kappa$  is **Ramsey** if  $\kappa \rightarrow (\kappa)^{<\omega}$ .

## Facts

- 1 If there exists a Ramsey cardinal then  $V \neq L$ .
- 2 Measurable cardinals are Ramsey limits of Ramsey cardinals.

# So what happens between $\omega$ and $\omega_1$ ?

## Definition

A cardinal  $\kappa$  is **Ramsey** if  $\kappa \rightarrow (\kappa)^{<\omega}$ .

## Facts

- 1 If there exists a Ramsey cardinal then  $V \neq L$ .
- 2 Measurable cardinals are Ramsey limits of Ramsey cardinals.

## Theorem (Mitchell '79, Dodd '82)

A cardinal  $\kappa$  is Ramsey iff every  $A \subseteq \kappa$  can be put into a  $\kappa$ -sized model  $M \models \text{ZFC}^-$  containing  $\kappa+1$  such that there exists a weakly amenable countably complete  $M$ -measure on  $\kappa$ .



# So what happens between $\omega$ and $\omega_1$ ?

## Definition

A cardinal  $\kappa$  is **Ramsey** if  $\kappa \rightarrow (\kappa)^{<\omega}$ .

## Facts

- 1 If there exists a Ramsey cardinal then  $V \neq L$ .
- 2 Measurable cardinals are Ramsey limits of Ramsey cardinals.

## Theorem (Mitchell '79, Dodd '82)

A cardinal  $\kappa$  is Ramsey iff every  $A \subseteq \kappa$  can be put into a  $\kappa$ -sized model  $M \models \text{ZFC}^-$  containing  $\kappa+1$  such that there exists a weakly amenable countably complete  $M$ -measure on  $\kappa$ .

## Proposition (N.)

Strategic  $(\omega+1)$ -Ramsey cardinals are Ramsey limits of Ramsey cardinals.

## So what happens between $\omega$ and $\omega_1$ ?

### Theorem (Mitchell '79, Dodd '82)

A cardinal  $\kappa$  is Ramsey iff every  $A \subseteq \kappa$  can be put into a  $\kappa$ -sized model  $M \models \text{ZFC}^-$  containing  $\kappa+1$  such that there exists a weakly amenable countably complete  $M$ -measure on  $\kappa$ .

### Proposition (N.)

Strategic  $(\omega+1)$ -Ramsey cardinals are Ramsey limits of Ramsey cardinals.

## So what happens between $\omega$ and $\omega_1$ ?

### Theorem (Mitchell '79, Dodd '82)

A cardinal  $\kappa$  is Ramsey iff every  $A \subseteq \kappa$  can be put into a  $\kappa$ -sized model  $M \models \text{ZFC}^-$  containing  $\kappa+1$  such that there exists a weakly amenable countably complete  $M$ -measure on  $\kappa$ .

### Proposition (N.)

Strategic  $(\omega+1)$ -Ramsey cardinals are Ramsey limits of Ramsey cardinals.

**Proof sketch.** Whenever  $\vec{\mathcal{M}}_\alpha * \vec{\mu}_\alpha$  is a play of  $\mathcal{G}_{\omega+1}^\theta(\kappa)$  then

$(\mathcal{M}_{\omega+1}, \in, \mu_{\omega+1}) \models \mu_{\omega+1}$  is countably complete.

## So what happens between $\omega$ and $\omega_1$ ?

### Theorem (Mitchell '79, Dodd '82)

A cardinal  $\kappa$  is Ramsey iff every  $A \subseteq \kappa$  can be put into a  $\kappa$ -sized model  $M \models \text{ZFC}^-$  containing  $\kappa+1$  such that there exists a weakly amenable countably complete  $M$ -measure on  $\kappa$ .

### Proposition (N.)

Strategic  $(\omega+1)$ -Ramsey cardinals are Ramsey limits of Ramsey cardinals.

**Proof sketch.** Whenever  $\vec{\mathcal{M}}_\alpha * \vec{\mu}_\alpha$  is a play of  $\mathcal{G}_{\omega+1}^\theta(\kappa)$  then

$$(\mathcal{M}_{\omega+1}, \in, \mu_{\omega+1}) \models \mu_{\omega+1} \text{ is countably complete.}$$

Then  $\mu_\omega \subseteq \mu_{\omega+1}$  implies that, wlog,

$\mathcal{M}_{\omega+1} \models \mu_\omega$  is countably complete and weakly amenable,  
making  $\mu_\omega$  countably complete and weakly amenable by elementarity.

## So what happens between $\omega$ and $\omega_1$ ?

### Theorem (Mitchell '79, Dodd '82)

A cardinal  $\kappa$  is Ramsey iff every  $A \subseteq \kappa$  can be put into a  $\kappa$ -sized model  $M \models \text{ZFC}^-$  containing  $\kappa+1$  such that there exists a weakly amenable countably complete  $M$ -measure on  $\kappa$ .

### Proposition (N.)

Strategic  $(\omega+1)$ -Ramsey cardinals are Ramsey limits of Ramsey cardinals.

**Proof sketch.** Whenever  $\vec{\mathcal{M}}_\alpha * \vec{\mu}_\alpha$  is a play of  $\mathcal{G}_{\omega+1}^\theta(\kappa)$  then

$$(\mathcal{M}_{\omega+1}, \in, \mu_{\omega+1}) \models \mu_{\omega+1} \text{ is countably complete.}$$

Then  $\mu_\omega \subseteq \mu_{\omega+1}$  implies that, wlog,

$$\mathcal{M}_{\omega+1} \models \mu_\omega \text{ is countably complete and weakly amenable,}$$

making  $\mu_\omega$  countably complete and weakly amenable by elementarity.

So  $\kappa$  is Ramsey.

So what happens between  $\omega$  and  $\omega_1$ ?

Proposition (N.)

Strategic  $(\omega+1)$ -Ramsey cardinals are Ramsey limits of Ramsey cardinals.

**Proof sketch.** Whenever  $\vec{\mathcal{M}}_\alpha * \vec{\mu}_\alpha$  is a play of  $\mathcal{G}_{\omega+1}^\theta(\kappa)$  then

$$(\mathcal{M}_{\omega+1}, \in, \mu_{\omega+1}) \models \mu_{\omega+1} \text{ is countably complete.}$$

Then  $\mu_\omega \subseteq \mu_{\omega+1}$  implies that, wlog,

$$\mathcal{M}_{\omega+1} \models \mu_\omega \text{ is countably complete and weakly amenable,}$$

making  $\mu_\omega$  countably complete and weakly amenable by elementarity.

So  $\kappa$  is Ramsey.

## So what happens between $\omega$ and $\omega_1$ ?

### Proposition (N.)

Strategic  $(\omega+1)$ -Ramsey cardinals are Ramsey limits of Ramsey cardinals.

**Proof sketch.** Whenever  $\vec{\mathcal{M}}_\alpha * \vec{\mu}_\alpha$  is a play of  $\mathcal{G}_{\omega+1}^\theta(\kappa)$  then

$$(\mathcal{M}_{\omega+1}, \in, \mu_{\omega+1}) \models \mu_{\omega+1} \text{ is countably complete.}$$

Then  $\mu_\omega \subseteq \mu_{\omega+1}$  implies that, wlog,

$$\mathcal{M}_{\omega+1} \models \mu_\omega \text{ is countably complete and weakly amenable,}$$

making  $\mu_\omega$  countably complete and weakly amenable by elementarity.

So  $\kappa$  is Ramsey. But then  $\mathcal{M}_\omega \models \kappa$  is Ramsey and so  $\text{Ult}(\mathcal{M}_\omega, \mu_\omega) \models \kappa$  is Ramsey, since  $\mathcal{M}_\omega$  has the same subsets as the ultrapower by weak amenability of  $\mu_\omega$ .

## So what happens between $\omega$ and $\omega_1$ ?

### Proposition (N.)

Strategic  $(\omega+1)$ -Ramsey cardinals are Ramsey limits of Ramsey cardinals.

**Proof sketch.** Whenever  $\vec{\mathcal{M}}_\alpha * \vec{\mu}_\alpha$  is a play of  $\mathcal{G}_{\omega+1}^\theta(\kappa)$  then

$$(\mathcal{M}_{\omega+1}, \in, \mu_{\omega+1}) \models \mu_{\omega+1} \text{ is countably complete.}$$

Then  $\mu_\omega \subseteq \mu_{\omega+1}$  implies that, wlog,

$$\mathcal{M}_{\omega+1} \models \mu_\omega \text{ is countably complete and weakly amenable,}$$

making  $\mu_\omega$  countably complete and weakly amenable by elementarity.

So  $\kappa$  is Ramsey. But then  $\mathcal{M}_\omega \models \kappa$  is Ramsey and so  $\text{Ult}(\mathcal{M}_\omega, \mu_\omega) \models \kappa$  is Ramsey, since  $\mathcal{M}_\omega$  has the same subsets as the ultrapower by weak amenability of  $\mu_\omega$ .

Łoś and elementarity then makes  $\kappa$  a limit of Ramseys. □



Alright then. What about when we exceed  $\omega_1$ ?

### Proposition (N.)

Let  $\lambda$  be uncountable regular. Then  $\kappa$  is strategic  $\lambda$ -Ramsey iff

Alright then. What about when we exceed  $\omega_1$ ?


### Proposition (N.)

Let  $\lambda$  be uncountable regular. Then  $\kappa$  is strategic  $\lambda$ -Ramsey iff there's a  $<\lambda$ -closed forcing  $\mathbb{P}$  such that, in  $V^{\mathbb{P}}$ , there's a transitive  $N$  with  $\mathcal{P}(\kappa)^V = \mathcal{P}(\kappa)^N$  and an elementary embedding  $j : V \rightarrow N$  with  $\text{crit}(j) = \kappa$ .

Alright then. What about when we exceed  $\omega_1$ ?

### Proposition (N.)

Let  $\lambda$  be uncountable regular. Then  $\kappa$  is strategic  $\lambda$ -Ramsey iff there's a  $<\lambda$ -closed forcing  $\mathbb{P}$  such that, in  $V^{\mathbb{P}}$ , there's a transitive  $N$  with  $\mathcal{P}(\kappa)^V = \mathcal{P}(\kappa)^N$  and an elementary embedding  $j : V \rightarrow N$  with  $\text{crit}(j) = \kappa$ .



Alright then. What about when we exceed  $\omega_1$ ?

### Proposition (N.)

Let  $\lambda$  be uncountable regular. Then  $\kappa$  is strategic  $\lambda$ -Ramsey iff there's a  $<\lambda$ -closed forcing  $\mathbb{P}$  such that, in  $V^{\mathbb{P}}$ , there's a transitive  $N$  with  $\mathcal{P}(\kappa)^V = \mathcal{P}(\kappa)^N$  and an elementary embedding  $j : V \rightarrow N$  with  $\text{crit}(j) = \kappa$ .

Alright then. What about when we exceed  $\omega_1$ ?

### Proposition (N.)

Let  $\lambda$  be uncountable regular. Then  $\kappa$  is strategic  $\lambda$ -Ramsey iff there's a  $<\lambda$ -closed forcing  $\mathbb{P}$  such that, in  $V^{\mathbb{P}}$ , there's a transitive  $N$  with  $\mathcal{P}(\kappa)^V = \mathcal{P}(\kappa)^N$  and an elementary embedding  $j : V \rightarrow N$  with  $\text{crit}(j) = \kappa$ .

### Theorem (Schindler-N.)

The “ $\Rightarrow$ ” direction in the  $\lambda = \omega$  case above.

Alright then. What about when we exceed  $\omega_1$ ?

### Proposition (N.)

Let  $\lambda$  be uncountable regular. Then  $\kappa$  is strategic  $\lambda$ -Ramsey iff there's a  $<\lambda$ -closed forcing  $\mathbb{P}$  such that, in  $V^{\mathbb{P}}$ , there's a transitive  $N$  with  $\mathcal{P}(\kappa)^V = \mathcal{P}(\kappa)^N$  and an elementary embedding  $j : V \rightarrow N$  with  $\text{crit}(j) = \kappa$ .

### Theorem (Schindler-N.)

The “ $\Rightarrow$ ” direction in the  $\lambda = \omega$  case above.

The “ $\Leftarrow$ ” direction is open – getting wellfoundedness of  $\text{Ult}(\mathcal{M}_\omega, \mu_\omega)$  seems hard. We've shown it in the case where  $N \subseteq V$ .

# What's not known?

## Question 1

Are the strategic  $\alpha$ -Ramseys equivalent to some kind of “generic embedding property” when  $\alpha$  is countably infinite, as in the uncountable regular case?

# What's not known?

## Question 1

Are the strategic  $\alpha$ -Ramseys equivalent to some kind of “generic embedding property” when  $\alpha$  is countably infinite, as in the uncountable regular case?


## Question 2

Do the strategic  $\alpha$ -Ramseys form a strict hierarchy for  $\alpha$  countably infinite? More specifically, does

$$\text{ZFC} + \exists \text{strategic } (\alpha+1)\text{-Ramsey} \vdash \text{Con}(\exists \text{strategic } \alpha\text{-Ramsey})?$$



# An overview



Thank you for your attention

Slides and preprint available at  
<https://dsnielsen.com>