


# Matrix Transformation

CE40282-1: Linear Algebra  
Hamid R. Rabiee and Maryam Ramezani  
Sharif University of Technology





# Linear Transformation

- Matrix is a linear transformation: map one vector to another vector

$$A \in \mathbb{R}^{m \times n}, x \in \mathbb{R}^n, y \in \mathbb{R}^m : \quad y_{m \times 1} = A_{m \times n} x_{n \times 1}$$

$$A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$$


- Input-output



# Matrix in applications

- Feature matrix
- Signal matrix
- Correlation matrix

# Linear Transformation



Domain, codomain, and range of  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

# Linear Transformation

**EXAMPLE 1** Let  $A = \begin{bmatrix} 1 & -3 \\ 3 & 5 \\ -1 & 7 \end{bmatrix}$ ,  $\mathbf{u} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $\mathbf{b} = \begin{bmatrix} 3 \\ 2 \\ -5 \end{bmatrix}$ ,  $\mathbf{c} = \begin{bmatrix} 3 \\ 2 \\ 5 \end{bmatrix}$ , and define a transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  by  $T(\mathbf{x}) = A\mathbf{x}$ , so that

$$T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} 1 & -3 \\ 3 & 5 \\ -1 & 7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 - 3x_2 \\ 3x_1 + 5x_2 \\ -x_1 + 7x_2 \end{bmatrix}$$

- Find  $T(\mathbf{u})$ , the image of  $\mathbf{u}$  under the transformation  $T$ .
- Find an  $\mathbf{x}$  in  $\mathbb{R}^2$  whose image under  $T$  is  $\mathbf{b}$ .
- Is there more than one  $\mathbf{x}$  whose image under  $T$  is  $\mathbf{b}$ ?
- Determine if  $\mathbf{c}$  is in the range of the transformation  $T$ .

# Linear mapping

- $V, W$  are vector spaces over  $\mathbb{F}$ .

- A function  $T : V \rightarrow W$  is called **linear** if

$$T(u + v) = T(u) + T(v)$$

for all  $u, v \in V$ ,

$$T(av) = aT(v)$$

for all  $a \in \mathbb{F}$  and  $v \in V$ .

# Linear mapping

- Example: which are linear mapping?
  - **zero** map  $0 : V \rightarrow W$
  - **identity** map  $I : V \rightarrow V$
  - Let  $T : \mathcal{P}(\mathbb{F}) \rightarrow \mathcal{P}(\mathbb{F})$  be the **differentiation** map defined as  $Tp(z) = p'(z)$ .
  - Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be the map given by  $T(x, y) = (x - 2y, 3x + y)$
  - $f(x) = e^x$
  - $f : \mathbb{F} \rightarrow \mathbb{F}$  given by  $f(x) = x - 1$

# Linear mapping

## ■ Theorem


*Let  $(v_1, \dots, v_n)$  be a basis of  $V$  and  $(w_1, \dots, w_n)$  an arbitrary list of vectors in  $W$ . Then there exists a unique linear map*

$$T : V \rightarrow W \quad \text{such that } T(v_i) = w_i.$$



# Projection



## ■ Example:



If  $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ , then the transformation  $\mathbf{x} \mapsto A\mathbf{x}$  projects points in  $\mathbb{R}^3$  onto the  $x_1x_2$ -plane because

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \mapsto \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \\ 0 \end{bmatrix}$$


# Projection

Transformation	Image of the Unit Square	Standard Matrix
Projection onto the $x_1$ -axis		$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$
Projection onto the $x_2$ -axis		$\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$

# Projection



The **projection** of a vector  $y \in \mathbb{R}^m$  onto the span of  $\{x_1, \dots, x_n\}$  is the vector  $v \in \text{span}(\{x_1, \dots, x_n\})$ , such that  $v$  is as close as possible to  $y$ , as measured by the Euclidean norm  $\|v - y\|_2$ .

$$\text{Proj}(y; \{x_1, \dots, x_n\}) = \operatorname{argmin}_{v \in \text{span}(\{x_1, \dots, x_n\})} \|y - v\|_2.$$





# Rotation

- $R = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$






# Reflection

- $R = \begin{bmatrix} \cos(2\theta) & \sin(2\theta) \\ \sin(2\theta) & -\cos(2\theta) \end{bmatrix}$




# Reflection

Transformation	Image of the Unit Square	Standard Matrix
Reflection through the $x_1$ -axis		$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$
Reflection through the $x_2$ -axis		$\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$
Reflection through the line $x_2 = x_1$		$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$


# Reflection

Reflection through  
the line  $x_2 = -x_1$



$$\begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$



Reflection through  
the origin



$$\begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

# Uniform Scaling



- $S = sI = \begin{bmatrix} s & 0 \\ 0 & s \end{bmatrix}$





# Non-uniform Scaling

$$\blacksquare S = \begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix}$$



# Shearing

## ■ Example

Let  $A = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$ . The transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

A typical shear matrix is of the form

$$S = \begin{pmatrix} 1 & 0 & 0 & \lambda & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$



sheep



sheared sheep


# Shearing

A shear parallel to the  $x$  axis results in  $x' = x + \lambda y$  and  $y' = y$ . In matrix form:

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

Similarly, a shear parallel to the  $y$  axis has  $x' = x$  and  $y' = y + \lambda x$ . In matrix form:

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$



# Difference Matrix

■

$$D_{(n-1) \times n} = \begin{bmatrix} -1 & 1 & 0 & 0 & \cdots & 0 \\ 0 & -1 & 1 & 0 & \cdots & 0 \\ \vdots & & \ddots & \ddots & & \vdots \\ 0 & 0 & \cdots & -1 & 1 & 0 \\ 0 & 0 & \cdots & 0 & -1 & 1 \end{bmatrix}$$

$$D : \mathbb{R}^n \longrightarrow \mathbb{R}^{n-1} \quad \Rightarrow \quad D \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} x_2 - x_1 \\ x_3 - x_2 \\ \vdots \\ x_n - x_{n-1} \end{bmatrix}$$

■ Example

$$\begin{bmatrix} -1 & 1 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ -1 \\ 3 \\ 2 \\ 5 \end{bmatrix} = \begin{bmatrix} -1 - 0 \\ 3 - (-1) \\ 2 - 3 \\ 5 - 2 \end{bmatrix} = \begin{bmatrix} -1 \\ 4 \\ -1 \\ 3 \end{bmatrix}$$

# Selectors

- an  $m \times n$  selector matrix: each row is a unit vector (transposed)

$$A = \begin{bmatrix} e_{k_1}^T \\ \vdots \\ e_{k_m}^T \end{bmatrix}$$

multiplying by  $A$  selects entries of  $x$ :

$$Ax = (x_{k_1}, x_{k_2}, \dots, x_{k_m})$$

- $A : \mathbb{R}^n \longrightarrow \mathbb{R}^m \quad \Rightarrow \quad A \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} x_{k_1} \\ x_{k_2} \\ \vdots \\ x_{k_m} \end{bmatrix}$

# Selectors

■ Example 
$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \\ 0 \\ -3 \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \end{bmatrix}$$

- Selecting first and last elements of vector:
- Reversing the elements of vector:

# Slicing

- Keeping  $m$  elements from  $r$  to  $s$  ( $m=s-r+1$ )

$$\begin{bmatrix} 0_{m \times (r-1)} & I_{m \times m} & 0_{m \times (n-s)} \end{bmatrix}$$

- Example: Slicing two first and one last elements:

$$\begin{bmatrix} -1 \\ 2 \\ 0 \\ -3 \\ 5 \end{bmatrix} = \begin{bmatrix} 0 \\ -3 \end{bmatrix}$$

# Down Sampling

- Down sampling with k: selecting one sample in every k samples
- Example: k=2?

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ \vdots \end{bmatrix} = \begin{bmatrix} x_1 \\ x_3 \\ x_5 \\ \vdots \end{bmatrix}$$




# Applications

- Rotation matrix

$$\begin{aligned} \text{(i)} \quad \sin 2A &= 2 \sin A \cos A \\ \text{(ii)} \quad \cos 2A &= \cos^2 A - \sin^2 A \end{aligned}$$

$$R = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \Rightarrow R^n = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}^n = \begin{bmatrix} \cos(n\theta) & -\sin(n\theta) \\ \sin(n\theta) & \cos(n\theta) \end{bmatrix}$$

- Adjacency matrix



$$A = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 \end{bmatrix}$$

$$A^3 = \begin{bmatrix} 1 & 0 & 1 & 0 & 0 & 1 \\ 1 & 2 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \\ 2 & 1 & 0 & 0 & 0 & 1 \end{bmatrix}$$

# Multiple Transformation

$$\blacksquare \quad x_{n \times 1} \xrightarrow{A_{m \times n}} y_{m \times 1} \xrightarrow{B_{p \times m}} z_{p \times 1} \implies \begin{cases} y = Ax \\ z = By \end{cases} \implies z = B(Ax) = BAx$$

## ■ Example

### ■ Difference Matrix

$$x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} \xrightarrow[4 \times 5]{D} y = \begin{bmatrix} x_2 - x_1 \\ x_3 - x_2 \\ x_4 - x_3 \\ x_5 - x_4 \end{bmatrix} \xrightarrow[3 \times 4]{D} z = \begin{bmatrix} x_3 - x_2 - (x_2 - x_1) \\ x_4 - x_3 - (x_3 - x_2) \\ x_5 - x_4 - (x_4 - x_3) \end{bmatrix} = \begin{bmatrix} x_3 - 2x_2 + x_1 \\ x_4 - 2x_3 + x_2 \\ x_5 - 2x_4 + x_3 \end{bmatrix}$$

$$x \longrightarrow z \begin{bmatrix} 1 & -2 & 1 & 0 & 0 \\ 0 & 1 & -2 & 1 & 0 \\ 0 & 0 & 1 & -2 & 1 \end{bmatrix}_{3 \times 5}$$

$$x \longrightarrow y \longrightarrow z$$


$$\begin{bmatrix} 1 & -1 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & -1 \end{bmatrix}_{3 \times 4} \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix}_{4 \times 5} = \begin{bmatrix} 1 & -2 & 1 & 0 & 0 \\ 0 & 1 & -2 & 1 & 0 \\ 0 & 0 & 1 & -2 & 1 \end{bmatrix}$$

# Multiple Transformation

- $x_{n \times 1} \xrightarrow{A_{m \times n}} y_{m \times 1} \xrightarrow{B_{p \times m}} z_{p \times 1} \implies \begin{cases} y = Ax \\ z = By \end{cases} \implies z = B(Ax) = BAx$

- Example

- Rotation



$$R_\theta = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$


$$x \rightarrow z \quad z = R_{\delta+\theta}x \quad \begin{bmatrix} \cos(\delta + \theta) & -\sin(\delta + \theta) \\ \sin(\delta + \theta) & \cos(\delta + \theta) \end{bmatrix}$$

$$x \rightarrow y \rightarrow z \quad \begin{cases} y = R_\theta x \\ z = R_\delta y \end{cases} \implies z = R_\delta R_\theta x \quad \begin{bmatrix} \cos \delta & -\sin \delta \\ \sin \delta & \cos \delta \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

$$= \begin{bmatrix} \cos \delta \cos \theta - \sin \delta \sin \theta & -\cos \delta \sin \theta - \sin \delta \cos \theta \\ \sin \delta \cos \theta + \cos \delta \sin \theta & -\sin \delta \sin \theta + \cos \delta \cos \theta \end{bmatrix}$$

$$= \begin{bmatrix} \cos(\delta + \theta) & -\sin(\delta + \theta) \\ \sin(\delta + \theta) & \cos(\delta + \theta) \end{bmatrix}$$

# Invertible Linear Transformations



## ■ Definition:

A linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is said to be **invertible** if there exists a function  $S : \mathbb{R}^n \rightarrow \mathbb{R}^n$  such that

$$S(T(\mathbf{x})) = \mathbf{x} \quad \text{for all } \mathbf{x} \text{ in } \mathbb{R}^n$$

$$T(S(\mathbf{x})) = \mathbf{x} \quad \text{for all } \mathbf{x} \text{ in } \mathbb{R}^n$$

# Invertible Linear Transformations

- Theorem:

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation and let  $A$  be the standard matrix for  $T$ . Then  $T$  is invertible if and only if  $A$  is an invertible matrix. In that case, the linear transformation  $S$  given by  $S(\mathbf{x}) = A^{-1}\mathbf{x}$  is the unique function satisfying

$$S(T(\mathbf{x})) = \mathbf{x} \quad \text{for all } \mathbf{x} \text{ in } \mathbb{R}^n$$


$$T(S(\mathbf{x})) = \mathbf{x} \quad \text{for all } \mathbf{x} \text{ in } \mathbb{R}^n$$

- Proof in HW3!


- Example:

# Mapping

A mapping  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is said to be **onto**  $\mathbb{R}^m$  if each  $\mathbf{b}$  in  $\mathbb{R}^m$  is the image of *at least one*  $\mathbf{x}$  in  $\mathbb{R}^n$ .





$T$  is not onto  $\mathbb{R}^m$



$T$  is onto  $\mathbb{R}^m$

A mapping  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is said to be **one-to-one** if each  $\mathbf{b}$  in  $\mathbb{R}^m$  is the image of *at most one*  $\mathbf{x}$  in  $\mathbb{R}^n$ .



Let  $T$  be the linear transformation whose standard matrix is

$$A = \begin{bmatrix} 1 & -4 & 8 & 1 \\ 0 & 2 & -1 & 3 \\ 0 & 0 & 0 & 5 \end{bmatrix}$$

Does  $T$  map  $\mathbb{R}^4$  onto  $\mathbb{R}^3$ ? Is  $T$  a one-to-one mapping?

# One-to-One Linear Transformation

## THEOREM

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then  $T$  is one-to-one if and only if the equation  $T(\mathbf{x}) = \mathbf{0}$  has only the trivial solution.




# One-to-One Linear Transformation

- Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation, and let  $A$  be the standard matrix for  $T$ . Then:
  - a.  $T$  maps  $\mathbb{R}^n$  onto  $\mathbb{R}^m$  if and only if the columns of  $A$  span  $\mathbb{R}^m$ ;
  - b.  $T$  is one-to-one if and only if the columns of  $A$  are linearly independent.

## ■ Example

Let  $T(x_1, x_2) = (3x_1 + x_2, 5x_1 + 7x_2, x_1 + 3x_2)$ . Show that  $T$  is a one-to-one linear transformation. Does  $T$  map  $\mathbb{R}^2$  onto  $\mathbb{R}^3$ ?



# Machine Learning Application

- The central problem in machine learning and deep learning is to meaningfully transform data: in other words, to learn useful representations of the input data at hand — representations that get us closer to the expected output.