Numerical Methods

Sharif University of Technology
Department of Computer Engineering
Fatemeh Baharifard
Mail: fateme.baharifard@gmail.com

©2021

©Sharif University Of Technology

Course Introduction

Webpage TAs Resources Grade

Join the Course Page on Quera.

Everything will be posted on Quera page.

©Sharif University Of Technology

Course Introduction

Webpage **TAs** Resources Grade

Kahbod Aeini Head
Mohammad Reza Daviran
Reza Abdollahzadeh
Mahdi Gheidi
Amir Mohammad Isazadeh
Masih Eskandar
Tarlan Bahadori
Neda Taghizadeh
Haniyeh Ehsani
Rosta Roghani

For any enquiries, please contact Mr. Aeini via
mailto: kahbodaeini@gmail.com

©Sharif University Of Technology

Course Introduction

Webpage TAs **Resources** Grade

- APPLIED NUMERICAL METHODS WITH MATLAB® for Engineers and Scientists -Steven C. Chapra -Third Edition 2012
- NUMERICAL METHODS IN ENGINEERING WITH MATLAB

  Jaan Kiusalaas, 3rd Edition

  Pennsylvania State University

  Cambridge University Press 2016
- محاسبات عددی، مسعود نیکوکار



©Sharif University Of Technology

Course Introduction

Webpage TAs Resources **Grade**

Grade		
Midterm	4	Ch(1-3) – 7 Azar
Final	7	Ch(1-6) – 25 Dei
Six HWs	6	Theoretical and Practical
Project	3	TBA
Random Quizzes	1	Extra points

©Sharif University Of Technology

Why Numerical Methods?

Some problems cannot be solved analytically or are too long and tedious to calculate.

$$\int_0^1 e^{-x^2}\,\mathrm{dx}$$

$$\int_0^1 \frac{1}{1+x^3}\,\mathrm{dx}$$

©Sharif University Of Technology

Numerical Methods

NM Methods:

- Iterative
- Direct

©Sharif University Of Technology

Contents

- Six chapters:
  - Errors
  - Numerical methods for solving nonlinear equations
  - Interpolation, extrapolation and curve fitting
  - Numerical Integration and differentiation
  - Ordinary differential equations
  - System of linear equations

©Sharif University Of Technology

Errors



©Sharif University Of Technology

Roots of Nonlinear Equations



Roots: Solve for x so that $f(x) = 0$

Optimization: Solve for x so that $f'(x) = 0$

©Sharif University Of Technology

Interpolation, Extrapolation, Curve Fitting





©Sharif University Of Technology

Integration and Differentiation



Integration: Find the area under the curve

Differentiation: Find the slope of the curve

©Sharif University Of Technology

Ordinary Differential Equations



Given

$$\frac{dy}{dt} \approx \frac{\Delta y}{\Delta t} = f(t, y)$$

solve for y as a function of t

$$y_{i+1} = y_i + f(t_i, y_i)\Delta t$$

©Sharif University Of Technology

System of Linear equations

Given the a's and the b's, solve for the x's

$$a_{11}x_1 + a_{12}x_2 = b_1$$

$$a_{21}x_1 + a_{22}x_2 = b_2$$



©Sharif University Of Technology

Chapter 1

Errors

©Sharif University Of Technology

Source of Error

Error Representation

Floating Point Representation

Types of Error

Error Propagation and Process Graph

©Sharif University Of Technology

Introduction

- What is error?
- Where does it come from?
- What types does it have?
- How can we minimize it?

©Sharif University Of Technology

Precision & Accuracy

Accuracy and Precision:

Accuracy refers to how closely a computed or measured value agrees with the true value.

Precision refers to how closely individual computed or measured values agree with each other.

Inaccuracy (also known as bias) is the systematic deviation from the truth.

Imprecision (uncertainty) refers to the magnitude of the scatter.

©Sharif University Of Technology

Precision & Accuracy



©Sharif University Of Technology

Examples of Errors

$$\sqrt{5} \stackrel{!}{=} 2\ldots$$

©Sharif University Of Technology

Examples of Errors

$$\sqrt{5} \overset{!}{=} 2.23.$$

©Sharif University Of Technology

Examples of Errors

$$\sqrt{5} \stackrel{!}{=} 2.2360679\ldots$$

©Sharif University Of Technology

Examples of Errors

Π = 3.14159265…

e = 2.71828182…

$^1/_3$ = 0.3333333…

Because of the limitation of showing numbers, we have inherent error!

©Sharif University Of Technology

Examples of Errors



In Euclidean plane geometry, π is defined as the ratio of a circle's circumference (C) to its diameter (d).

$$\pi = \frac{C}{d}$$

As the number of sides of a polygon increases, its area approximates the area of a circle more accurately, showing that the value of π can be estimated with regular polygons.

$$\pi = 3.14159265358979323846264338327950 28 \ldots$$

©Sharif University Of Technology

Examples of Errors



In Euclidean plane geometry, π is defined as the ratio of a circle's circumference (C) to its diameter (d).

$$\pi = \frac{C}{d}$$

As the number of sides of a polygon increases, its area approximates the area of a circle more accurately, showing that the value of π can be estimated with regular polygons.

$$\pi = 3.14159265358979323846264338327950 28 \ldots$$

©Sharif University Of Technology

Examples of Errors





Actual Size	$17.2\ cm$	$82.49\ m$
Measured Size	$18.2\ cm$	$82.5\ m$

©Sharif University Of Technology

Sources of Error

- Measurement

  Measurement contains error.
- Mathematical Models

  Some parameters are ignored in mathematical modeling.
- Truncation Errors
- Roundoff Errors
- Operation Errors (propagation)

©Sharif University Of Technology

Error Representation

Absolute Relative Example

True Value Approximate Absolute Error

$$x \dashrightarrow \bar{x} \dashrightarrow e(\bar{x}) = |x - \bar{x}|$$

The error does not have sign (It is always positive)!

©Sharif University Of Technology

An Example





Actual Size	$17.2\ cm$	$82.49\ m$
Measured Size	$18.2\ cm$	$82.5\ m$
Absolute Error	$1\ cm$	$1\ cm$

©Sharif University Of Technology

Error Representation

Absolute **Relative** Example

True Value Approximate Relative Error

$$x \dashrightarrow \bar{x} \dashrightarrow \delta(\bar{x}) = \left| \frac{e(\bar{x})}{x} \right|$$

The error does not have sign (It is always positive)!

©Sharif University Of Technology

An Example





Actual Size	$17.2\ cm$	$82.49\ m$
Measured Size	$18.2\ cm$	$82.5\ m$
Absolute Error	$1\ cm$	$1\ cm$
Relative Error	0.05	0.00012

©Sharif University Of Technology

Error Representation

Absolute Relative Example

$$e(\bar{x}) \leq e_{\bar{x}}$$

Example: $x = \sqrt{2}$

$\bar{x} = 1.41$

$e(\bar{x}) = |x - \bar{x}| = |\sqrt{2} - 1.41| < 0.005$

$$\boxed{|x - \bar{x}| \leq e_{\bar{x}} \longleftrightarrow x = \bar{x} \pm e_{\bar{x}}}$$

©Sharif University Of Technology

Error Representation

Absolute **Relative** Example

$$\delta(\bar{x}) = \frac{e(\bar{x})}{|x|} \leq \frac{e_{\bar{x}}}{|x|} \cong \frac{e_{\bar{x}}}{|\bar{x}|}$$

©Sharif University Of Technology

Representation of Floating-Point Numbers

$$23.1 = 2.31 \times 10^1 = 0.231 \times 10^2 = 0.0231 \times 10^3$$

$$231 \times 10^{-1} = 2310 \times 10^{-2} = 23100 \times 10^{-3}$$

Which form is normalized?

©Sharif University Of Technology

Representation of Floating-Point Numbers

$$23.1 = 2.31 \times 10^1 = 0.231 \times 10^2 = 0.0231 \times 10^3$$

$$231 \times 10^{-1} = 2310 \times 10^{-2} = 23100 \times 10^{-3}$$

Which form is normalized?

©Sharif University Of Technology

Normalized Representation

$$z = \sigma \times (a_0.a_1a_2a_3\ldots)_\beta \times \beta^e = \sigma \times m \times \beta^e$$

σ is the sign (+ or -).

β is the base, e is the exponent.

binary : $\beta = 2$

decimal : $\beta = 10$

$\boldsymbol{m}$ is mantissa $(significant)$:

$1 \leq \boldsymbol{m} < \beta \quad , \quad (a_0 \neq 0 \ \ and \ \ 0 \leq a_i \leq \beta - 1)$

binary: $1 \leq m < 2$

decimal: $1 \leq m < 10$

©Sharif University Of Technology

Normalized Representation

Example: z = 0.005678

decimal normalized : z = 5.678* 10^{-3}

$x = 3.5$ *(2 significant digits)* ➡ $3.45 \leq x < 3.55$

©Sharif University Of Technology

IEEE-754 Double-Precision Format

The manner in which a floating-point number is stored in an 8-byte word in IEEE double precision format:



©Sharif University Of Technology

IEEE-754 Double-Precision Format

Since binary numbers consist exclusively of 0s and 1s, a bonus occurs when they are normalized:
The bit to the left of the binary point will always be one and does not have to be stored.

$$\pm(1+f)\times 2^{e}$$

f **= the *mantissa* (i.e., the fractional part of the significand).**

Example:
Normalized the binary number 1101.1:

1.1011×2^{-3} **or** $(1+0.1011)\times 2^{-3}$

only have to store the four fractional bits instance of five significant bits.

©Sharif University Of Technology

IEEE-754 Double-Precision Format

Range. In a fashion similar to the way in which integers are stored, the 11 bits used for the exponent translates into a range from -1022 to 1023. The largest positive number can be represented in binary as

$$\text{Largest value} = +1.1111\ldots1111 \times 2^{+1023}$$

where the 52 bits in the mantissa are all 1. Since the significand is approximately 2 (it is actually $2 - 2^{-52}$), the largest value is therefore $2^{1024} = 1.7977 \times 10^{308}$. In a similar fashion, the smallest positive number can be represented as

$$\text{Smallest value} = +1.0000\ldots0000 \times 2^{-1022}$$

This value can be translated into a base-10 value of $2^{-1022} = 2.2251 \times 10^{-308}$.

©Sharif University Of Technology

IEEE-754 Double-Precision Format

Precision. The 52 bits used for the mantissa correspond to about 15 to 16 base-10 digits. Thus, π would be expressed as

```
>> format long
>> pi

ans =
   3.14159265358979
```

Note that the machine epsilon is $2^{-52} = 2.2204 \times 10^{-16}$.

©Sharif University Of Technology

Types of Errors

Inherent Round Off Truncation

Length = 23.47

$$23.465 < \text{length} < 23.475$$

©Sharif University Of Technology

Types of Errors

Inherent **Round Off** Truncation

Decimal : $\frac{1}{3} = 0.\overline{3}$

Binary : $(0.1)_{10} = (0.0\overline{0011})_2$

©Sharif University Of Technology

Types of Errors

Inherent **Round Off** Truncation

	Chopping	Symmetric
0.00065	0.0006	0.0007

Rounded off to 4 digits (4D).

Example (symmetric): 1.23456

1.23 (2D)

1.235 (3D)

1.2346 (4D)

©Sharif University Of Technology

Maximum Round Off Error

Absolute:

Chopping	$\lvert e_x \rvert < 10^{-t}$
Symmetric	$\lvert e_x \rvert \leq \frac{1}{2} \times 10^{-t}$

Rounded off to t digits.

©Sharif University Of Technology

Types of Errors

Inherent Round Off **Truncation**

$$\sin x = x - \frac{x^{۳}}{۳!} + \frac{x^{۵}}{۵!} - \cdots + (-۱)^n \frac{x^{۲n+۱}}{(۲n+۱)!} \pm \cdots$$

$$e^x = ۱ + \frac{x}{۱!} + \frac{x^{۲}}{۲!} + \cdots + \frac{x^{n}}{n!} + E_n(x)$$

$$f(x_{i+1}) = f(x_i) + f'(x_i)h + \frac{f''(x_i)}{2!}h^2 + \frac{f^{(3)}(x_i)}{3!}h^3 + \cdots + \frac{f^{(n)}(x_i)}{n!}h^n + E_n$$

©Sharif University Of Technology

Types of Errors

Inherent Round Off **Truncation**

مثال ۱۴. مقدار تقریبی تابع $\sin x$ را به ازای $x = \frac{\pi}{۷}$ و با خطای کمتر از $۱۰^{-۳}$ حساب کنید.

حل : داریم

$$\sin x = x - \frac{x^{۳}}{۳!} + \frac{x^{۵}}{۵!} - \cdots + (-۱)^{n} \frac{x^{۲n+۱}}{(۲n+۱)!} \pm \cdots$$

در اینجا قرار می‌دهیم $|E_n(x)| = \frac{x^{۲n+۱}}{(۲n+۱)!}$

$$x = \frac{\pi}{۷} = \pi \frac{۱}{۷} = ۳/۱۴۱۶ \times ۰/۱۴۲۹ = ۰/۴۴۸۹$$

©Sharif University Of Technology

Types of Errors

Inherent Round Off **Truncation**

بنابراین بایستی n را طوری تعیین کنیم که

$$\frac{(۰٫۴۴۸۹)^{۲n+۱}}{(۲n+۱)!} \leq \frac{۱}{۲} \times ۱۰^{-۳} = ۵ \times ۱۰^{-۴}$$

برای $n \geq ۲$ نامساوی فوق برقرار می‌باشد، در نتیجه

$$\sin\frac{\pi}{۷} \simeq ۰٫۴۴۸۹ - \frac{(۰٫۴۴۸۹)^{۳}}{۳!} + \frac{(۰٫۴۴۸۹)^{۵}}{۵!}$$

$$= ۰٫۴۴۸۹ - ۰٫۰۱۵۱ + ۰٫۰۰۰۲$$

$$= ۰٫۴۳۴۰ \quad (۴D)$$

$$\sin\frac{\pi}{۷} \simeq ۰٫۴۳۴(۳D)$$

©Sharif University Of Technology

Error Propagation

When does error propagation occur?

1) When we want to substitute parameters of formulas with non-exact values.

$$s = \pi r^2$$

2) When we have two algebraically equivalent equations and like to discover which one is better for implementation.

$$a^2 - b^2$$

$$(a - b)(a + b)$$

©Sharif University Of Technology

Error Propagation

Absolute Error Relative Error

Addition (+):

$\bar{x}$ and $\bar{y}$ are approximations of x and y $(\bar{x}, \bar{y} > 0)$

$|x - \bar{x}| \leq e_{\bar{x}}$ and $|y - \bar{y}| \leq e_{\bar{y}}$

$$\bar{x} - e_{\bar{x}} \leq x \leq \bar{x} + e_{\bar{x}}$$

$$\bar{y} - e_{\bar{y}} \leq y \leq \bar{y} + e_{\bar{y}}$$

$$\bar{x} + \bar{y} - (e_{\bar{x}} + e_{\bar{y}}) \leq x + y \leq \bar{x} + \bar{y} + (e_{\bar{x}} + e_{\bar{y}})$$

$$|(x + y) - (\bar{x} + \bar{y})| \leq (e_{\bar{x}} + e_{\bar{y}})$$

$$e_{\bar{x}+\bar{y}} \leq e_{\bar{x}} + e_{\bar{y}}$$

©Sharif University Of Technology

Error Propagation

Absolute Error Relative Error

Subtraction (-):

$\bar{x}$ and $\bar{y}$ are approximations of x and y $(\bar{x}, \bar{y} > 0)$

$|x - \bar{x}| \leq e_{\bar{x}}$ and $|y - \bar{y}| \leq e_{\bar{y}}$

$$\bar{x} - e_{\bar{x}} \leq x \leq \bar{x} + e_{\bar{x}}$$

$$\bar{y} - e_{\bar{y}} \leq y \leq \bar{y} + e_{\bar{y}}$$

$$-\bar{y} - e_{\bar{y}} \leq -y \leq -\bar{y} + e_{\bar{y}}$$

$$\bar{x} - \bar{y} - (e_{\bar{x}} + e_{\bar{y}}) \leq x - y \leq \bar{x} - \bar{y} + (e_{\bar{x}} + e_{\bar{y}})$$

$$e_{\bar{x}-\bar{y}} \leq e_{\bar{x}} + e_{\bar{y}}$$

©Sharif University Of Technology

Error Propagation

Absolute Error

Relative Error

$$e_{\bar{x}+\bar{y}} \leq e_{\bar{x}} + e_{\bar{y}}$$

$$e_{\bar{x}-\bar{y}} \leq e_{\bar{x}} + e_{\bar{y}}$$

©Sharif University Of Technology

Error Propagation

Absolute Error

Relative Error

مثال ۸. هرگاه اعداد $\sqrt{۱۷}$ و $\sqrt{۵}$ را تا سه رقم اعشار گرد کنیم، مطلوبست محاسبه $\sqrt{۱۷} \pm \sqrt{۵}$ و محاسبه حداکثر خطای حاصل جمع و تفاضل.

حل : داریم : $\sqrt{۱۷} = ۴/۱۲۳ + e_۱, \qquad \sqrt{۵} = ۲/۲۳۶ + e_۲$

منظور از $e_۱$ و $e_۲$ خطای مرتکب شده در نمایش $\sqrt{۱۷}$ و $\sqrt{۵}$ می‌باشد. چون اعداد تا سه رقم اعشار گرد شده‌اند، پس

$$e_۱ \leq \frac{۱}{۲} \times ۱۰^{-۳}, \qquad e_۲ \leq \frac{۱}{۲} \times ۱۰^{-۳}$$

داریم : $\sqrt{۱۷} + \sqrt{۵} = (۴/۱۲۳ + ۲/۲۳۶) + e_۳ = ۶/۳۵۹ + e_۳$

و چون $e_۳ \leq e_۱ + e_۲$ لذا $e_۳ \leq ۱۰^{-۳}$. در نتیجه

$$۶/۳۵۹ - ۱۰^{-۳} \leq \sqrt{۱۷} + \sqrt{۵} \leq ۶/۳۵۹ + ۱۰^{-۳}$$

همچنین $\sqrt{۱۷} - \sqrt{۵} = ۱/۸۸۷ + e_۴$ که در اینجا نیز

$$e_۴ \leq e_۱ + e_۲ \leq ۱۰^{-۳}$$

بنابراین $۱/۸۸۷ - ۱۰^{-۳} \leq \sqrt{۱۷} - \sqrt{۵} \leq ۱/۸۸۷ + ۱۰^{-۳}$

©Sharif University Of Technology

Error Propagation

Absolute Error

Relative Error

Addition (+):

$\bar{x}$ and $\bar{y}$ are approximations of x and y $(\bar{x}, \bar{y} > 0)$

$$\delta_{\bar{x}} \leq \frac{e_{\bar{x}}}{\bar{x}} \qquad and \qquad \delta_{\bar{y}} \leq \frac{e_{\bar{y}}}{\bar{y}}$$

$$\delta_{\bar{x}+\bar{y}} \leq \frac{e_{\bar{x}+\bar{y}}}{\bar{x}+\bar{y}} \leq \frac{e_{\bar{x}} + e_{\bar{y}}}{\bar{x}+\bar{y}} = \frac{\bar{x}}{\bar{x}+\bar{y}} * \frac{e_{\bar{x}}}{\bar{x}} + \frac{\bar{y}}{\bar{x}+\bar{y}} * \frac{e_{\bar{y}}}{\bar{y}} = \frac{\bar{x}}{\bar{x}+\bar{y}}\delta_{\bar{x}} + \frac{\bar{y}}{\bar{x}+\bar{y}}\delta_{\bar{y}}$$

$$\delta_{\bar{x}+\bar{y}} \leq \frac{\bar{x}}{\bar{x}+\bar{y}}\delta_{\bar{x}} + \frac{\bar{y}}{\bar{x}+\bar{y}}\delta_{\bar{y}}$$

©Sharif University Of Technology

Error Propagation

Absolute Error **Relative Error**

Subtraction (-):

$\bar{x}$ and $\bar{y}$ are approximations of x and y $(\bar{x}, \bar{y} > 0)$

$$\delta_{\bar{x}} \leq \frac{e_{\bar{x}}}{\bar{x}} \qquad and \qquad \delta_{\bar{y}} \leq \frac{e_{\bar{y}}}{\bar{y}}$$

$$\delta_{\bar{x}-\bar{y}} \leq \frac{e_{\bar{x}-\bar{y}}}{\bar{x}-\bar{y}} \leq \frac{e_{\bar{x}}+e_{\bar{y}}}{\bar{x}-\bar{y}} = \frac{\bar{x}}{\bar{x}-\bar{y}} * \frac{e_{\bar{x}}}{\bar{x}} + \frac{\bar{y}}{\bar{x}-\bar{y}} * \frac{e_{\bar{y}}}{\bar{y}} = \frac{\bar{x}}{\bar{x}-\bar{y}}\delta_{\bar{x}} + \frac{\bar{y}}{\bar{x}-\bar{y}}\delta_{\bar{y}}$$

$$\delta_{\bar{x}-\bar{y}} \leq \frac{\bar{x}}{\bar{x}-\bar{y}}\delta_{\bar{x}} + \frac{\bar{y}}{\bar{x}-\bar{y}}\delta_{\bar{y}}$$

©Sharif University Of Technology

Error Propagation

Absolute Error

Relative Error

$$\delta_{\bar{x}+\bar{y}} \leq \frac{\bar{x}}{\bar{x}+\bar{y}}\delta_{\bar{x}} + \frac{\bar{y}}{\bar{x}+\bar{y}}\delta_{\bar{y}} \qquad \bar{x}, \bar{y} > 0$$

$$\delta_{\bar{x}-\bar{y}} \leq \frac{\bar{x}}{\bar{x}-\bar{y}}\delta_{\bar{x}} + \frac{\bar{y}}{\bar{x}-\bar{y}}\delta_{\bar{y}} \qquad \bar{x} > \bar{y} > 0$$

Nearly identical amounts for $\bar{x}$ and $\bar{y}$ increase error propagation.

©Sharif University Of Technology

Error Propagation

Absolute Error Relative Error

Multiplication (*):

$\bar{x}$ and $\bar{y}$ are approximations of x and y $(\bar{x}, \bar{y} > 0)$

$|x - \bar{x}| \leq e_{\bar{x}}$ and $|y - \bar{y}| \leq e_{\bar{y}}$

$$\bar{x} - e_{\bar{x}} \leq x \leq \bar{x} + e_{\bar{x}}$$

$$\bar{y} - e_{\bar{y}} \leq y \leq \bar{y} + e_{\bar{y}}$$

$$\bar{x}\bar{y} - (\bar{y}e_{\bar{x}} + \bar{x}e_{\bar{y}}) + e_{\bar{x}}e_{\bar{y}} \leq xy \leq \bar{x}\bar{y} + (ye_{\bar{x}} + xe_{\bar{y}}) + e_{\bar{x}}e_{\bar{y}}$$

$$e_{\bar{x}\bar{y}} \leq \bar{y}e_{\bar{x}} + \bar{x}e_{\bar{y}}$$

©Sharif University Of Technology

Error Propagation

Absolute Error Relative Error

Division (/):

$\bar{x}$ and $\bar{y}$ are approximations of x and y $(\bar{x}, \bar{y} > 0)$

$|x-\bar{x}| \le e_{\bar{x}}$ and $|y-\bar{y}| \le e_{\bar{y}}$

$$\bar{x} - e_{\bar{x}} \le x \le \bar{x} + e_{\bar{x}}$$

$$\bar{y} - e_{\bar{y}} \le y \le \bar{y} + e_{\bar{y}}$$

$$\frac{\bar{x} - e_{\bar{x}}}{\bar{y} + e_{\bar{y}}} \le \frac{x}{y} \le \frac{\bar{x} + e_{\bar{x}}}{\bar{y} - e_{\bar{y}}}$$

$$\frac{\bar{x} - e_{\bar{x}}}{\bar{y} + e_{\bar{y}}} * \frac{\bar{y} - e_{\bar{y}}}{\bar{y} - e_{\bar{y}}} = \frac{\bar{x}\bar{y} - \bar{y}e_{\bar{x}} - \bar{x}e_{\bar{y}} + e_{\bar{x}}e_{\bar{y}}}{\bar{y}^2 - {e_{\bar{y}}}^2} = \frac{\bar{x}}{\bar{y}} - \frac{\bar{y}e_{\bar{x}} + \bar{x}e_{\bar{y}}}{\bar{y}^2}$$

$$\frac{\bar{x} + e_{\bar{x}}}{\bar{y} - e_{\bar{y}}} * \frac{\bar{y} + e_{\bar{y}}}{\bar{y} + e_{\bar{y}}} = \frac{\bar{x}\bar{y} + \bar{y}e_{\bar{x}} + \bar{x}e_{\bar{y}} + e_{\bar{x}}e_{\bar{y}}}{\bar{y}^2 - {e_{\bar{y}}}^2} = \frac{\bar{x}}{\bar{y}} + \frac{\bar{y}e_{\bar{x}} + \bar{x}e_{\bar{y}}}{\bar{y}^2}$$

$$e_{\frac{\bar{x}}{\bar{y}}} \le \frac{\bar{y}e_{\bar{x}} + \bar{x}e_{\bar{y}}}{\bar{y}^2}$$

©Sharif University Of Technology

Error Propagation

Absolute Error

Relative Error

$$e_{\bar{x}\times\bar{y}} \leq e_{\bar{x}} \times |\bar{y}| + e_{\bar{y}} \times |\bar{x}|$$

$$e_{\frac{\bar{x}}{\bar{y}}} \leq \frac{|\bar{y}|e_{\bar{x}}+|\bar{x}|e_{\bar{y}}}{|\bar{y}|^2}$$

$$\bar{x}, \bar{y} > 0$$

Absolute error is too sensitive to the value of the parameters shown:

- Large amounts for $\bar{x}, \bar{y}$ increase error propagation in multiplication
- Small amounts for $\bar{y}$ increase error propagation in division.

©Sharif University Of Technology

Error Propagation

Absolute Error

Relative Error

مثال ۱۰. مقدار $\pi\sqrt{۲}$ را با چهار رقم اعشار محاسبه نموده و حداکثر خطای این حاصل ضرب را نیز به دست آورید.

حل : داریم:

$$\pi = ۳/۱۴۱۶ + e_۱$$

$$\sqrt{۲} = ۱/۴۱۴۲ + e_۲$$

$$e_۱ \leq \frac{۱}{۲} \times ۱۰^{-۴}, \qquad e_۲ \leq \frac{۱}{۲} \times ۱۰^{-۴}$$

$$\pi\sqrt{۲} = (۳/۱۴۱۶ \times ۱/۴۱۴۲) + e_۳$$

©Sharif University Of Technology

Error Propagation

Absolute Error

Relative Error

$$e_۳ \leq ۳,۱۴۱۶ e_۲ + ۱,۴۱۴۲ e_۱$$

$$e_۳ \leq \frac{۱}{۲} \times ۱۰^{-۴}(۳,۱۴۱۶ + ۱,۴۱۴۲)$$

$$e_۳ \leq ۰,۵ \times ۱۰^{-۴}(۴,۵۵۵۸) = ۲,۲۷۷۹ \times ۱۰^{-۴}$$

اما

$$\pi\sqrt{۲} = ۴,۴۴۲۹ + e'_۳$$

©Sharif University Of Technology

Error Propagation

Absolute Error

Relative Error

چون حاصل‌ضرب اعداد ۳٫۱۴۱۶ و ۱٫۴۱۴۲ در محاسبهٔ $\pi\sqrt{۲}$ بیشتر از چهار رقم اعشار دارد، هنگام نمایش حاصل‌ضرب دو عدد مذکور با چهار رقم اعشار خطای دیگری مرتکب شده‌ایم و خطای حدی کل را با $e'_۳$ نشان داده‌ایم. برای $e'_۳$ داریم:

$$e'_۳ \leq \frac{۱}{۲} \times ۱۰^{-۴} + e_۳$$

$$e'_۳ \leq ۰٫۵ \times ۱۰^{-۴} + ۲٫۲۷۷۹ \times ۱۰^{-۴} = ۲٫۷۷۷۹ \times ۱۰^{-۴}$$

$$۴٫۴۴۲۹ - ۲٫۷۷۷۹ \times ۱۰^{-۴} \leq \pi\sqrt{۲} \leq ۴٫۴۴۲۹ + ۲٫۷۷۷۹ \times ۱۰^{-۴}$$

$$۴٫۴۴۲۶ \leq \pi\sqrt{۲} \leq ۴٫۴۴۳۲$$

©Sharif University Of Technology

Error Propagation

Absolute Error

Relative Error

Multiplication (*):

$\bar{x}$ and $\bar{y}$ are approximations of x and y $(\bar{x}, \bar{y} > 0)$

$$\delta_{\bar{x}} \leq \frac{e_{\bar{x}}}{\bar{x}} \qquad and \qquad \delta_{\bar{y}} \leq \frac{e_{\bar{y}}}{\bar{y}}$$

$$\delta_{\bar{x}\bar{y}} \leq \frac{e_{\bar{x}\bar{y}}}{\bar{x}\bar{y}} \leq \frac{\bar{y}e_{\bar{x}} + \bar{x}e_{\bar{y}}}{\bar{x}\bar{y}} = \frac{e_{\bar{x}}}{\bar{x}} + \frac{e_{\bar{y}}}{\bar{y}} = \delta_{\bar{x}} + \delta_{\bar{y}}$$

$$\delta_{\bar{x}\bar{y}} \leq \delta_{\bar{x}} + \delta_{\bar{y}}$$

©Sharif University Of Technology

Error Propagation

Absolute Error

Relative Error

Division (/):

$\bar{x}$ and $\bar{y}$ are approximations of x and y $(\bar{x}, \bar{y} > 0)$

$$\delta_{\bar{x}} \leq \frac{e_{\bar{x}}}{\bar{x}} \qquad and \qquad \delta_{\bar{y}} \leq \frac{e_{\bar{y}}}{\bar{y}}$$

$$\delta_{\frac{\bar{x}}{\bar{y}}} \leq \frac{\frac{e_{\bar{x}}}{\bar{y}}}{\frac{\bar{x}}{\bar{y}}} \leq \frac{\frac{\bar{y}e_{\bar{x}}+\bar{x}e_{\bar{y}}}{\bar{y}^2}}{\frac{\bar{x}}{\bar{y}}} = \frac{\frac{\bar{x}}{\bar{y}}\left(\frac{e_{\bar{x}}}{\bar{x}}+\frac{e_{\bar{y}}}{\bar{y}}\right)}{\frac{\bar{x}}{\bar{y}}} = \frac{e_{\bar{x}}}{\bar{x}} + \frac{e_{\bar{y}}}{\bar{y}} = \delta_{\bar{x}} + \delta_{\bar{y}}$$

$$\delta_{\frac{\bar{x}}{\bar{y}}} \leq \delta_{\bar{x}} + \delta_{\bar{y}}$$

©Sharif University Of Technology

Error Propagation

Absolute Error

Relative Error

$$\delta_{\frac{\bar{x}}{\bar{y}}} \leq \delta_{\bar{x}} + \delta_{\bar{y}} \qquad \bar{y} \neq 0$$

$$\delta_{\bar{x}\bar{y}} \leq \delta_{\bar{x}} + \delta_{\bar{y}} \qquad \bar{x}, \bar{y} > 0$$

©Sharif University Of Technology

An Example

Suppose a cylinder with a radius of 5.7 cm and a height of 10.5 cm. Estimate the absolute and relative errors of calculating the volume of it considering that all the values have been rounded using chopping model.(consider π = 3.14)



©Sharif University Of Technology

An Example

Suppose a cylinder with a radius of 5.7 cm and a height of 10.5 cm. Estimate the absolute and relative errors of calculating the volume of it considering that all the values have been rounded using chopping model.(consider π = 3.14)

Solution :

V = $h\,\pi\,r^2$

π **is symmetrically rounded off into two floating digits,**

thus $\rightarrow e_\pi \leq 0.5 \times 10^{-2}$

h **and** r **are measured by a device with the maximum error of** 10^{-1}

thus $\rightarrow$ $e_r \leq 10^{-1}$ **and** $e_h \leq 10^{-1}$

$$e(x \times y) \leq |x|e_y + |y|e_x \rightarrow \begin{cases} e(h\pi) \leq 10.5 \times 0.5 \times 10^{-2} + 3.14 \times 10^{-1} = 0.3665 \\ e(r \times r) \leq 2|r|e_r = 1.14 \end{cases}$$

$$\delta(xy) \leq \delta_x + \delta_y \qquad \rightarrow \begin{cases} \delta(h\pi) \leq 0.111162 \times 10^{-1} \\ \delta(r^2) \leq \delta_r + \delta_r = 0.35087 \times 10^{-1} \end{cases}$$



$$e_v = e_{h\pi \times r^2} \leq |h\pi|e_{r^2} + |r^2| \times e_{h\pi} = 0.494933 \times 10^2\,, \delta_v \leq 0.46204 \times 10^{-1}$$

©Sharif University Of Technology

An Example

Suppose a cylinder with a radius of 5.7 cm and a height of 10.5 cm. Estimate the absolute and relative errors of calculating the volume of it considering that all the values have been rounded.

Solution :

	$value$	$max(e)$	$max(\delta)$
h	10.5	0.1	0.00952
π	3.14	0.005	0.00159
r	5.7	0.01	0.01754
hπ	32.97	0.3665	0.01111
$r^2 = r \times r$	32.49	1.14	0.03508
$v = (h\pi).(r^2)$	1071.1953	49.49338	0.046204



©Sharif University Of Technology

An Example

Suppose a cylinder with a radius of 5.7 cm and a height of 10.5 cm. Estimate the absolute and relative errors of calculating the volume of it considering that all the values have been rounded.

Solution :

	$value$	$max(e)$	$max(\delta)$
h	10.5	0.1	0.00952
π	3.14	0.005	0.00159
r	5.7	0.01	0.01754
$h\pi$	32.97	0.3665	0.01111
$r^2 = r \times r$	32.49	1.14	0.03508
$v = (h\pi).(r^2)$	1071.1953	49.49338	0.046204



What if our computer supports just 3 digits mantissa?

©Sharif University Of Technology

An Example

Suppose a cylinder with a radius of 5.7 cm and a height of 10.5 cm. Estimate the absolute and relative errors of calculating the volume of it considering that all the values have been rounded.

Solution :

	$value$	$max(e)$	$max(\delta)$
h	10.5	$0.1 + \lvert 10.5 - 10.5 \rvert$	$0.00952 + \frac{\lvert 10.5 - 10.5 \rvert}{10.5}$
π	3.14	$0.005 + \lvert 3.14 - 3.14 \rvert$	$0.00159 + \frac{\lvert 3.14 - 3.14 \rvert}{3.14}$
r	5.7	$0.1 + \lvert 5.7 - 5.7 \rvert$	$0.01754 + \frac{\lvert 5.7 - 5.7 \rvert}{5.7}$
$h\pi$	33	$0.3665 + \lvert 33 - 32.97 \rvert$	$0.011116 + \frac{\lvert 33 - 32.97 \rvert}{33}$
$r^2 = r \times r$	32.5	$1.14 + \lvert 32.5 - 32.49 \rvert$	$0.035087 + \frac{\lvert 32.5 - 32.49 \rvert}{32.5}$
$v = (h\pi).(r^2)$	107	$49.4933 + \lvert 107 - 1071.1953 \rvert$	$0.04620 + \frac{\lvert 107 - 1071.1953 \rvert}{107}$

©Sharif University Of Technology

Formula Error

The Taylor series of $f(x)$ at a number a:

$$f(x) = f(a) + (x-a)f'(a) + \frac{(x-a)^2}{2!}f''(a) + \cdots$$

$$|f(x) - f(a)| \cong |x-a||f'(a)| = e(a)|f'(a)|$$

$$e_f \leq e_a|f'(a)|$$

©Sharif University Of Technology

Formula Error

The Taylor series of $f(x_1, x_2)$ at (a_1, a_2):

$$f(x_1, x_2) = f(a_1, a_2) + (x_1 - a_1)\frac{\partial f(a_1, a_2)}{\partial x_1} + (x_2 - a_2)\frac{\partial f(a_1, a_2)}{\partial x_2} + \cdots$$

$$|f(x_1, x_2) - f(a_1, a_2)| \cong e(a_1)\left|\frac{\partial f(a_1, a_2)}{\partial x_1}\right| + e(a_2)\left|\frac{\partial f(a_1, a_2)}{\partial x_2}\right|$$

$$e_f \leq e_{a_1}\left|\frac{\partial f(a_1, a_2)}{\partial x_1}\right| + e_{a_2}\left|\frac{\partial f(a_1, a_2)}{\partial x_2}\right|$$

©Sharif University Of Technology

Formula Error

Error of $f(x_1, x_2, \dots, x_n)$ at $\bar{a}=(a_1, a_2, \dots, a_n)$:

$$e_f = |f(x_1, x_2, \dots, x_n) - f(a_1, a_2, \dots, a_n)| \leq$$

$$e_{a_1} \left| \frac{\partial f}{\partial x_1} \right|_{\bar{a}} + e_{a_2} \left| \frac{\partial f}{\partial x_2} \right|_{\bar{a}} + \dots + e_{a_n} \left| \frac{\partial f}{\partial x_n} \right|_{\bar{a}}$$

©Sharif University Of Technology

An Example

Suppose a cylinder with a radius of 5.7 cm and a height of 10.5 cm. Estimate the absolute and relative errors of calculating the volume of it considering that all the values have been rounded using chopping model.(consider π = 3.14)

Solution :

V = $h\,\pi\,r^2$

$f = x\,y\,z^2$

$$e_h = e_x \leq 10^{-1}$$
$$e_\pi = e_y \leq 0.5 \times 10^{-2}$$
$$e_r = e_z \leq 10^{-1}$$



$$e_f \leq e_x\, yz^2 + e_y\, xz^2 + e_z\, 2xyz =$$

$$10^{-1} \times 3.14 \times (5.7)^2 + 0.5 \times 10^{-2} \times 10.5 \times (5.7)^2 +$$

$$10^{-1} \times 2 \times 10.5 \times 3.14 \times 5.7 = 0.494933 \times 10^2$$

$$\delta_f \leq \frac{e_f}{|f(\bar{a})|} = 0.46204 \times 10^{-1}$$

©Sharif University Of Technology

An Example

Compute the following expression with 4 digits mantissa and symmetric round-off for $x = 3.209$.

$$1.076x^3 + 0.319x^2 - 0.017x + 1.107$$

a) From left to right.

b) From right to left.

c) Compute the exact value.

d) What is the difference and why?

©Sharif University Of Technology

An Example

Compute the following expression with 4 digits mantissa and symmetric round-off for $x = 3.209$.

$$1.076x^3 + 0.319x^2 - 0.017x + 1.107$$

a) From left to right.

b) From right to left.

c) Compute the exact value.

d) What is the difference and why?

Solution:

$$((1.076 \times 3.209) \times 3.209) \times 3.209 = 35.56$$

$$(0.319 \times 3.209) \times 3.209 = 3.286$$

$$0.017 \times 3.209 = 0.054553 \rightarrow 0.05455$$

$$1.107 \rightarrow 1.107$$

©Sharif University Of Technology

An Example

$$1.076x^3 + 0.319x^2 - 0.017x + 1.107$$

a) From left to right $\rightarrow 39.91$

b) From right to left

c) Compute the exact value

d) What is the difference and why?

$$1.076x^3 \rightarrow 35.56$$
$$0.319x^2 \rightarrow 3.286$$
$$0.017x \rightarrow 0.05455$$
$$1.107 \rightarrow 1.107$$

Solution:

$$35.56 + 3.286 = 38.85$$
$$38.85 - 0.05455 = 38.80$$
$$38.80 + 1.107 = 39.91$$

©Sharif University Of Technology

An Example

$$1.076x^3 + 0.319x^2 - 0.017x + 1.107$$

a) From left to right $\rightarrow 39.91$

b) From right to left

c) Compute the exact value

d) What is the difference and why?

Solution:

$$\big(3.209 \times (3.209 \times 3.209)\big) \times 1.076 = 35.56$$

$$(3.209 \times 3.209) \times 0.319 = 3.286$$

$$3.209 \times 0.017 = 0.054553 \rightarrow 0.05455$$

$$1.107 \rightarrow 1.107$$

©Sharif University Of Technology

An Example

$$1.076x^3 + 0.319x^2 - 0.017x + 1.107$$

a) From left to right $\rightarrow 39.91$

b) From right to left $\rightarrow 39.90$

c) Compute the exact value

d) What is the difference and why?

$$1.076x^3 \rightarrow 35.56$$
$$0.319x^2 \rightarrow 3.286$$
$$0.017x \rightarrow 0.05455$$
$$1.107 \rightarrow 1.107$$

Solution:

$$1.107 - 0.05455 = 1.052$$
$$1.052 + 3.286 = 4.338$$
$$4.338 + 35.56 = 39.90$$

©Sharif University Of Technology

An Example

$$1.076x^3 + 0.319x^2 - 0.017x + 1.107$$

a) From left to right $\rightarrow 39.91$

b) From right to left $\rightarrow 39.90$

c) Compute the exact value $\rightarrow 39.894105201004$

d) What is the difference and why?

Solution:

$$1.076x^3 \rightarrow 35.56$$
$$0.319x^2 \rightarrow 3.285$$
$$0.017x \rightarrow 0.05455$$
$$1.107 \rightarrow 1.107$$

$$1.076x^3 + 0.319x^2 - 0.017x + 1.107 = 39.894105201004$$

©Sharif University Of Technology

An Example

$$1.076x^3 + 0.319x^2 - 0.017x + 1.107$$

a) From left to right $\rightarrow 39.91$

b) From right to left $\rightarrow 39.90$

c) Compute the exact value

d) What is the difference and why?

Solution:

We better initially deal with <u>the least significant numbers</u> in any computational system where the number of digits are limited, i.e. small numbers show themselves better if used prior to others.

©Sharif University Of Technology

Process Graph

$$u = (x + y) * z$$



©Sharif University Of Technology

An Example

Draw process graph of $\mathrm{v} = \pi r^2 h$

1. From left to right
2. From right to left

©Sharif University Of Technology

An Example

Draw process graph of $\mathrm{v} = \pi r^2 h$

1. From left to right
2. From right to left

Solution :

$\mathrm{v} = \pi r^2 h$

$= \pi r r h$

$= (\pi . r).(r . h)$



©Sharif University Of Technology

An Example

Draw process graph of $\mathrm{v} = \pi r^2 h$

1. From left to right
2. From right to left

Solution :

$$\mathrm{v} = \pi r^2 h$$
$$= hrr\pi$$
$$= (h.r).(r.\pi)$$



©Sharif University Of Technology

Stability

Algorithm (method)

- Stable : $E_n \approx cE_0$ (linearly)
- Unstable : $E_n \approx c^n E_0$ c≥ 1 (exponentially)

problem

- Inherent unstable → Example: Wilkinson problem,
- Induced unstable

Wilkinson problem: roots of

$$P_{20}(x) = (x-1)(x-2)\dots(x-20) = x^{20} - 210x^{19} + \cdots + 20!$$

©Sharif University Of Technology

Any questions?