Assignment 9: Spatial Analysis in R

Sophia Bryson

OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics (ENV872L) on spatial analysis.

Directions

- 1. Change "Student Name" on line 3 (above) with your name.
- 2. Use the lesson as a guide. It contains code that can be modified to complete the assignment.
- 3. Work through the steps, creating code and output that fulfill each instruction.
- 4. Be sure to answer the questions in this assignment document. Space for your answers is provided in this document and is indicated by the ">" character. If you need a second paragraph be sure to start the first line with ">". You should notice that the answer is highlighted in green by RStudio.
- 5. When you have completed the assignment, **Knit** the text and code into a single HTML file.
- 6. After Knitting, please submit the completed exercise (PDF file) in Sakai. Please add your last name into the file name (e.g., "Fay A10 SpatialAnalysis.pdf") prior to submission.

DATA WRANGLING

Set up your session

library(mapview)

- 1. Check your working directory
- 2. Import libraries: tidyverse, sf, leaflet, and mapview


```
#1.
getwd()
## [1] "Z:/ENV872/Environmental_Data_Analytics_2022/Assignments"
library(tidyverse)
## -- Attaching packages ------ tidyverse 1.3.1 --
## v ggplot2 3.3.5
                     v purrr
                               0.3.4
## v tibble 3.1.6
                     v dplyr
                               1.0.8
## v tidyr
            1.2.0
                     v stringr 1.4.0
                     v forcats 0.5.1
## v readr
            2.1.2
## -- Conflicts -----
                                           ----- tidyverse_conflicts() --
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()
                   masks stats::lag()
library(sf)
## Linking to GEOS 3.9.1, GDAL 3.2.1, PROJ 7.2.1; sf_use_s2() is TRUE
library(leaflet)
```

Read (and filter) county features into an sf dataframe and plot

In this exercise, we will be exploring stream gage height data in Nebraska corresponding to floods occurring there in 2019. First, we will import from the US Counties shapefile we've used in lab lessons, filtering it this time for just Nebraska counties. Nebraska's state FIPS code is 31 (as North Carolina's was 37).

- 3. Read the cb_2018_us_county_20m.shp shapefile into an sf dataframe, filtering records for Nebraska counties (State FIPS = 31)
- 4. Reveal the dataset's coordinate reference system
- 5. Plot the records as a map (using mapview or ggplot)

```
#3. Read in Counties shapefile into an sf dataframe, filtering for just NE counties
counties.sf <- st_read("../Data/Spatial/cb_2018_us_county_20m.shp") %>%
               filter(STATEFP == "31")
## Reading layer `cb_2018_us_county_20m' from data source
     `Z:\ENV872\Environmental_Data_Analytics_2022\Data\Spatial\cb_2018_us_county_20m.shp'
##
##
     using driver `ESRI Shapefile'
## Simple feature collection with 3220 features and 9 fields
## Geometry type: MULTIPOLYGON
## Dimension:
                  XY
## Bounding box:
                  xmin: -179.1743 ymin: 17.91377 xmax: 179.7739 ymax: 71.35256
## Geodetic CRS: NAD83
#4. Reveal the CRS of the counties features
st_crs(counties.sf)
## Coordinate Reference System:
##
     User input: NAD83
##
     wkt:
##
  GEOGCRS["NAD83",
##
       DATUM["North American Datum 1983",
##
           ELLIPSOID["GRS 1980",6378137,298.257222101,
##
               LENGTHUNIT["metre",1]]],
       PRIMEM["Greenwich",0,
##
           ANGLEUNIT["degree", 0.0174532925199433]],
##
##
       CS[ellipsoidal,2],
##
           AXIS["latitude", north,
##
               ORDER[1],
               ANGLEUNIT["degree", 0.0174532925199433]],
##
##
           AXIS["longitude", east,
               ORDER[2],
##
               ANGLEUNIT["degree", 0.0174532925199433]],
##
##
       ID["EPSG",4269]]
#5. Plot the data
ggplot() +
 geom_sf(data = counties.sf)
```



6. What is the EPSG code of the Counties dataset? Is this a geographic or a projected coordinate reference system? (Or, does this CRS use angular or planar coordinate units?) To what datum is this CRS associated? (Tip: look the EPSG code on https://spatialreference.org)

ANSWER: The counties dataset is in NAD83 (North American Datum 1983), a projected coordinate system. It is associated with the 1983 North American Datum (EPSG 4269).

Read in gage locations csv as a dataframe, then display the column names it contains

Next we'll read in some USGS/NWIS gage location data added to the Data/Raw folder. These are in the NWIS_SiteInfo_NE_RAW.csv file.(See NWIS_SiteInfo_NE_RAW.README.txt for more info on this dataset.)

- 7. Read the NWIS SiteInfo NE RAW.csv file into a standard dataframe.
- 8. Display the column names of this dataset.

9. What columns in the dataset contain the x and y coordinate values, respectively?

> ANSWER: > The x coordinate values are contained in the column "dec_long_va". The y coordinate values are contained in the column "dec_lat_va".

Convert the dataframe to a spatial features ("sf") dataframe

- 10. Convert the dataframe to an sf dataframe.
 - Note: These data use the same coordinate reference system as the counties dataset
- 11. Display the column names of the resulting sf dataframe

12. What new field(s) appear in the sf dataframe created? What field(s), if any, disappeared?


ANSWER: In the spatial dataframe, the longitude and latitude columns were replaced by a single unified geometry column containing lat/long pairs.

Plot the gage locations on top of the counties

- 13. Use ggplot to plot the county and gage location datasets.
- Be sure the datasets are displayed in different colors
- Title your plot "NWIS Gage Locations in Nebraska"
- Subtitle your plot with your name

```
#13. Plot the gage locations atop the county features
ggplot() +
   geom_sf(data = counties.sf, color = "olivedrab", fill = "olivedrab3") +
   geom_sf(data = gages.sf, alpha = 0.8, size = 1.3, color = "tomato4") +
   labs(title = "NWIS Gage Locations in Nebraska",
        subtitle = "Sophia Bryson")
```

NWIS Gage Locations in Nebraska Sophia Bryson



Read in the gage height data and join the site location data to it.


Lastly, we want to attach some gage height data to our site locations. I've constructed a csv file listing many of the Nebraska gage sites, by station name and site number along with stream gage heights (in meters) recorded during the recent flood event. This file is titled NWIS_SiteFlowData_NE_RAW.csv and is found in the Data/Raw folder.

- 14. Read the NWIS_SiteFlowData_NE_RAW.csv dataset in as a dataframe.
- 15. Show the column names .
- 16. Join our site information (already imported above) to these gage height data.
 - The site_no and station_nm can both/either serve as joining attributes.
 - Construct this join so that the result only includes spatial features where both tables have data.
- 17. Show the column names in this resulting spatial features object
- 18. Show the dimensions of the resulting joined dataframe

Map the pattern of gage height data

Now we can examine where the flooding appears most acute by visualizing gage heights spatially. 19. Plot the gage sites on top of counties (using mapview, ggplot, or leaflet) * Show the magnitude of gage height by color, shape, other visualization technique.

Gage height of Nebraska NWIS gages



SPATIAL ANALYSIS


Up next we will do some spatial analysis with our data. To prepare for this, we should transform our data into a projected coordinate system. We'll choose UTM Zone 14N (EPGS = 32614).

Transform the counties and gage site datasets to UTM Zone 14N

- 20. Transform the counties and gage sf datasets to UTM Zone 14N (EPGS = 32614).
- 21. Using mapview or ggplot, plot the data so that each can be seen as different colors

```
#20 Transform the counties and gage location datasets to UTM Zone 14
counties.trans.sf <- counties.sf %>% st_transform(crs = 32614)
gages.trans.sf <- gages.sf %>% st_transform(crs = 32614)

#21 Plot the data
ggplot() +
   geom_sf(data = counties.trans.sf, color = "olivedrab", fill = "olivedrab3") +
   geom_sf(data = gages.trans.sf, alpha = 0.8, size = 1.3, color = "tomato4")
```



Select the gages falling within a given county

Now let's zoom into a particular county and examine the gages located there. 22. Select Lancaster county from your county sf dataframe 23. Select the gage sites falling within that county * Use either matrix subsetting or tidy filtering 24. Create a plot showing: * all Nebraska counties, * the selected county, * and

the gage sites in that county

```
#22 Select the county
lancaster <- counties.trans.sf %>% filter(NAME == "Lancaster")

#23 Select gages within the selected county
lancasterGages <- gages.trans.sf[lancaster,]

#24 Plot
ggplot() +
   geom_sf(data = counties.trans.sf, color = "olivedrab", fill = "olivedrab3") +
   geom_sf(data = lancaster, color = "olivedrab", fill = "olivedrab4") +
   geom_sf(data = lancasterGages, color = "tomato4", size = 1, alpha = 0.7) +
   labs(title = "NWIS gages in Lancaster County, Nebraska")</pre>
```

NWIS gages in Lancaster County, Nebraska

