## **LABORATORY 2**

This assignment is due for Week 2.

Write a C application with a menu based console interface which solves the problems below. Each requirement must be resolved using at least one function. All functions need to be specified.

- 1. a. Generate all the prime numbers smaller than a given natural number n.
  - b. Given a vector of numbers, find the longest increasing contiguous subsequence, such the sum of that any 2 consecutive elements is a prime number.
- 2. a. Generate the first n prime numbers (n is a given natural number).
  - b. Given a vector of numbers, find the longest contiguous subsequence such that any two consecutive elements are relatively prime.
- 3. a. Print the Pascal triangle of dimension n of all combinations C(m,k) of m objects taken by k, k = 0, 1, ..., m, for line m, where m = 1, 2, ..., n.
  - b. Given a vector of numbers, find the longest contiguous subsequence of prime numbers.
- 4. a. Compute the approximated value of square root of a positive real number. Use a given precision.
  - b. Given a vector of numbers, find the longest contiguous subsequence such that the difference of any two consecutive elements is a prime number.
- 5. a. Print the exponent of a prime number p from the decomposition in prime factors of a given number n (n is a non-null natural number).
  - b. Given a vector of numbers, find the longest contiguous subsequence such that any two consecutive elements are relatively prime.
- 6. a. Read a sequence of natural numbers (sequence ended by 0) and determine the number of 0 digits of the product of the read numbers.
  - b. Given a vector of numbers, find the longest contiguous subsequence such that the sum of any two consecutive elements is a prime number.
- 7. a. Read sequences of positive integer numbers (reading of each sequence ends by 0, reading of all the sequences ends by -1) and determine the maximum element of each sequence and the maxim element of the global sequence.
  - b. Given a vector of numbers, find the longest contiguous subsequence such that all elements are in a given interval.

- 8. a. Determine the value x^n, where x is a real number and n is a natural number, by using multiplication and squared operations.
  - b. Given a vector of numbers, find the longest contiguous subsequence such that any two consecutive elements have contrary signs.
- 9. a. Decompose a given natural number in its prime factors.
  - b. Given a vector of numbers, find the longest contiguous subsequence such that any consecutive elements contain the same digits.
- 10. a. Decompose a given even natural number, greater than 2, as a sum of two prime numbers (Goldbach's conjecture).
  - b. Given a vector of numbers, find the longest contiguous subsequence such that any consecutive elements have at least 2 distinct digits in common.
- 11. a. Determine the first n pairs of twin numbers, where n is a given natural and non-null number. Two prime numbers p and q are called twin if q p = 2.
  - b. Given a vector of numbers, find the longest decreasing contiguous subsequence.
- 12. a. Determine all the numbers smaller than a given natural and non-null number n and that are relatively prime to n.
  - b. Given a vector of numbers, find the longest contiguous subsequence with the maximum sum.
- 13. a. Determine the first 8 natural numbers (x1, x2, ..., x8) greater than 2 with the following property: all the natural numbers smaller than  $x_i$  and that are relatively prime with  $x_i$  (except for the number 1) are prime, i = 1, 2, ..., n.
  - b. Given a vector of numbers, find the longest contiguous subsequence such that any consecutive elements contain the same digits.