

1.3 1 Solve the following congruence

**d**  $19x \equiv 1 \pmod{36}$

**Ans**

$$19x \equiv 1 \pmod{36}$$

$$19x = 1 + 36n, \text{ for } n \in \mathbb{Z}$$

$$\Rightarrow 1 = 19x - 36n$$

$$1 = 19(19) - 36(10)$$

Therefore,  $x \equiv 19 \pmod{36}$

□

**4** Solve the following congruence:  $20x \equiv 12 \pmod{72}$

**Ans** Since  $(20, 72) = 4$ , there exists 4 solutions.

$$20x \equiv 12 \pmod{72}$$

$$20x = 12 + 72n, \text{ for } n \in \mathbb{Z}$$

$$\Rightarrow 5x = 3 + 18n$$

$$5x \equiv 3 \pmod{18}$$

Then,  $x \equiv 15 \pmod{18} \Rightarrow 18 \mid (5x - 3)$

Therefore,

$$x \equiv 15 \pmod{18}$$

$$x \equiv 33 \pmod{18}$$

$$x \equiv 51 \pmod{18}$$

$$x \equiv 69 \pmod{18}$$

□

**7** The smallest positive solution of the congruence  $ax \equiv 0 \pmod{n}$  is called the additive order of  $a$  modulo  $n$ . Find the additive orders of each of the following elements, by solving the appropriate congruences.

**b** 7 modulo 12

**Ans** The smallest positive solution:  $7x \equiv 0 \pmod{12}$

That is, the smallest positive integer  $x$  such that  $12 \mid 7x \Rightarrow x = 4$

Therefore, the additive order of 7 modulo 12 is  $x = 12$

□

**d** 12 modulo 18

**Ans** The smallest positive solution:  $12x \equiv 0 \pmod{18}$

That is, the smallest positive integer  $x$  such that  $18 \mid 12x \Rightarrow x = 3$

Therefore, the additive order of 12 modulo 18 is  $x = 3$

□

**14** Find the units digit of  $3^{29} + 11^{12} + 15$ .

*Hint:* Choose an appropriate modulus  $n$ , and then reduce modulo  $n$ .

**Ans** Since  $3^4 = 81$  with a units digit of 1,

then  $3^{29} = (3^4)^7 \cdot 3$  with a units digit of 3

Since  $11^2 = 121$  with a units digit of 1,

then  $11^{12} = (11^2)^6$  with a units digit of 1

Therefore, the units digit of  $3^{29} + 11^{12} + 15$  is:  $1 + 3 + 5 = 9$

□

**16** Solve the following congruences by trial and error.

**a**  $x^3 + 2x + 2 \equiv 0 \pmod{5}$

**Ans** By trial and error

$$x = 1 \Rightarrow 5 \mid (1)^3 + 2(1) + 2 = 5$$

$$x = 2 \Rightarrow 5 \nmid (2)^3 + 2(2) + 2 = 14$$

$$x = 3 \Rightarrow 5 \mid (3)^3 + 2(3) + 2 = 35$$

$$x = 4 \Rightarrow 5 \nmid (4)^3 + 2(4) + 2 = 74$$

Therefore,

$x \equiv 1 \pmod{5}$  and  $x \equiv 3 \pmod{5}$

□

**20** Solve the following system of congruences.

$$2x \equiv 5 \pmod{7}$$

$$3x \equiv 4 \pmod{8}$$

**Ans** Simplifying the congruences first,

$$2x \equiv 5 \pmod{7}$$

$$2x \equiv 5 \pmod{7}$$

$$2v \equiv 1 \pmod{7}$$

$$2v = 1 - 7n, \text{ for } n \in \mathbb{Z}$$

$$\Rightarrow 1 = 2v + 7n$$

$$1 = 2(4) + 7(-1)$$

$$\Rightarrow x \equiv 4v \pmod{7}$$

Therefore,

$$2x \equiv 4 \cdot 5 \pmod{7}$$

$$x \equiv 6 \pmod{7}$$

And  $3x \equiv 4 \pmod{8}$

$$3x \equiv 4 \pmod{8}$$

$$3v \equiv 1 \pmod{8}$$

$$3v = 1 - 8n, \text{ for } n \in \mathbb{Z}$$

$$\Rightarrow 1 = 3v + 8n$$

$$1 = 3(3) + 8(-1)$$

$$\Rightarrow x \equiv 3v \pmod{8}$$

Therefore,

$$3x \equiv 3 \cdot 4 \pmod{8}$$

$$x \equiv 4 \pmod{8}$$

Now the system can be solved using the Chinese Remainder Theorem:

$$x \equiv 6 \pmod{7}$$

$$x \equiv 4 \pmod{8}$$

Since  $(n_1, n_2) = (7, 8) = 1$ , let  $u_1 = 7k_1$  and  $u_2 = 8k_2$

Then

$$\begin{aligned}u_1 + u_2 = 1 &\Rightarrow 7k_1 + 8k_2 = 1 \\1 &= 7(-1) + 8(1)\end{aligned}$$

Thus

$$\begin{aligned}u_1 &= 7(-1) = -7 \equiv 1 \pmod{8} \\u_1 &= 7(-1) = -7 \equiv 0 \pmod{7}\end{aligned}$$

And

$$\begin{aligned}u_2 &= 8(1) = 8 \equiv 0 \pmod{8} \\u_2 &= 8(1) = 8 \equiv 1 \pmod{7}\end{aligned}$$

Therefore,

$$\begin{aligned}x &= 6u_1 + 4u_2 \\&= 6(-7) + 4(8) \\&= -10\end{aligned}$$

Therefore, the general solution with the smallest nonnegative integer is

$$\begin{aligned}x &\equiv -10 \pmod{n_1 n_2} \\x &\equiv -10 \pmod{56} \\x &\equiv 46 \pmod{56}\end{aligned}$$

□

**1.4    2** Make multiplication tables for the following sets.

□

□

**Table 1: b:** Multiplication table of  $\mathbb{Z}_7$ 

$\times$	[0]	[1]	[2]	[3]	[4]	[5]	[6]
[0]	[0]	[0]	[0]	[0]	[0]	[0]	[0]
[1]	[0]	[1]	[2]	[3]	[4]	[5]	[6]
[2]	[0]	[2]	[4]	[6]	[1]	[3]	[5]
[3]	[0]	[3]	[6]	[2]	[5]	[1]	[4]
[4]	[0]	[4]	[1]	[5]	[2]	[6]	[3]
[5]	[0]	[5]	[3]	[1]	[6]	[4]	[2]
[6]	[0]	[6]	[5]	[4]	[3]	[2]	[1]

**Table 2: c:** Multiplication table of  $\mathbb{Z}_8$ 

$\times$	[0]	[1]	[2]	[3]	[4]	[5]	[6]	[7]
[0]	[0]	[0]	[0]	[0]	[0]	[0]	[0]	[0]
[1]	[0]	[1]	[2]	[3]	[4]	[5]	[6]	[7]
[2]	[0]	[2]	[4]	[6]	[0]	[2]	[4]	[6]
[3]	[0]	[3]	[6]	[1]	[4]	[7]	[2]	[5]
[4]	[0]	[4]	[0]	[4]	[0]	[4]	[0]	[4]
[5]	[0]	[5]	[2]	[7]	[4]	[1]	[6]	[3]
[6]	[0]	[6]	[4]	[2]	[0]	[6]	[4]	[2]
[7]	[0]	[7]	[5]	[4]	[3]	[2]	[1]	[1]

- 6 Let  $m$  and  $n$  be positive integers such that  $m \mid n$ . Show that for any integer  $a$ , the congruence class  $[a]_m$  is the union of the congruence classes  $[a]_n, [a+m]_n, [a+2m]_n, \dots, [a+n-m]_n$

**Ans**

□

- 9 Let  $(a, n) = 1$ . The smallest positive integer  $k$  such that  $a^k \equiv 1 \pmod{n}$  is called the **multiplicative order** of  $[a]$  in  $\mathbb{Z}_n^\times$

**b** Find the multiplicative orders of  $[2]$  and  $[5]$  in  $\mathbb{Z}_{17}^\times$ .

**Ans** Show  $2^k \equiv 1 \pmod{17}$ , for  $k \in \mathbb{Z}$

Then,  $2^k = 1 + 17n$ , for  $n \in \mathbb{Z}$

Then,  $n = (2^k - 1)/17$

Therefore, for  $n$  to be an integer,  $k = 8$ .

Similarly, show  $5^k \equiv 1 \pmod{17}$ , for  $k \in \mathbb{Z}$

Then,  $5^k = 1 + 17n$ , for  $n \in \mathbb{Z}$

Then,  $n = (5^k - 1)/17$

Therefore, for  $n$  to be an integer,  $k = 16$ .

Therefore, the multiplicative order of  $[2]$  and  $[5]$  in  $\mathbb{Z}_{17}^\times$  is  $k = 8$

□

- 10 Let  $(a, n) = 1$ . If  $[a]$  has multiplicative order  $k$  in  $\mathbb{Z}_n^\times$ , show that  $k \mid \varphi(n)$ .

**Ans**

□

- 13 An element  $[a]$  of is said to be **idempotent** if  $[a]^2 = [a]$ .

**b** Find all idempotent elements of  $\mathbb{Z}_{10}^\times$  and  $\mathbb{Z}_{30}^\times$ .

**Ans**

□

- 15 If  $n$  is not a prime power, show that  $\mathbb{Z}_n$  has an idempotent element different from  $[0]$  and  $[1]$ .

*Hint:* Suppose that  $n = bc$ , with  $(b, c) = 1$ . Solve the simultaneous congruences  $x \equiv 1 \pmod{b}$  and  $x \equiv 0 \pmod{c}$ .

**Ans**

□

- 20 Show that  $\varphi(1) + \varphi(p) + \dots + \varphi(p^\alpha) = \varphi^\alpha$  for any prime number  $p$  and any positive integer  $\alpha$ .

**Ans**

□

**26** Let  $p = 2k + 1$  be a prime number. Show that if  $a$  is an integer such that  $p \nmid a$ , then either  $a^k \equiv 1 \pmod{p}$  or  $a^k \equiv -1 \pmod{p}$

**Ans**

□