

1. With 3 n -sided rolls, there are n^3 possibilities.

The probability that either of the pair of persons roll the same face of the die is therefore $n/n^3 = 1/n^2$.

Therefore

$$\begin{aligned} P(A_{12}) = P(A_{13}) = P(A_{23}) &= \frac{n}{n^3} \\ &= \frac{1}{n^2} \end{aligned}$$

But if both the events A_{12} and A_{13} takes place, that is both persons 1 and 2 and persons 1 and 3 roll the same face, then that yields A_{23} .

That is, if both persons 1 and 2 and persons 1 and 3 roll the same face, then that implies persons 1 and 3 rolled the same face.

But the outcome of person 3's roll is not dependent on the other persons.

That is, pairwise A_{12} and A_{13} , A_{12} and A_{23} , and A_{13} and A_{23} are independent.

But if considered individually, they are dependent. □

2. Consider the following counter-example with two independent tosses of a fair coin.

Let events $B = \{HT, HH\}$ and $C = \{HT, TT\}$ represent tosses where they landed heads and tails respectively.

Let $A = \{HT, TH\}$ be the event that exactly one toss resulted in heads.

Then,

$$P(A) = P(B) = P(C) = \frac{1}{2}$$

And,

$$P(A \cup B) = P(A \cup C) = \frac{1}{4}$$

Therefore, A and B and A and C are both independent events.

Therefore, B and C are also independent events.

However,

$$P(A \cap (B \cup C)) = \frac{1}{4} \neq P(A)P(B \cup C) = \frac{1}{2} \cdot \frac{3}{4}$$

Therefore, A and $B \cup C$ are dependent □

3.

□

4. Let p_5 denote the longer path of 5 links from A to B ,
and p_3 denote the shorter path of 3 links.

Given, the probability of links failing independently is q .

Therefore, the probability of links not failing is $1 - q$.

For a successful transmission, all of the links have to not fail.

Therefore, for path p_5 the probability is $P(p_5) = (1 - q)^5$

and for path p_3 the probability is $P(p_3) = (1 - q)^3$.

Since the paths are independent of each other,

$$\begin{aligned} P(p_5 \cap p_3) &= P(p_5) \cdot P(p_3) \\ &= (1 - q)^5 \cdot (1 - q)^3 \\ &= (1 - q)^8 \end{aligned}$$

That is, the probability of both the paths not failing is $(1 - q)^8$.

Therefore, the probability of either the paths not failing for a successful transmission from terminal A to B is:

$$\begin{aligned} P(p_5 \cup p_3) &= P(p_5) + P(p_3) - P(p_5 \cap p_3) \\ &= (1 - q)^5 + (1 - q)^3 - (1 - q)^8 \end{aligned}$$

□

5.

□

6.

□

7. Given:

$$P(\text{qualified}) = q,$$

$$P(\text{not qualified}) = 1 - q,$$

$$P(\text{correct answer} \mid \text{qualified}) = p$$

$$P(\text{incorrect answer} \mid \text{not qualified}) = p$$

Therefore,

$$\begin{aligned}
 P(>15 \text{ correct} \mid \text{qualified}) &= \frac{P(>15 \text{ correct} \cap \text{qualified})}{P(\text{qualified})} \\
 &= \frac{q \sum_{k=15}^{20} \binom{20}{15} (p)^{15} (1-p)^5}{q \sum_{k=15}^{20} \binom{20}{15} (p)^{15} (1-p)^5 + (1-q) \sum_{k=15}^{20} \binom{20}{15} (p)^{15} (1-p)^5} \\
 &= \frac{q \sum_{k=15}^{20} \binom{20}{15} (p)^{15} (1-p)^5}{\sum_{k=15}^{20} \binom{20}{15} (p)^{15} (1-p)^5} \\
 &= q
 \end{aligned}$$

□

8. Let C represent the 20 random distinct cars chosen for a test drive.

(a) To find $P(K = 0 \mid C)$, without replacement:

$$\begin{aligned}
 P(K = 0 \mid C) &= \frac{P(K = 0)P(C \mid K = 0)}{\sum_{i=0}^9 P(K = i)P(C \mid K = i)} \\
 &= \frac{\binom{100}{20}}{\sum_{i=0}^9 \binom{100-i}{20}} \\
 &\approx 0.227
 \end{aligned}$$

□

(b) To find $P(K = 0 \mid C)$, with replacement:

$$\begin{aligned}
 P(K = 0 \mid C) &= \frac{P(K = 0)P(C \mid K = 0)}{\sum_{i=0}^9 P(K = i)P(C \mid K = i)} \\
 &= \frac{100^{20}}{\sum_{i=0}^9 (100-i)^{20}} \\
 &\approx 0.213
 \end{aligned}$$

□

9.

□

10. The permutations of a word is given by:

$$\frac{(\text{length of word})!}{(\text{repetitions of A})!(\text{repetitions of B})! \dots (\text{repetitions of Z})!}$$

(a) Since there are no repeating characters, the permutation is simply:

$$\text{permutations} = \text{length}(\text{children})!$$

$$= 8!$$

$$= 40320$$

□

(b) Since the characters o repeats 2 times, k repeats 2 times, and e repeats 3 times:

$$\text{permutations} = \frac{\text{length}(\text{bookkeeper})!}{(\text{repetitions of } o)! (\text{repetitions of } k)! (\text{repetitions of } e)!}$$

$$= \frac{10!}{2!2!3!}$$

$$= 151200$$

□