


This article was downloaded by: [University of Cambridge]

On: 20 December 2014, At: 19:04

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Stochastic Analysis and Applications

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Isaa20>

Stochastic nagumo's viability theorem

J.P. Aubin ^a & G. DA Prato ^b

^a CEREMADE , Université de Paris-Dauphine , Paris, 75775, France

^b Scuola Normale Superiore di Pisa , Pisa, 56126, Italy

Published online: 03 Apr 2007.

To cite this article: J.P. Aubin & G. DA Prato (1995) Stochastic nagumo's viability theorem, Stochastic Analysis and Applications, 13:1, 1-11, DOI: [10.1080/07362999508809379](https://doi.org/10.1080/07362999508809379)

To link to this article: <http://dx.doi.org/10.1080/07362999508809379>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

STOCHASTIC NAGUMO'S VIABILITY THEOREM

J. P. Aubin

CEREMADE, Université de Paris-Dauphine, 75775 Paris, France

G. Da Prato

Scuola Normale Superiore di Pisa, 56126 Pisa, Italy

ABSTRACT

This paper is devoted to viability of random set-valued variables by stochastic differential equations, characterized in terms of stochastic tangent sets to random set-valued variables.

1 INTRODUCTION

The main aim of this paper is to extend to the stochastic case Nagumo's Theorem on viability properties of closed subsets with respect to a differential equation. In [2, Aubin & Da Prato], only invariance theorems were presented : we proved that under adequate stochastic tangential conditions, any solution to the stochastic differential equation starting from \mathcal{K} , if any, remains in \mathcal{K} , whereas in viability theorems, we prove the existence of one solution which is viable in \mathcal{K} .

Let us consider a closed subset K of $X := \mathbb{R}^n$ and a stochastic differential equation

$$d\xi = f(\xi(t))dt + g(\xi(t))dW(t)$$

the solution of which is given by the formula

$$\xi(t) = \xi(0) + \int_0^t f(\xi(s))ds + \int_0^t g(\xi(s))dW(s)$$

when one of the following conditions is satisfied:

- f and g are Lipschitz functions
- f and g are uniformly continuous and monotone

We want to characterize the (stochastic) viability property of K with respect to the pair (f, g) : for any random variable x in K , there exists a solution ξ to the stochastic differential equation starting at x which is viable in K , in the sense that

$$\forall t \in [0, T], \text{ for almost all } \omega \in \Omega, \xi_\omega(t) \in K_\omega$$

The construction of approximate viable solutions we provide when f and g are uniformly continuous is interesting for its own sake, and may be useful in other problems.

For that purpose, we use the concept of contingent set to a subset introduced in [2, Aubin & Da Prato]. Let us consider a \mathcal{F}_t -random variable $x \in K$.

We define the stochastic contingent set $\mathcal{T}_K(t, x)$ to K at x (with respect to \mathcal{F}_t) as the set of pairs (γ, v) of \mathcal{F}_t -random variables satisfying the following property: There exist sequences of $h_n > 0$ converging to 0 and of \mathcal{F}_{t+h_n} -measurable random variables a^n and b^n such that

$$\begin{cases} i) & \mathbf{E}(\|a^n\|^2) \rightarrow 0 \\ ii) & \mathbf{E}(\|b^n\|^2) \rightarrow 0 \\ iii) & \mathbf{E}(b^n) = 0 \\ iv) & b^n \text{ is independent of } \mathcal{F}_t \end{cases}$$

and satisfying for almost all $\omega \in \Omega$,

$$\forall n \geq 0, x_\omega + \gamma_\omega h_n + v_\omega(W_\omega(t+h_n) - W_\omega(t)) + h_n a_\omega^n + \sqrt{h_n} b_\omega^n \in K_\omega$$

Then we shall prove in essence that the following conditions are equivalent:

1. — The subset K enjoys the viability property with respect to the pair (f, g)

2. — for every \mathcal{F}_t -random variable x viable in K ,

$$(f(x), g(x)) \in T_K(t, x)$$

For instance, this condition means that for every \mathcal{F}_t -random variable x viable in K

•

$$f(x) \in K \text{ \& } g(x) \in K$$

when K is a vector subspace,

•

$$\langle x, g(x) \rangle = 0 \text{ \& } \langle x, f(x) \rangle + \frac{1}{2} \|g(x)\|^2 = 0$$

when K is the unit sphere

•

$$\langle x, g(x) \rangle = 0 \text{ \& } \langle x, f(x) \rangle + \frac{1}{2} \|g(x)\|^2 \leq 0$$

when K is the unit ball.

We mention that an elementary calculus of stochastic tangent sets to direct images, inverse images and intersections of closed subsets can be found in [2, Aubin & Da Prato].

2 STOCHASTIC TANGENT SETS

Let us consider a complete probability space (Ω, \mathcal{F}, P) , an increasing family of σ -sub- algebras $\mathcal{F}_t \subset \mathcal{F}$ and a finite dimensional vector-space $X := \mathbb{R}^n$. Moreover $W(t), t \geq 0$ is a real standard Brownian motion such that $W(t)$ is \mathcal{F}_t -measurable and $W(t+h) - W(t)$ is independent of \mathcal{F}_t for any $h \geq 0$.

The constraints are defined by closed subsets $K_\omega \subset X$, where the set-valued map

$$K : \omega \in \Omega \mapsto K_\omega \subset X$$

is assumed to be \mathcal{F}_0 - measurable (which can be regarded as a random set-valued variable).

We denote by \mathcal{K} the subset

$$\mathcal{K} := \{u \in L^2(\Omega, \mathcal{F}, P) \mid \text{for almost all } \omega \in \Omega, \ u_\omega \in K_\omega\}$$

For simplicity, we restrict ourselves to scalar \mathcal{F}_t -Wiener processes $W(t)$.

Definition 2.1 (Stochastic Contingent Set) *Let us consider a \mathcal{F}_t -random variable $x \in K$ (i.e., a \mathcal{F}_t -measurable selection of K).*

We define the stochastic contingent set $T_K(t, x)$ to K at x (with respect to \mathcal{F}_t) as the set of pairs (γ, v) of \mathcal{F}_t -random variables satisfying the following property: For any $\alpha, \rho > 0$, there exist $h \in]0, \alpha[$ and \mathcal{F}_{t+h} -random variables a^h and b^h such that

$$\begin{cases} i) & \mathbb{E}(\|a^h\|^2) \leq \rho^2 \\ ii) & \mathbb{E}(\|b^h\|^2) \leq \rho^2 \\ iii) & \mathbb{E}(b^h) = 0 \\ iv) & b^h \text{ is independent of } \mathcal{F}_t \end{cases} \quad (2.1)$$

and satisfying

$$x + v(W(t+h) - W(t)) + h\gamma + ha^h + \sqrt{h}b^h \in K \quad (2.2)$$

We consider the stochastic differential equation

$$d\xi = f(\xi(t))dt + g(\xi(t))dW(t) \quad (2.3)$$

where f and g are Lipschitz.

We say that a stochastic process $\xi(t)$ defined by

$$\xi(t) = \xi(0) + \int_0^t f(\xi(s))ds + \int_0^t g(\xi(s))dW(s) \quad (2.4)$$

is a solution to the stochastic differential equation (2.3) if the functions f and g satisfy:

for almost all $\omega \in \Omega$, $f(\xi(\cdot)) \in L^1(0, T; X)$ & $g(\xi(\cdot)) \in L^2(0, T; X)$

Definition 2.2 *We shall say that a stochastic process $x(\cdot)$ is viable in K if and only if*

$$\forall t \in [0, T], \quad x(t) \in K \quad (2.5)$$

i.e., if and only if

$$\forall t \in [0, T], \text{ for almost all } \omega \in \Omega, \quad \xi_\omega(t) \in K_\omega$$

We shall say that K enjoys the (stochastic) viability property with respect to the pair (f, g) if for any random variable x in K , there exists a solution ξ to the stochastic differential equation starting at x which is viable in K .

3 STOCHASTIC VIABILITY

Theorem 3.1 (Stochastic Viability) *Let K be a closed subset of X . We assume that either*

- *the maps f and g are Lipschitz*
- *the maps f and g are uniformly continuous and monotone in the sense that there exists $\nu \in \mathbf{R}$ such that*

$$\forall x, y \in X, \quad 2\langle f(x) - f(y), x - y \rangle + \|g(x) - g(y)\|^2 \leq \nu \|x - y\|^2,$$

Then the following conditions are equivalent:

1. — *From any initial stochastic process $\xi_0 \in \mathcal{K}$ starts a solution to the stochastic differential equation which is viable in \mathcal{K} .*
2. — *for every \mathcal{F}_t -random variable x in \mathcal{K} ,*

$$(f(x), g(x)) \in \mathcal{T}_{\mathcal{K}}(t, x) \quad (3.1)$$

We already proved in [2, Aubin & Da Prato] that the condition was necessary. It remains to prove that it is sufficient.

We begin by constructing approximate viable solutions to the stochastic differential equation.

Lemma 3.2 *Let K be a closed subset of X . We assume that the maps f and g are uniformly continuous. Then, for any $\varepsilon > 0$, the set $\mathcal{S}_\varepsilon(\xi_0)$ of stochastic processes $\xi(\cdot)$ on $[0, 1]$ satisfying $\xi(0) = \xi_0$ and*

$$\begin{cases} i) & \forall t \in [0, 1], \quad \mathbf{E}(\mathbf{d}^2(\xi(t), \mathcal{K})) \leq \varepsilon^2 \\ ii) & \forall t \in [0, 1], \quad \mathbf{E} \left\| \xi(t) - \xi(0) - \int_0^t f(\xi(s)) ds - \int_0^t g(\xi(s)) dW(s) \right\|^2 \leq \varepsilon^2 \end{cases} \quad (3.2)$$

is not empty.

Proof — Let us fix $\varepsilon > 0$. Since f and g are uniformly continuous with concave uniform continuity modulus¹ ω , we choose $\eta \in]0, \varepsilon]$ such that

$$\omega(\eta^2) \leq \frac{\varepsilon^2}{4}.$$

¹Set $\omega(t) = \sup_{\|x-y\|^2 \leq t} \|f(x) - f(y)\|^2$. Then ω is a non decreasing, subadditive continuity modulus of f . One can check that the concave envelope of ω is still a uniform continuity.

We denote by $\mathcal{A}_\varepsilon(\xi_0)$ the set of pairs $(T_\xi, \xi(\cdot))$ where $T_\xi \in [0, 1]$ and $\xi(\cdot)$ is a stochastic process satisfying $\xi(0) = \xi_0$ and

$$\begin{cases} i) & \forall t \in [0, T_\xi], \mathbf{E} d^2(\xi(T_\xi), \mathcal{K}) \leq \eta^2 T_\xi \\ ii) & \forall t \in [0, T_\xi], \mathbf{E} d^2(\xi(t), \mathcal{K}) \leq \eta^2 \\ iii) & \forall t \in [0, T_\xi], \mathbf{E} \left\| \xi(t) - \xi(0) - \int_0^t f(\xi(s)) ds - \int_0^t g(\xi(s)) dW(s) \right\|^2 \leq \varepsilon^2 \end{cases} \quad (3.3)$$

The set $\mathcal{A}_\varepsilon(\xi)$ is not empty: take $T_\xi = 0$ and $\xi(0) \equiv \xi_0$. It is an inductive set for the order relation

$$(T_{\xi_1}, \xi_1(\cdot)) \preceq (T_{\xi_2}, \xi_2(\cdot))$$

if and only if

$$T_{\xi_1} \leq T_{\xi_2} \text{ \& } \xi_2(\cdot)|_{[0, T_{\xi_1}]} = \xi_1(\cdot)$$

Zorn's Lemma implies that there exists a maximal element $(T_\xi, \xi(\cdot)) \in \mathcal{A}_\varepsilon(\xi_0)$. The Lemma follows from the claim that for such a maximal element, we have $T_\xi = 1$. If not, we shall extend $\xi(\cdot)$ by a stochastic process $\hat{\xi}(\cdot)$ on an interval $[T_\xi, S_\xi]$ where $S_\xi > T_\xi$, contradicting the maximal character of $(T_\xi, \xi(\cdot))$.

Since K_ω and $\xi_\omega(T_\xi)$ are \mathcal{F}_{T_ξ} measurable, the projection map $\Pi_{K_\omega}(\xi_\omega(T_\xi))$ is also \mathcal{F}_{T_ξ} -measurable (see [3, Theorem 8.2.13, p. 317]). Then there exists a \mathcal{F}_{T_ξ} -measurable selection $y_\omega \in \Pi_{K_\omega}(\xi_\omega(T_\xi))$, which we call a projection of the random variable $\xi(T_\xi)$ onto the random set-valued variable \mathcal{K} . For simplicity, we set $x = \xi(T_\xi)$ and thus choose a projection $y \in \Pi_{\mathcal{K}}(x)$.

We take

$$\rho := \frac{\eta \sqrt{1 - T_\xi}}{2} > 0$$

and we set

$$c^2 := \max(\mathbf{E}(\|f(y)\|^2), \mathbf{E}(\|g(y)\|^2)) < +\infty \quad (3.4)$$

We then introduce

$$\alpha := \min \left(\eta, \frac{(1 - T_\xi)\eta^2}{\eta^2 + 4c^2} \right) > 0$$

which is positive whenever $T_\xi < 1$.

We know that $(f(y), g(y))$ belongs to the stochastic contingent set $\mathcal{T}_{\mathcal{K}}(T_x, y)$: There exist $h_x \in]0, \alpha]$ and $\mathcal{F}_{T_x+h_x}$ -random variables a^{h_x} and b^{h_x} such that

$$\begin{cases} i) & \mathbf{E}(\|a^{h_x}\|^2) \leq \rho^2 \\ ii) & \mathbf{E}(\|b^{h_x}\|^2) \leq \rho^2 \\ iii) & \mathbf{E}(b^{h_x}) = 0 \\ iv) & b^{h_x} \text{ is independent of } \mathcal{F}_t \end{cases} \quad (3.5)$$

and satisfying

$$y + g(y)(W(T_x + h_x) - W(T_x)) + h_x f(y) + h_x a^h + \sqrt{h_x} b^{h_x} \in \mathcal{K} \quad (3.6)$$

We then set $S_x := T_x + h_x > T_x$ and we define the stochastic process $\hat{\xi}(t)$ on the interval $[T_x, S_x]$ by

$$\hat{\xi}(t) := x + (t - T_x)f(y) + (W(t) - W(T_x))g(y)$$

Therefore, setting $h := t - T_x$,

$$d_{\mathcal{K}}^2(\hat{\xi}(t)) - d_{\mathcal{K}}^2(\hat{\xi}(T_x)) \leq \|x - y - ha^h - \sqrt{h}b^h\|^2 - \|x - y\|^2 =:$$

$$\|ha^h + \sqrt{h}b^h\|^2 - 2\langle x - y, ha^h \rangle - 2\langle x - y, \sqrt{h}b^h \rangle$$

We take the expectation in both sides of this inequality and estimate each term of the right hand-side. First, we use estimate

$$\mathbf{E}(\|ha^h + \sqrt{h}b^h\|^2) \leq 2h(h\mathbf{E}(\|a^h\|^2) + \mathbf{E}(\|b^h\|^2))$$

because

$$\mathbf{E} \left(\left\| \int_0^t \varphi(s) ds \right\|^2 \right) \leq t \int_0^t \mathbf{E}(\|\varphi(s)\|^2) ds$$

and

$$\mathbf{E} \left(\left\| \int_0^t \varphi(s) dW(s) \right\|^2 \right) = \int_0^t \mathbf{E}(\|\varphi(s)\|^2) ds$$

Next,

$$\mathbf{E} \langle x - y, a^h \rangle \leq \mathbf{E} (\|x - y\|^2)^{\frac{1}{2}} \left(\mathbf{E} (\|a^h\|^2) \right)^{\frac{1}{2}}$$

and we observe that

$$\mathbf{E} \left\langle x - y, \frac{1}{\sqrt{h}} b^h \right\rangle = 0$$

since b^h is independent of $x - y$ and $\mathbf{E}(b^h) = 0$.

We obtain, by the very choice of ρ ,

$$\begin{aligned} \mathbf{E}(d^2(\hat{\xi}(S_x), \mathcal{K})) &= \mathbf{E}(d^2(\hat{\xi}(T_x + h_x), \mathcal{K})) \\ &\leq \mathbf{E}d^2(\hat{\xi}(T_x), \mathcal{K}) + 2h_x \mathbf{E} (\|x - y\|^2)^{\frac{1}{2}} \left(\mathbf{E} (\|a^{h_x}\|^2) \right)^{\frac{1}{2}} \\ &\quad + 2h_x (h_x \mathbf{E}(\|a^{h_x}\|^2) + \mathbf{E}(\|b^{h_x}\|^2)) \end{aligned}$$

$$\begin{aligned}
&\leq \mathbf{E} \mathbf{d}^2(\widehat{\xi}(T_x), \mathcal{K}) + h_x \left(\mathbf{E}(\|x - y\|^2) + 3\mathbf{E}(\|a^{h_x}\|^2) + \mathbf{E}(\|b^{h_x}\|^2) \right) \\
&\leq \eta^2 T_x + h_x(\eta^2 T_x + 4\rho^2) \leq \eta^2 T_x + h_x \eta^2 = \eta^2 S_x
\end{aligned}$$

by (3.3)i).

Hence $\widehat{\xi}(\cdot)$ satisfies (3.3)i) for S_x .

We observe also that for any $t \in [T_x, S_x]$,

$$\mathbf{d}_{\mathcal{K}}^2(\widehat{\xi}(t)) \leq \|\widehat{\xi}(t) - y\|^2$$

and that

$$\begin{aligned}
\|\widehat{\xi}(t) - y\|^2 &= \|x - y + (t - T_x)f(y) + (W(t) - W(T_x))g(y)\|^2 \\
&= \mathbf{d}_{\mathcal{K}}^2(x) + 2\langle x - y, (t - T_x)f(y) + (W(t) - W(T_x))g(y) \rangle \\
&\quad + \|(t - T_x)f(y) + (W(t) - W(T_x))g(y)\|^2
\end{aligned}$$

By taking the expectations, we obtain

$$\begin{aligned}
&\mathbf{E}(\|\widehat{\xi}(t) - y\|^2) - \mathbf{E}(\mathbf{d}_{\mathcal{K}}^2(\widehat{\xi}(T_x))) \\
&\leq (t - T_x)(\mathbf{E}(\mathbf{d}_{\mathcal{K}}^2(\widehat{\xi}(T_x))) + (1 + 2(t - T_x))\mathbf{E}(\|f(y)\|^2) + \mathbf{E}(\|g(y)\|^2))
\end{aligned}$$

Therefore, since $\max(\mathbf{E}(\|f(y)\|^2), \mathbf{E}(\|g(y)\|^2)) = c^2$ by (3.4), we deduce that

$$\mathbf{E}\|\widehat{\xi}(t) - y\|^2 \leq \eta^2 T_x + (t - T_x)(\eta^2 T_x + 4c^2) \leq \eta^2 T_x + \alpha(\eta^2 + 4c^2) \leq \eta^2 \quad (3.7)$$

since, by the choice of α , we have $\alpha(\eta^2 + 4c^2) \leq (1 - T_x)\eta^2$. Therefore,

$$\mathbf{E}(\mathbf{d}_{\mathcal{K}}^2(\widehat{\xi}(t))) \leq \mathbf{E}(\|\widehat{\xi}(t) - y\|^2) \leq \eta^2$$

Hence $\widehat{\xi}(\cdot)$ satisfies (3.3)ii) for S_x .

We also observe that

$$\begin{aligned}
&\mathbf{E} \left(\left\| \widehat{\xi}(t) - x - \int_{T_x}^t f(\widehat{\xi}(s))ds - \int_0^t g(\widehat{\xi}(s))dW(s) \right\|^2 \right) \\
&= \mathbf{E} \left(\left\| \int_{T_x}^t (f(y) - f(\widehat{\xi}(s)))ds + \int_{T_x}^t (g(y) - g(\widehat{\xi}(s)))ds \right\|^2 \right) \\
&\leq 2 \left(\mathbf{E} \left(\int_{T_x}^t \|f(y) - f(\widehat{\xi}(s))\|^2 ds \right) + \mathbf{E} \left(\int_{T_x}^t \|g(y) - g(\widehat{\xi}(s))\|^2 ds \right) \right)
\end{aligned}$$

Since the functions f and g are uniformly continuous, we deduce from the concavity of the continuous modulus $\omega(\cdot)$ that

$$\begin{aligned} & \mathbf{E} \left(\left\| \widehat{\xi}(t) - x - \int_{T_x}^t f(\widehat{\xi}(s)) ds - \int_{T_x}^t g(\widehat{\xi}(s)) dW(s) \right\|^2 \right) \\ & \leq 2 \left(\mathbf{E} \left(\int_{T_x}^t \omega \left(\|y - \widehat{\xi}(s)\|^2 \right) ds \right) + \mathbf{E} \left(\int_{T_x}^t \omega \left(\|y - \widehat{\xi}(s)\|^2 \right) ds \right) \right) \\ & \leq 4 \mathbf{E} \left(\int_{T_x}^t \omega \left(\|y - \widehat{\xi}(s)\|^2 \right) ds \right) \leq 4 \omega \left(\int_{T_x}^t \mathbf{E} \left(\|y - \widehat{\xi}(s)\|^2 \right) ds \right) \\ & \leq 4 \omega(\eta^2) \leq \varepsilon^2. \end{aligned}$$

since we have already proved that

$$\mathbf{E}(\|\widehat{\xi}(t) - y\|^2) \leq \eta^2$$

so that $\widehat{\xi}(\cdot)$ satisfies (3.3)iii). Therefore, we have extended the maximal solution $(T_{\xi}, \xi(\cdot))$ on the interval $[0, S_x]$ and obtained the desired contradiction. Hence the proof of Lemma 3.2 is completed. ■

It remains now to prove that the limit of the sequence of approximate solutions to a viable stochastic process exists and is a solution to the stochastic differential equation.

Let us choose for every ε an approximate solution ξ_ε which can be written in the form

$$\xi_\varepsilon(t) = \xi_0 + \int_0^t f(\xi_\varepsilon(s)) ds + \int_0^t g(\xi_\varepsilon(s)) dW(z) + \zeta_\varepsilon(t)$$

where $\sup_{t \in [0,1]} \mathbf{E}(\|\zeta_\varepsilon(t)\|^2) \leq \varepsilon^2$. Then for any $\varepsilon, \eta > 0$,

$$\mathbf{E} \|\xi_\varepsilon(t) - \xi_\eta(t)\|^2 \leq$$

$$\mathbf{E} \left\| \int_0^t (f(\xi_\varepsilon(s)) - f(\xi_\eta(s))) ds + \int_0^t (g(\xi_\varepsilon(s)) - g(\xi_\eta(s))) dW(z) + \zeta_\varepsilon(t) - \zeta_\eta(t) \right\|^2$$

It follows that

Lipschitz case

$$\mathbf{E} \left(\|\xi_\varepsilon(t) - \xi_\eta(t)\|^2 \right) \leq 4l^2 \left(\int_0^t \mathbf{E} \left(\|\xi_\varepsilon(s) - \xi_\eta(s)\|^2 \right) ds \right) + 2(\varepsilon^2 + \eta^2)$$

Gronwall's Lemma implies that

$$\mathbf{E} \left(\|\xi_\varepsilon(t) - \xi_\eta(t)\|^2 \right) \leq 2(\varepsilon^2 + \eta^2)e^{4l^2t}$$

Monotone case We use Ito formula for the function $\|\xi_\varepsilon - \xi_\eta\|^2$ to obtain

$$\begin{aligned} & \mathbf{E} \|\xi_\varepsilon(t) - \xi_\eta(t)\|^2 \\ & \leq \mathbf{E} \int_0^t (\langle \xi_\varepsilon(s) - \xi_\eta(s), f(\xi_\varepsilon(s)) - f(\xi_\eta(s)) \rangle + \|g(\xi_\varepsilon(s)) - g(\xi_\eta(s))\|^2) ds \\ & \quad + 2(\varepsilon^2 + \eta^2) \\ & \leq \nu^2 \int_0^t \mathbf{E}(\|\xi_\varepsilon(s) - \xi_\eta(s)\|^2) ds + 2(\varepsilon^2 + \eta^2) \end{aligned}$$

Gronwall's Lemma implies that

$$\mathbf{E} \|\xi_\varepsilon(t) - \xi_\eta(t)\|^2 \leq 2(\varepsilon^2 + \eta^2)e^{\nu^2 t}$$

In both cases we deduce that the above Cauchy sequences converge to some $\xi(\cdot)$:

$$\forall t \in [0, 1], \quad \lim_{\varepsilon \rightarrow 0} \mathbf{E}(\|\xi_\varepsilon(t) - \xi(t)\|^2) = 0$$

Furthermore, inequalities (3.2)ii) imply that

$$\mathbf{E} \left(d_{\mathcal{K}}^2(\xi(t)) \right) = 0$$

so that the solution is viable in \mathcal{K} . ■.

References

- [1] AUBIN J.-P. & CELLINA A. (1984) DIFFERENTIAL INCLUSIONS, Springer-Verlag
- [2] AUBIN J.-P. & DA PRATO G. (1990) *Stochastic Viability and Invariance*, Annali Scuola Normale di Pisa, 27, 595-694
- [3] AUBIN J.-P. & FRANKOWSKA H. (1990) SET-VALUED ANALYSIS, Birkhäuser, Boston, Basel
- [4] AUBIN J.-P. (1991) VIABILITY THEORY Birkhäuser, Boston, Basel
- [5] BENSOUSSAN A. (1982) STOCHASTIC CONTROL BY FUNCTIONAL ANALYSIS METHODS, North-Holland
- [6] BENSOUSSAN A. & LIONS J.-L. (1978) APPLICATIONS DES INÉQUATIONS VARIATIONNELLES EN CONTRÔLE STOCHASTIQUE, Dunod

- [7] DA PRATO G. & FRANKOWSKA H. (preprint) *A Stochastic Filippov Theorem*
- [8] DA PRATO G. & ZABCZYK J. (1992) *STOCHASTIC EVOLUTION EQUATIONS IN INFINITE DIMENSIONS* Cambridge University Press
- [9] MILIAN A. (preprint) *Characterization of coefficients of Ito equation whose solutions live in some subsets of \mathbb{R}^m .*