## SERUM BILIRUBIN(TSB)

**NORMAL RANGE UNITS** Adults: 0.4 - 1.2 Total Bilirubin 16.9 mg/dl (Method: Modified Jendrassik) Children: F١ Name : Baby of. SWATHI Bill Number :0-**M739**: 2.0 - 6.0 Age/Gender : 25DAYS/MALE Bill Date :1-25daps:2084 18:67 PM Sample Type : SERUM Sample Collection 3-35dAps-2084 82048 PM Reff By : C/O SRADDHA HOSPITAL Sample Received PREMATERAL 12:48 PM TypedBy : Bharat Saini Reporting Date C254APT02024001:38 PM 0-1 day: 1.0 - 8.0 1-2 days: 6.0 - 12.0 3-5 days: 10.0 - 14.0 1.0 Up to 0.25 Direct Bilirubin mg/dl

Elevation in serum unconjugated bilirubin levels occur in haemolytic jaundice due to excessive haemolysis.B). Hepatic jaundice is associated with increase in both conjugated and unconjugated bilirubin in serum. Ref for BRI: Carl.A.Burtis, David.E.Burn et.al Tietz Fundamentals of Clinical Chemistry and Molecular Diagnostics. 7th ed., page No. 955

15.9

-----End of the Report-----

mg/dl

up to 1

AR INCHARGE

Authorized Signatory

(Method: Modified Jendrassik)

Indirect Bilirubin

(Method: Calculated)

, Opp. Bajaj Electronics, Beside Govt. Hospital, New Market Road, Ambedkar TANCHERU, Sangareddy Dist- 502 319. T.S. 🛍 : 08455 296155, 🙃: 960349617