

## Use of bone and otolith measures for size-estimation of fish in predator-prey studies

Ali Serhan TARKAN<sup>1</sup>\*, Çiğdem GÜRSOY GAYGUSUZ<sup>2</sup>, Özcan GAYGUSUZ<sup>1</sup> and Hasan ACIPINAR<sup>3</sup>

<sup>1</sup> *Istanbul University, Faculty of Fisheries, Ordu Cad. No. 200, 34470 Laleli, Istanbul, Turkey;*  
\*e-mail: serhantarkan@yahoo.com, serhan@istanbul.edu.tr

<sup>2</sup> *Çanakkale Onsekiz Mart University, Faculty of Fisheries, 17020 Çanakkale, Turkey*

<sup>3</sup> *Istanbul University, Natural and Applied Sciences Institute, 34850 Avcılar, Istanbul, Turkey*

Received 15 June 2007; Accepted 1 October 2007

**Abstract.** To estimate the size of fish taken as prey by piscivorous predators, linear or non-linear relationships between bone measures (pharyngeal, opercula, cleithra, anal and dorsal spine bones, otoliths) and body length were elaborated for eleven Eurasian cyprinid fish species captured in three lakes of Turkey: rudd *Scardinius erythrophthalmus*, Baltic vimba *Vimba vimba*, Danube bleak *Chalcalburnus chalcoides*, gibel carp *Carassius gibelio*, roach *Rutilus rutilus*, silver bream *Blicca bjoerkna*, common carp *Cyprinus carpio*, chub *Leuciscus cephalus*, Dnieper chub *Petroleuciscus borythenicus*, tench *Tinca tinca* and tarek *Alburnus tarichi* (endemic species for Lake Van). All calculated regressions were highly significant, with coefficients of determination >81% in most of cases. The results suggest that the biometric relationships between fish length and some bones (pharyngeal, opercula, cleithra) are well suited for use in prey-predator studies of all the studied species, but otoliths and the dorsal and anal spines can be used for some fish species only (rudd, Baltic vimba, roach, silver bream, gibel carp).

**Key words:** diet analysis; prey size estimation; piscivorous predators; cyprinids

## Introduction

The determination of prey identity and size is a crucial aspect in dietary studies of piscivorous animals (e.g. Mann & Beaumont 1980, Hansel et al. 1988, Pierce & Boyle 1991). In Eurasian fresh waters, cyprinids are often the predominant taxonomic family in the fish assemblages and consequently knowledge of their bone and body biometry is useful in the dietary studies of the piscivorous animals that feed on them (e.g. Prenda & Granado-Lorencio 1992, Radke et al. 2000, Beyer et al. 2006). Although there have been several studies that have used freshwater fish bones to estimate the original lengths and weights of prey consumed by predators (e.g. Mann & Beaumont 1980, Prenda & Granado-Lorencio 1992, Copp & Kováč 2003, Miranda & Escala 2005), there are no known studies for freshwater fishes that have examined the relationships between otolith size and fish length, including different measurements of both the left and right sides. Similarly, few studies have attempted to increase the precision and accuracy of the back-calculated lengths by comparing more than one measurement with both sides of any given bone structures (Radke et al. 2000, Hájková et al. 2003). The aim of the present study was to examine these prerequisites for diet analysis. We present the equations for estimating the original lengths and weights of the fishes consumed, from both sides' measurements of diagnostic bones (pharyngeal, opercula, cleithra, dorsal and anal spine and otoliths) of eleven Eurasian cyprinid fish species for which relatively little

\* Corresponding author

or no biometric information exists. These bones were chosen because of their taxonomic importance (Miranda & Escala 2005), usage in diet analyses of piscivorous animals (e.g. Copp & Roche 2003, Britton & Shepherd 2005) and in paleontological studies (e.g. Yasuno 1997).

## Material and Methods

A total of 2256 specimens, representing eleven species, were collected from January 2002 to June 2003 from two water bodies (Ömerli Reservoir and Lake Sapanca) in northwest Turkey and from one lake (Lake Van) in eastern Turkey using various fishing methods (gill-netting, beach-seine netting, fyke netting, electrofishing). The fish were measured for total length (TL) to the nearest mm and weighed to the nearest 0.1 g (Table 1). The otoliths were then extracted and the fish were placed in boiling water (duration depended on fish size) to permit recovery of the pharyngeal bone, opercula, cleithra, dorsal and anal spines. To examine the relationships with fish size, replicate measurements were taken (to the nearest 0.01 mm) of the bones and otoliths using a digital calliper or a micrometer under a binocular microscope from both the left and right sides: five measurements of the pharyngeal bone, six measurements of the opercula, four measurements of the cleithra, three measurements of the otolith and one measurement of the anal and dorsal spines (Fig. 1). Otolith pairs of the lapillus were used because of their relatively bigger size and regular shape as compared to other otolith pairs in cyprinids. However, some species have fragile and small otoliths, which are likely to be broken or strongly eroded in diet samples (e.g. tarek *Alburnus tarichi*, Danube bleak *Chalcalburnus chalcoides*, common carp *Cyprinus carpio*, chub *Leuciscus cephalus*, Dnieper chub *Petroleuciscus borysthenicus* and tench *Tinca tinca*). Therefore, their otolith size vs. body size relationships was not calculated. Linear ( $y = ax + b$ ) and non-linear ( $y = ax^b$ , power model) regression equations and analysis of variance (ANOVA), where  $y = TL$ , were fitted to determine what equations best described the relationships between fish size and bone/otolith dimensions. Relationships with the highest coefficient of determination ( $r^2$ ) were

**Table 1.** The species, common names, locations, sample sizes and total length (TL) ranges of fish used in the study.

| Species                             | Common name   | Location         | n   | TL range |
|-------------------------------------|---------------|------------------|-----|----------|
| <i>Alburnus tarichi</i>             | Tarek         | Lake Van         | 63  | 191–246  |
| <i>Blicca bjoerkna</i>              | Silver bream  | Lake Sapanca     | 157 | 120–212  |
| <i>Carassius gibelio</i>            | Gibel carp    | Ömerli Reservoir | 229 | 86–357   |
| <i>Chalcalburnus chalcoides</i>     | Danube bleak  | Ömerli Reservoir | 36  | 115–284  |
| "                                   | "             | Lake Sapanca     | 24  | 180–233  |
| <i>Cyprinus carpio</i>              | Common carp   | Ömerli Reservoir | 7   | 222–321  |
| "                                   | "             | Lake Sapanca     | 8   | 168–224  |
| <i>Leuciscus cephalus</i>           | European chub | Ömerli Reservoir | 12  | 97–257   |
| <i>Petroleuciscus borysthenicus</i> | Dnieper chub  | Ömerli Reservoir | 34  | 98–149   |
| <i>Rutilus rutilus</i>              | Roach         | Lake Sapanca     | 563 | 141–381  |
| <i>Scardinius erythrophthalmus</i>  | Rudd          | Ömerli Reservoir | 509 | 93–270   |
| "                                   | "             | Lake Sapanca     | 246 | 85–277   |
| <i>Tinca tinca</i>                  | Tench         | Lake Sapanca     | 33  | 135–278  |
| <i>Vimba vimba</i>                  | Baltic vimba  | Ömerli Reservoir | 293 | 99–294   |
| "                                   | "             | Lake Sapanca     | 42  | 174–262  |

adopted as the best predictor (Z a r 1999). To test whether left and right side provided similar results, the linear and non-linear regression were calculated for all measures respectively, and the maximum relative deviation between the predicted regression lines was determined. Data of left and right sides of a measure were pooled for further analysis only if maximum relative difference within the range measured was <2.5% (R a d k e et al. 2000). The measures providing the most accurate estimates of the back-calculated lengths were determined by calculating the confidence limits (95%) and then comparing the maximum values of relative error (confidence limit/calculated length) between measures (R a d k e et al. 2000). Analysis of covariance, ANCOVA (Z a r 1999), was used to analyze effect of site on bone and otolith measures of same species (Danube bleak, common carp, rudd *Scardinius erythrophthalmus* and Baltic vimba *Vimba vimba*) occurring in two different locations studied.




Fig. 1. The bones and otoliths of the fish showing the measures used in the present study.

Results

The relationships between fish TL and bone measurements did not differ between Ömerli Reservoir and Lake Sapanca within the same species (ANCOVA,  $P > 0.005$ ), so data from these locations were combined. The left and right side measurements of the pharyngeal bone were pooled in 69% of all cases. Measurements of tooth width of pharyngeal bone (*ce*) were analyzed separately in all cases except tench and common carp (Appendix 2 and 4). Operculum and cleithrum data from left and right sides were not pooled only for two species (roach *Rutilus rutilus* and silver bream *Blicca bjoerkna*) (Appendices 1 and 3). Non-linear and linear functions provided the best fit for 57% and 43% of all regressions, respectively. All regressions were highly significant ( $P < 0.001$ ) and analysis of bone morphometric parameters vs. TL showed that the regression model explained more than 81% of the variance in all species, with exception of silver bream and tarek in some cases (Appendix 1). The maximum

values of relative error ranged from 1.2 to 27.6% in pharyngeal, from 1.3 to 24.9% in opercula, from 1.2 to 22.1% in cleithra bones, from 1.3 to 7.0% in otolith, from 5.0 to 13.6% in dorsal spine and from 3.0 to 13.7% in anal spine. Chub was found to give considerably high maximum values of relative error while tarek, silver bream and roach gave low maximum values of relative error for all structures measured. Anal and dorsal spine percent errors were lower for gibel carp *Carassius gibelio* than those in common carp (Appendices 1–4).

## Discussion

Linear functions were usually adequate to describe bone size – fish length relationships (Mann & Beaumont 1980, Hansel et al. 1988, Prenda & Granado-Lorenzo 1992), but some authors (Newsome 1977, Radke et al. 2000) found that curvilinear relationships provided the best fit for some fish as seen in the present study. Left and right measurements of some bone structures do not always provide the same estimate of prey fish length (see Raczisky & Szuba 1997), so the pooling left and right measurements should be undertaken after adequate statistical analysis only (e.g. Radke et al. 2000, Copp & Kováč 2003, Hájková et al. 2003). The data for cleithra, opercula, dorsal and anal spine and otolith (Appendix 1–4) could not be compared with those from the literature due to differences in the models and lengths used. However, the observed relationships between pharyngeal bone length and TL for silver bream, gibel carp, common carp, rudd and tench (Appendix 1–4) are similar to those reported by Radke et al. (2000). This suggests that the relationship between bone size and body length are relatively constant within species across geographical ranges (e.g. Copp & Kováč 2003). However, in the present study the slope and intercept values within Cyprinidae vary greatly, reflecting natural variability in ontogenesis between the species compared. The onset of skeleton formation during early development of fish is species specific, and also strongly depends on environmental conditions, especially temperature and oxygen concentrations (e.g. Balon 1981). Both bone measures and otoliths are exposed to a variable degree of chemical and mechanical abrasion in the digestive track of predators. Hence, some small otoliths and bones are likely to be totally dissolved and eroded. Furthermore, partial digestion will bias estimates of prey size (Jobling & Breiby 1986, Pierce & Boyle 1991). Use of hard structures may also bias data on food habits by favoring larger over smaller prey fish because their bones may be more resistant to digestion (Hansel et al. 1988). Generally, the bones are relatively resistant to digestion and occasionally they are the only recognizable remains (Mann & Beaumont 1980). Similarly, *in vitro* (Pierce et al. 1993) and *in vivo* (Pierce et al. 1993, Carss & Elston 1996) experiments provided some evidence of a higher resistance of bones to acidic digestion in relation to otoliths, which represents an advantage of the former to back-calculate prey size. Even though otoliths showed relatively low percent errors compared to other structures in the present study (Appendix 1–4), only five of eleven fish species could be suitable for otolith analyses. Fish weight can also be estimated by two-step procedures, first using a relationship between structure length and fish length and then applying a fish length/fish weight equation. Length-weight relationships of fishes studied in the present study were given in Tarkan et al. (2006). Our results suggested that the biometric relationship between measurements of pharyngeal, opercula, cleithra bones and TL are well suited to the prediction of prey fish lengths from the partly digested remains for all fish species studied. However, dorsal and anal spine bones and otoliths may be used for some fish species only (rudd, Baltic vimba, roach, silver bream, gibel carp). The outputs of this

study provide a tool for biometric relationships that enable estimation of length and weight using the bones. This information should facilitate the assessment of the diet of piscivorous fauna in Turkey, but potentially wherever these eleven fish species occur.

## Acknowledgements

We thank G.H. Copp, J. Cucherousset and R.J. Radke for editorial and scientific comments on the manuscript as well as B. Öztürk, N. Gökalp and A. Sümer for facilitating this study, which are greatly appreciated.

## LITERATURE

- Balon E.K. 1981: Saltatory processes and altricial to precocial forms in the ontogeny of fishes. *American Zoologist* 21: 573–596.
- Beyer K., Miranda R., Copp G.H. & Gozlan R.E. 2006: Biometric data and bone identification of topmouth gudgeon, *Pseudorasbora parva* and sunbleak, *Leucaspis delineatus*. *Folia Zool.* 55: 287–292.
- Britton J. R. & Shepherd J. S. 2005: Biometric data to facilitate the diet reconstruction of piscivorous fauna. *Folia Zool.* 54: 193–200.
- Carss D. & Elston D. 1996: Errors associated with otter *Lutra lutra* faecal analysis. II. Estimating prey size distribution from bones recovered in spraints. *J. Zool. (London)* 238: 319–332.
- Copp G.H. & Kováč V. 2003: Biometric relationships between body size and bone lengths in fish prey of the Eurasian otter *Lutra lutra*: chub *Leuciscus cephalus* and perch *Perca fluviatilis*. *Folia Zool.* 52: 109–112.
- Copp G.H. & Roche K. 2003: Range and diet of Eurasian otters *Lutra lutra* (L.) in the catchment of the river Lee (south-east England) since re-introduction. *Aquat. Conserv.: Mar. Freshwat. Ecosyst.* 13: 65–76.
- Hájková P., Roche K. & Kocián L'. 2003: On the use of diagnostic bones of brown trout, *Salmo trutta* m. *fario*, grayling, *Thymallus thymallus* and Carpathian sculpin, *Cottus poecilopus* in Eurasian otter, *Lutra lutra* diet analysis. *Folia Zool.* 52: 389–398.
- Jobling M. & Breiby A. 1986: The use and abuse of fish otoliths in studies of feeding habits of marine piscivores. *Sarsia* 71: 265–274.
- Hansel H.C., Duke S.D., Lofy P.T. & Gray G.A. 1988: Use of diagnostic bones to identify and estimate original lengths of ingested prey fishes. *Trans. Amer. Fish. Soc.* 117: 55–62.
- Mann R.H.K. & Beaumont W.R.C. 1980: The collection, identification and reconstruction of lengths of fish prey from their remains in pike stomachs. *Fish. Manag.* 11: 169–172.
- Miranda R. & Escala M. C. 2005: Morphometrical comparison of cleithra, opercular and pharyngeal bones of autochthonous Leuciscinae (Cyprinidae) of Spain. *Folia Zool.* 54: 173–188.
- Newsome G.E. 1977: Use of opercular bones to identify and estimate lengths of prey consumed by piscivores. *Can. J. Zool.* 55: 733–736.
- Pierce G.J. & Boyle P.R. 1991: A review of methods for diet analysis in piscivorous marine mammals. *Oceanogr. Mar. Biol.* 2: 409–486.
- Pierce G.J., Boyle P.R., Watt J. & Grisley M. 1993: Recent advances in diet analysis of marine mammals. *Symp. Zool. Soc.* 66: 214–261.
- Prenda J. & Granado-Lorencio C. 1992: Biometric analysis of some cyprinid bones of prey fishes to estimate original lengths and weights. *Folia Zool.* 14: 175–185.
- Raczynski M. & Szuba Z. 1997: Description of the inferior pharyngeal bones, ossa pharynges inferiora, in the common bream *Abramis brama* (L.) from the Szczecin lagoon with special regard to bilateral asymmetry. *Acta Ichthyol. Pisc.* 27: 59–67.
- Radke R.J., Petzoldt T. & Wolter C. 2000: Suitability of pharyngeal bone measures commonly used for reconstruction of prey fish length. *J. Fish Biol.* 57: 961–967.
- Tarkan A.S., Gaygusuz Ö., Acıpinar H., Gürsoy Ç. & Özüluğ M. 2006: Length-weight relationship of fishes from the Marmara region (NW-Turkey). *J. Appl. Ichthyol.* 22: 271–273.
- Yasuno T. 1997: Fossil pharyngeal teeth of Cyprinid from the Late Pleistocene in Fukui Prefecture, central Japan. *Bull. Japan Sea Res. Inst.* 28: 65–75.
- Zar J.H. 1999: Biostatistical Analysis, 4<sup>th</sup> Edition. Prentice-Hall, Englewood Cliffs, NJ, 663 pp.

**Appendix 1.** Regression statistics for linear and non-linear relationships relating measurements (mm) of bones (pharyngeal, operculum, cleithrum, dorsal and anal spin) to total length and percent relative errors of each structure for *Alburnus tarichi*, *Blicca bjoerkna* and *Carassius gibelio*. See Figure 1 for measure descriptions. Linear:  $TL = ax + b$ ; non-Linear:  $TL = ax^b$ . Coefficient of determination ( $r^2$ ) and number of data pairs in regression (n) and are indicated.

| Species            | Structure  | Measure | Orientation | Type of regressions | a       | b       | $r^2$ | n   | Relative error (%) |
|--------------------|------------|---------|-------------|---------------------|---------|---------|-------|-----|--------------------|
| <i>A. tarichi</i>  | Pharyngeal | ab      | Pooled      | Linear              | 12.147  | 84.066  | 0.61  | 52  | 1.42               |
|                    |            | bd      | Left        | Linear              | 13.443  | 86.767  | 0.65  | 55  | 1.98               |
|                    |            | bd      | Right       | Linear              | 14.418  | 79.687  | 0.66  | 53  | 1.87               |
|                    |            | ce      | Left        | Linear              | 27.612  | 80.294  | 0.65  | 56  | 1.86               |
|                    |            | ce      | Right       | Linear              | 19.798  | 110.080 | 0.52  | 57  | 1.92               |
|                    |            | da      | Pooled      | Non-linear          | 63.053  | 0.620   | 0.62  | 54  | 1.33               |
|                    |            | bf      | Pooled      | Linear              | 11.884  | 71.968  | 0.72  | 52  | 1.40               |
|                    |            | a       | Pooled      | Linear              | 10.729  | 108.120 | 0.61  | 58  | 1.25               |
|                    |            | b       | Pooled      | Linear              | 6.308   | 103.120 | 0.73  | 59  | 1.26               |
|                    |            | c       | Pooled      | Non-linear          | 32.087  | 0.690   | 0.78  | 59  | 1.31               |
|                    | Operculum  | d       | Pooled      | Non-linear          | 39.736  | 0.591   | 0.74  | 58  | 1.29               |
|                    |            | e       | Pooled      | Non-linear          | 60.925  | 0.514   | 0.62  | 58  | 1.29               |
|                    |            | f       | Pooled      | Non-linear          | 52.300  | 0.512   | 0.75  | 59  | 1.28               |
|                    |            | a       | Pooled      | Non-linear          | 50.800  | 0.543   | 0.65  | 58  | 1.24               |
|                    |            | b       | Pooled      | Linear              | 6.848   | 82.572  | 0.69  | 59  | 1.28               |
|                    |            | c       | Pooled      | Linear              | 10.644  | 91.423  | 0.69  | 56  | 1.26               |
|                    | Cleithrum  | d       | Pooled      | Non-linear          | 25.296  | 0.675   | 0.73  | 56  | 1.28               |
|                    |            | ab      | Pooled      | Linear              | 15.978  | 3.195   | 0.82  | 121 | 2.02               |
|                    |            | bd      | Pooled      | Linear              | 20.967  | 17.716  | 0.77  | 121 | 2.00               |
|                    |            | ce      | Left        | Non-linear          | 47.078  | 0.815   | 0.67  | 124 | 2.83               |
| <i>B. bjoerkna</i> | Pharyngeal | ce      | Right       | Linear              | 27.213  | 23.190  | 0.66  | 119 | 3.00               |
|                    |            | da      | Pooled      | Linear              | 17.892  | 13.247  | 0.81  | 121 | 2.01               |
|                    |            | bf      | Pooled      | Linear              | 15.719  | 6.194   | 0.81  | 121 | 1.99               |
|                    | Operculum  | a       | Pooled      | Linear              | 23.373  | 20.642  | 0.80  | 57  | 2.80               |
|                    |            | b       | Left        | Linear              | 13.491  | 0.740   | 0.91  | 68  | 3.67               |
|                    |            | b       | Right       | Linear              | 13.674  | -3.619  | 0.91  | 67  | 3.70               |
|                    |            | c       | Pooled      | Linear              | 11.002  | 4.681   | 0.91  | 68  | 2.51               |
|                    |            | d       | Pooled      | Linear              | 10.772  | 0.155   | 0.92  | 65  | 2.72               |
|                    |            | e       | Pooled      | Non-linear          | 14.039  | 1.069   | 0.91  | 57  | 2.77               |
|                    |            | f       | Pooled      | Linear              | 14.119  | 7.679   | 0.89  | 65  | 2.48               |
|                    |            | a       | Pooled      | Linear              | 10.031  | 15.334  | 0.89  | 115 | 2.05               |
|                    | Cleithrum  | b       | Pooled      | Linear              | 10.831  | 10.654  | 0.89  | 127 | 1.96               |
|                    |            | c       | Pooled      | Linear              | 7.143   | 16.163  | 0.85  | 113 | 1.95               |
|                    |            | d       | Pooled      | Linear              | 17.880  | 24.482  | 0.82  | 120 | 2.01               |
|                    |            | a       | Pooled      | Linear              | 97.983  | -46.861 | 0.86  | 75  | 2.62               |
|                    | Otolith    | b       | Pooled      | Linear              | 120.940 | -14.211 | 0.83  | 75  | 2.71               |
|                    |            | c       | Pooled      | Linear              | 148.640 | -22.899 | 0.82  | 75  | 2.78               |
|                    | Pharyngeal | ab      | Pooled      | Non-linear          | 10.941  | 0.990   | 0.94  | 207 | 5.09               |
|                    |            | bd      | Pooled      | Non-linear          | 22.139  | 0.990   | 0.92  | 212 | 4.58               |
|                    |            | ce      | Left        | Non-linear          | 26.443  | 1.151   | 0.92  | 207 | 5.26               |
|                    |            | ce      | Right       | Non-linear          | 24.220  | 1.173   | 0.92  | 210 | 5.37               |
|                    |            | da      | Pooled      | Non-linear          | 14.971  | 0.973   | 0.94  | 209 | 4.90               |
|                    |            | bf      | Pooled      | Non-linear          | 11.531  | 0.983   | 0.94  | 207 | 5.33               |
|                    | Operculum  | a       | Pooled      | Non-linear          | 21.499  | 0.900   | 0.93  | 200 | 5.12               |
|                    |            | b       | Pooled      | Non-linear          | 11.970  | 0.914   | 0.94  | 200 | 5.10               |
|                    |            | c       | Pooled      | Non-linear          | 10.569  | 0.936   | 0.95  | 200 | 4.94               |
|                    |            | d       | Pooled      | Non-linear          | 9.797   | 0.924   | 0.95  | 178 | 5.30               |
|                    |            | e       | Pooled      | Non-linear          | 13.309  | 0.937   | 0.93  | 133 | 6.39               |
|                    |            | f       | Pooled      | Non-linear          | 15.008  | 0.947   | 0.94  | 135 | 5.74               |
|                    | Cleithrum  | a       | Pooled      | Non-linear          | 10.186  | 0.894   | 0.89  | 205 | 4.91               |
| <i>C. gibelio</i>  | Pharyngeal | a       | Pooled      | Non-linear          | 10.186  | 0.894   | 0.89  | 205 | 4.91               |
|                    |            | b       | Pooled      | Non-linear          | 10.186  | 0.894   | 0.89  | 205 | 4.91               |
|                    |            | c       | Pooled      | Non-linear          | 10.186  | 0.894   | 0.89  | 205 | 4.91               |
|                    |            | d       | Pooled      | Non-linear          | 10.186  | 0.894   | 0.89  | 205 | 4.91               |
|                    |            | e       | Pooled      | Non-linear          | 10.186  | 0.894   | 0.89  | 205 | 4.91               |
|                    |            | f       | Pooled      | Non-linear          | 10.186  | 0.894   | 0.89  | 205 | 4.91               |
|                    |            | a       | Pooled      | Non-linear          | 10.186  | 0.894   | 0.89  | 205 | 4.91               |
|                    |            | b       | Pooled      | Non-linear          | 10.186  | 0.894   | 0.89  | 205 | 4.91               |
|                    |            | c       | Pooled      | Non-linear          | 10.186  | 0.894   | 0.89  | 205 | 4.91               |
|                    |            | d       | Pooled      | Non-linear          | 10.186  | 0.894   | 0.89  | 205 | 4.91               |

|              |   |        |            |         |       |      |     |      |
|--------------|---|--------|------------|---------|-------|------|-----|------|
|              | b | Pooled | Non-linear | 14.847  | 0.970 | 0.90 | 192 | 4.59 |
|              | c | Pooled | Non-linear | 7.358   | 0.970 | 0.92 | 188 | 4.81 |
| Dorsal spine | d | Pooled | Non-linear | 26.515  | 0.909 | 0.91 | 193 | 4.78 |
| Anal spine   | a | Pooled | Non-linear | 12.627  | 0.875 | 0.89 | 145 | 5.04 |
| Otolith      | b | Pooled | Non-linear | 16.538  | 0.810 | 0.85 | 146 | 3.04 |
|              | a | Left   | Non-linear | 55.895  | 1.394 | 0.90 | 160 | 6.68 |
|              | a | Right  | Non-linear | 55.525  | 1.382 | 0.91 | 158 | 6.76 |
|              | b | Left   | Non-linear | 86.571  | 1.316 | 0.92 | 156 | 6.95 |
|              | b | Right  | Non-linear | 87.125  | 1.282 | 0.91 | 153 | 6.71 |
|              | c | Pooled | Non-linear | 136.020 | 1.358 | 0.84 | 154 | 4.80 |

**Appendix 2.** Regression statistics and percent relative errors of each structure for *Chalcalburnus chalcoides*, *Cyprinus carpio* and *Leuciscus cephalus*. See Appendix 1 for symbols, measures.

| Species              | Structure    | Measure | Orientation | Type of regressions | a      | b       | r <sup>2</sup> | n  | Relative error (%) |
|----------------------|--------------|---------|-------------|---------------------|--------|---------|----------------|----|--------------------|
| <i>C. chalcoides</i> | Pharyngeal   | ab      | Pooled      | Non-linear          | 20.865 | 0.947   | 0.96           | 47 | 8.22               |
|                      |              | bd      | Pooled      | Non-linear          | 31.757 | 0.833   | 0.96           | 47 | 8.13               |
|                      |              | ce      | Left        | Non-linear          | 49.581 | 0.890   | 0.93           | 48 | 11.30              |
|                      |              | ce      | Right       | Non-linear          | 45.340 | 0.895   | 0.93           | 48 | 10.97              |
|                      |              | da      | Pooled      | Non-linear          | 37.020 | 0.800   | 0.93           | 48 | 8.23               |
|                      |              | bf      | Pooled      | Non-linear          | 21.646 | 0.897   | 0.95           | 47 | 8.24               |
|                      | Operculum    | a       | Pooled      | Linear              | 21.941 | 44.113  | 0.82           | 16 | 8.03               |
|                      |              | b       | Pooled      | Linear              | 10.221 | 63.073  | 0.89           | 15 | 9.41               |
|                      |              | c       | Pooled      | Linear              | 10.565 | 52.312  | 0.88           | 15 | 8.65               |
|                      |              | d       | Pooled      | Linear              | 9.724  | 54.227  | 0.93           | 16 | 8.39               |
|                      |              | e       | Pooled      | Linear              | 11.947 | 86.672  | 0.81           | 16 | 8.71               |
|                      |              | f       | Pooled      | Linear              | 12.853 | 52.743  | 0.86           | 16 | 8.61               |
|                      | Cleithrum    | a       | Pooled      | Linear              | 11.763 | 44.180  | 0.82           | 28 | 8.96               |
|                      |              | b       | Pooled      | Non-linear          | 16.914 | 0.862   | 0.91           | 24 | 9.00               |
|                      |              | c       | Pooled      | Non-linear          | 11.793 | 0.917   | 0.90           | 24 | 8.81               |
|                      |              | d       | Pooled      | Non-linear          | 26.165 | 0.834   | 0.88           | 24 | 8.99               |
| <i>C. carpio</i>     | Pharyngeal   | ab      | Pooled      | Non-linear          | 7.386  | 1.103   | 0.97           | 15 | 8.36               |
|                      |              | bd      | Pooled      | Linear              | 16.998 | 26.804  | 0.96           | 15 | 8.45               |
|                      |              | ce      | Pooled      | Linear              | 38.884 | -48.468 | 0.85           | 15 | 12.20              |
|                      |              | da      | Pooled      | Non-linear          | 10.897 | 1.071   | 0.96           | 15 | 8.85               |
|                      |              | bf      | Pooled      | Non-linear          | 10.628 | 0.992   | 0.97           | 15 | 8.61               |
|                      | Operculum    | a       | Pooled      | Non-linear          | 19.787 | 0.899   | 0.97           | 13 | 8.18               |
|                      |              | b       | Pooled      | Non-linear          | 8.415  | 1.043   | 0.96           | 13 | 8.30               |
|                      |              | c       | Pooled      | Non-linear          | 7.237  | 1.056   | 0.98           | 13 | 8.76               |
|                      |              | d       | Pooled      | Linear              | 8.589  | -20.124 | 0.98           | 13 | 8.85               |
|                      |              | e       | Pooled      | Linear              | 15.796 | -45.813 | 0.83           | 13 | 8.11               |
|                      |              | f       | Pooled      | Non-linear          | 6.374  | 1.226   | 0.88           | 13 | 8.55               |
|                      | Cleithrum    | a       | Pooled      | Non-linear          | 6.027  | 1.072   | 0.97           | 14 | 8.98               |
|                      |              | b       | Pooled      | Linear              | 6.742  | 53.688  | 0.96           | 13 | 8.68               |
|                      |              | c       | Pooled      | Linear              | 8.045  | -71.063 | 0.95           | 13 | 9.22               |
|                      |              | d       | Pooled      | Non-linear          | 43.723 | 0.610   | 0.94           | 15 | 8.08               |
|                      | Dorsal spine | a       | Pooled      | Non-linear          | 29.153 | 0.641   | 0.90           | 4  | 15.63              |
|                      | Anal spine   | b       | Pooled      | Linear              | 5.649  | 87.095  | 0.90           | 4  | 15.30              |
| <i>L. cephalus</i>   | Pharyngeal   | ab      | Pooled      | Non-linear          | 12.161 | 1.047   | 0.97           | 10 | 19.24              |
|                      |              | bd      | Pooled      | Linear              | 18.555 | -1.054  | 0.97           | 11 | 17.75              |
|                      |              | ce      | Left        | Linear              | 23.784 | 16.716  | 0.97           | 11 | 27.62              |
|                      |              | ce      | Right       | Linear              | 22.421 | 13.546  | 0.95           | 11 | 27.40              |
|                      |              | da      | Pooled      | Linear              | 16.877 | 4.845   | 0.99           | 10 | 19.17              |
|                      |              | bf      | Pooled      | Non-linear          | 14.434 | 0.954   | 0.98           | 10 | 19.08              |
|                      | Operculum    | a       | Pooled      | Linear              | 23.929 | 4.187   | 0.98           | 11 | 24.49              |
|                      |              | b       | Pooled      | Linear              | 12.113 | 12.451  | 0.99           | 11 | 23.06              |

|           |   |        |            |        |       |      |    |       |
|-----------|---|--------|------------|--------|-------|------|----|-------|
|           | c | Pooled | Non-linear | 14.441 | 0.942 | 0.99 | 11 | 24.10 |
|           | d | Pooled | Linear     | 11.604 | 6.922 | 0.98 | 11 | 23.52 |
|           | e | Pooled | Linear     | 19.444 | 0.164 | 0.95 | 11 | 23.31 |
|           | f | Pooled | Non-linear | 15.997 | 0.990 | 0.98 | 11 | 24.87 |
| Cleithrum | a | Pooled | Non-linear | 10.836 | 1.034 | 0.98 | 12 | 21.19 |
|           | b | Pooled | Non-linear | 12.348 | 0.944 | 0.99 | 12 | 21.54 |
|           | c | Pooled | Non-linear | 8.673  | 0.983 | 0.99 | 12 | 21.38 |
|           | d | Pooled | Non-linear | 19.368 | 0.968 | 0.98 | 12 | 22.09 |

**Appendix 3.** Regression statistics and percent relative errors of each structure for *Petroleuciscus borysthenicus*, *Rutilus rutilus* and *Scardinius erythrophthalmus*. See Appendix 1 for symbols and measures.

| Species                    | Structure  | Measure | Orientation | Type of regressions | a       | b      | r <sup>2</sup> | n    | Relative error (%) |
|----------------------------|------------|---------|-------------|---------------------|---------|--------|----------------|------|--------------------|
| <i>P. borysthenicus</i>    | Pharyngeal | ab      | Pooled      | Non-linear          | 15.931  | 0.962  | 0.94           | 19   | 4.70               |
|                            |            | bd      | Pooled      | Linear              | 15.996  | 19.981 | 0.92           | 19   | 4.73               |
|                            |            | ce      | Left        | Linear              | 24.234  | 17.821 | 0.89           | 19   | 7.06               |
|                            |            | ce      | Right       | Linear              | 18.929  | 30.655 | 0.87           | 19   | 6.86               |
|                            |            | da      | Pooled      | Non-linear          | 20.997  | 0.945  | 0.91           | 19   | 4.74               |
|                            |            | bf      | Pooled      | Non-linear          | 19.795  | 0.828  | 0.92           | 19   | 4.83               |
|                            | Operculum  | a       | Pooled      | Linear              | 29.648  | 2.355  | 0.87           | 12   | 6.06               |
|                            |            | b       | Pooled      | Non-linear          | 14.077  | 1.009  | 0.91           | 15   | 5.41               |
|                            |            | c       | Pooled      | Non-linear          | 13.287  | 0.979  | 0.92           | 15   | 5.60               |
|                            |            | d       | Pooled      | Linear              | 12.723  | -3.194 | 0.93           | 13   | 5.93               |
|                            |            | e       | Pooled      | Non-linear          | 23.668  | 0.887  | 0.86           | 13   | 5.90               |
|                            |            | f       | Pooled      | Non-linear          | 18.038  | 0.905  | 0.87           | 16   | 5.44               |
|                            | Cleithrum  | a       | Pooled      | Non-linear          | 12.390  | 0.972  | 0.86           | 17   | 5.06               |
|                            |            | b       | Pooled      | Non-linear          | 13.656  | 0.892  | 0.92           | 19   | 4.93               |
|                            |            | c       | Pooled      | Non-linear          | 10.184  | 0.902  | 0.93           | 17   | 5.05               |
|                            |            | d       | Pooled      | Non-linear          | 29.825  | 0.780  | 0.87           | 19   | 4.88               |
| <i>R. rutilus</i>          | Pharyngeal | ab      | Pooled      | Non-linear          | 17.627  | 0.943  | 0.94           | 636  | 1.24               |
|                            |            | bd      | Left        | Non-linear          | 21.982  | 0.955  | 0.95           | 640  | 1.78               |
|                            |            | bd      | Right       | Non-linear          | 22.088  | 0.957  | 0.95           | 633  | 1.76               |
|                            |            | ce      | Left        | Non-linear          | 37.056  | 0.859  | 0.92           | 655  | 1.90               |
|                            |            | ce      | Right       | Non-linear          | 34.055  | 0.857  | 0.92           | 650  | 1.71               |
|                            |            | da      | Pooled      | Non-linear          | 22.977  | 0.880  | 0.96           | 1501 | 1.22               |
|                            |            | bf      | Pooled      | Non-linear          | 16.453  | 0.939  | 0.95           | 636  | 1.25               |
|                            | Operculum  | a       | Pooled      | Non-linear          | 33.515  | 0.880  | 0.94           | 503  | 1.76               |
|                            |            | b       | Pooled      | Non-linear          | 15.878  | 0.914  | 0.97           | 506  | 1.73               |
|                            |            | c       | Pooled      | Non-linear          | 14.232  | 0.925  | 0.97           | 511  | 1.76               |
|                            |            | d       | Pooled      | Non-linear          | 13.886  | 0.917  | 0.97           | 499  | 1.92               |
|                            |            | e       | Pooled      | Non-linear          | 19.184  | 0.922  | 0.92           | 491  | 1.63               |
|                            |            | f       | Pooled      | Non-linear          | 17.411  | 0.925  | 0.96           | 502  | 1.72               |
|                            | Cleithrum  | a       | Pooled      | Non-linear          | 16.075  | 0.854  | 0.94           | 575  | 1.24               |
|                            |            | b       | Pooled      | Non-linear          | 12.562  | 0.935  | 0.97           | 565  | 1.26               |
|                            |            | c       | Pooled      | Non-linear          | 10.230  | 0.902  | 0.95           | 544  | 1.31               |
|                            |            | d       | Pooled      | Non-linear          | 23.366  | 0.882  | 0.92           | 566  | 1.76               |
|                            | Otolith    | a       | Left        | Non-linear          | 6.182   | 1.283  | 0.89           | 531  | 1.82               |
|                            |            | a       | Right       | Non-linear          | 6.299   | 1.284  | 0.88           | 512  | 1.83               |
|                            |            | b       | Pooled      | Non-linear          | 10.757  | 1.293  | 0.86           | 514  | 1.34               |
|                            |            | c       | Pooled      | Linear              | 104.310 | 39.263 | 0.66           | 237  | 1.43               |
| <i>S. erythrophthalmus</i> | Pharyngeal | ab      | Pooled      | Non-linear          | 17.627  | 0.943  | 0.94           | 636  | 4.44               |
|                            |            | bd      | Left        | Non-linear          | 21.982  | 0.955  | 0.95           | 640  | 5.18               |
|                            |            | bd      | Right       | Non-linear          | 22.088  | 0.957  | 0.95           | 633  | 5.32               |
|                            |            | ce      | Left        | Non-linear          | 37.056  | 0.859  | 0.92           | 655  | 5.30               |
|                            |            | ce      | Right       | Non-linear          | 34.055  | 0.857  | 0.92           | 650  | 5.79               |
|                            |            | da      | Pooled      | Non-linear          | 22.977  | 0.880  | 0.96           | 651  | 4.19               |



|           |    |        |            |        |       |      |     |      |
|-----------|----|--------|------------|--------|-------|------|-----|------|
| Operculum | bf | Pooled | Non-linear | 16.453 | 0.939 | 0.95 | 636 | 4.09 |
|           | a  | Pooled | Non-linear | 33.515 | 0.880 | 0.94 | 503 | 6.12 |
|           | b  | Pooled | Non-linear | 15.878 | 0.914 | 0.97 | 506 | 4.76 |
|           | c  | Pooled | Non-linear | 14.232 | 0.925 | 0.97 | 510 | 4.70 |
|           | d  | Pooled | Non-linear | 13.886 | 0.917 | 0.97 | 498 | 4.79 |
|           | e  | Pooled | Non-linear | 19.184 | 0.922 | 0.92 | 491 | 5.01 |
| Cleithrum | f  | Pooled | Non-linear | 17.411 | 0.925 | 0.96 | 502 | 4.83 |
|           | a  | Pooled | Non-linear | 16.075 | 0.854 | 0.94 | 575 | 3.07 |
|           | b  | Pooled | Non-linear | 12.562 | 0.935 | 0.97 | 565 | 3.81 |
|           | c  | Pooled | Non-linear | 10.230 | 0.902 | 0.95 | 544 | 3.25 |
| Otolith   | d  | Pooled | Non-linear | 23.366 | 0.882 | 0.92 | 567 | 3.98 |
|           | a  | Left   | Non-linear | 6.182  | 1.283 | 0.89 | 531 | 3.62 |
|           | a  | Right  | Non-linear | 6.299  | 1.284 | 0.88 | 512 | 3.84 |
|           | b  | Pooled | Non-linear | 10.757 | 1.293 | 0.86 | 514 | 2.62 |
|           | c  | Pooled | Non-linear | 13.914 | 1.212 | 0.79 | 495 | 2.73 |

**Appendix 4.** Regression statistics and percent relative errors of each structure for *Tinca tinca* and *Vimba vimba*. See Appendix 1 for symbols and measures.

| Species         | Structure  | Measure | Orientation | Type of regressions | a      | b      | r <sup>2</sup> | n    | Relative error (%) |
|-----------------|------------|---------|-------------|---------------------|--------|--------|----------------|------|--------------------|
| <i>T. tinca</i> | Pharyngeal | ab      | Pooled      | Linear              | 11.935 | 0.236  | 0.91           | 21   | 7.56               |
|                 |            | bd      | Pooled      | Linear              | 17.942 | 42.495 | 0.93           | 21   | 7.02               |
|                 |            | ce      | Pooled      | Non-linear          | 42.002 | 0.911  | 0.82           | 21   | 7.54               |
|                 |            | da      | Pooled      | Non-linear          | 18.337 | 0.922  | 0.91           | 21   | 7.57               |
|                 |            | bf      | Pooled      | Linear              | 12.476 | 11.437 | 0.93           | 21   | 7.54               |
|                 | Operculum  | a       | Pooled      | Linear              | 14.282 | 32.091 | 0.93           | 15   | 7.42               |
|                 |            | b       | Pooled      | Linear              | 8.859  | 28.257 | 0.94           | 14   | 8.10               |
|                 |            | c       | Pooled      | Linear              | 8.414  | 21.716 | 0.95           | 15   | 7.63               |
|                 |            | d       | Pooled      | Linear              | 8.211  | 17.927 | 0.97           | 15   | 7.36               |
|                 |            | e       | Pooled      | Linear              | 13.039 | 25.180 | 0.93           | 14   | 7.94               |
|                 | Cleithrum  | f       | Pooled      | Linear              | 10.484 | 20.395 | 0.97           | 14   | 8.13               |
|                 |            | a       | Pooled      | Linear              | 7.679  | 17.983 | 0.93           | 23   | 7.25               |
|                 |            | b       | Pooled      | Linear              | 10.591 | 4.426  | 0.93           | 23   | 7.14               |
|                 |            | c       | Pooled      | Linear              | 6.131  | 11.682 | 0.93           | 27   | 7.37               |
|                 | d          | Pooled  | Linear      | 15.412              | 50.188 | 0.86   | 23             | 6.97 |                    |
| <i>V. vimba</i> | Pharyngeal | ab      | Pooled      | Non-linear          | 18.362 | 1.007  | 0.97           | 399  | 5.79               |
|                 |            | bd      | Pooled      | Non-linear          | 36.227 | 0.857  | 0.96           | 300  | 5.96               |
|                 |            | ce      | Left        | Non-linear          | 46.501 | 0.943  | 0.96           | 302  | 9.34               |
|                 |            | ce      | Right       | Non-linear          | 41.880 | 0.954  | 0.96           | 300  | 8.59               |
|                 |            | da      | Pooled      | Non-linear          | 26.313 | 0.937  | 0.97           | 399  | 5.68               |
|                 |            | bf      | Pooled      | Non-linear          | 20.930 | 0.928  | 0.98           | 300  | 5.69               |
|                 | Operculum  | a       | Pooled      | Non-linear          | 30.512 | 0.879  | 0.97           | 265  | 4.10               |
|                 |            | b       | Pooled      | Non-linear          | 18.523 | 0.893  | 0.98           | 265  | 4.01               |
|                 |            | c       | Pooled      | Non-linear          | 17.600 | 0.884  | 0.98           | 265  | 3.99               |
|                 |            | d       | Pooled      | Non-linear          | 16.475 | 0.887  | 0.98           | 264  | 3.93               |
|                 |            | e       | Pooled      | Non-linear          | 28.417 | 0.851  | 0.96           | 265  | 4.11               |
|                 |            | f       | Pooled      | Non-linear          | 21.448 | 0.914  | 0.97           | 266  | 3.94               |
|                 | Cleithrum  | a       | Pooled      | Non-linear          | 16.810 | 0.892  | 0.96           | 303  | 3.90               |
|                 |            | b       | Pooled      | Non-linear          | 15.895 | 0.895  | 0.98           | 310  | 3.98               |
|                 |            | c       | Pooled      | Non-linear          | 11.224 | 0.917  | 0.97           | 299  | 4.33               |
|                 |            | d       | Pooled      | Non-linear          | 28.236 | 0.849  | 0.97           | 311  | 3.97               |
|                 | Otolith    | a       | Pooled      | Non-linear          | 68.769 | 1.324  | 0.91           | 215  | 4.02               |
|                 |            | b       | Pooled      | Non-linear          | 124.85 | 1.279  | 0.90           | 199  | 4.42               |
|                 |            | c       | Pooled      | Non-linear          | 157.36 | 1.172  | 0.86           | 201  | 4.15               |