

# **Development of methods in support of a head-only sampling program for Sablefish (*Anoplopoma fimbria*) in British Columbia**

Lisa C. Lacko, Kathryn L. Temple, Kendra R. Holt, Janine x. Supernault,  
Allen R. Kronlund, Malcolm R. Wyeth and Brendan M. Connors

Pacific Biological Station  
Fisheries and Oceans Canada, 3190 Hammond Bay Road  
Nanaimo, British Columbia, V9T 6N7, Canada

2022

**Canadian Technical Report of  
Fisheries and Aquatic Sciences ####**



Fisheries and Oceans  
Canada      Pêches et Océans  
Canada

**Canada**

## **Canadian Technical Report of Fisheries and Aquatic Sciences**

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of Fisheries and Oceans Canada, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in the data base *Aquatic Sciences and Fisheries Abstracts*.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page.

Numbers 1-456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457-714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715-924 were issued as Department of Fisheries and Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

## **Rapport technique canadien des sciences halieutiques et aquatiques**

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques de Pêches et Océans Canada, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications à part entière. Le titre exact figure au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la base de données *Résumés des sciences aquatiques et halieutiques*.

Les rapports techniques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre.

Les numéros 1 à 456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

Canadian Technical Report of  
Fisheries and Aquatic Sciences nnn

2022

DEVELOPMENT OF METHODS IN SUPPORT OF A HEAD-ONLY SAMPLING PROGRAM FOR  
SABLEFISH (*ANOPLOPOMA FIMBRIA*) IN BRITISH COLUMBIA

by

Lisa C. Lacko and Kathryn L. Temple and Kendra R. Holt and Janine x. Supernault and Allen R. Kronlund and Malcolm R. Wyeth and Brendan M. Connors

Pacific Biological Station  
Fisheries and Oceans Canada, 3190 Hammond Bay Road  
Nanaimo, British Columbia, V9T 6N7, Canada

© Her Majesty the Queen in Right of Canada, 2022  
Cat. No. Fs97-6/nnnE-PDF ISBN ISSN 1488-5379

Correct citation for this publication:

L.C. Lacko, Temple K.L., Holt K.R., Supernault J.x., and Kronlund A.R, Wyeth M.R., and Connors, B.M. 2022. Development of methods in support of a head-only sampling program for Sablefish (*Anoplopoma fimbria*) in British Columbia. Can. Tech. Rep. Fish. Aquat. Sci. nnn: vi + 17 p.

## CONTENTS

<b>ABSTRACT</b>	v
<b>RÉSUMÉ</b>	vi
<b>1 Introduction</b>	1
<b>2 Methods</b>	2
2.1 Experimental Study 2016 . . . . .	2
2.1.1 Data Collection . . . . .	2
2.1.2 Data Analysis . . . . .	2
2.1.3 Gender Test Development . . . . .	2
2.2 Fishery Pilot Study 2017 . . . . .	3
2.2.1 Data Collection . . . . .	3
2.2.2 Data Analysis . . . . .	3
<b>3 Results</b>	4
3.1 Experimental Study 2016 . . . . .	4
3.1.1 Genetic sex-determination . . . . .	4
3.2 Fishery Pilot Study 2017 . . . . .	4
<b>4 Discussion</b>	5
4.1 Identification of Preferred Head Cranial Dimension . . . . .	5
4.2 Implementation of Head-Only Catch Sampling in 2018 . . . . .	5
4.3 Future Monitoring and Research Requirements . . . . .	5
<b>5 Acknowledgments</b>	6
<b>6 Tables</b>	7
<b>7 Figures</b>	10

<b>APPENDICES</b>	<b>13</b>
<b>A IMAGES OF THE SIX CRANIAL DIMENSION MEASUREMENTS.</b>	<b>13</b>
<b>B SEX DETERMINATION BY OPERCULUM MARKING</b>	<b>14</b>
<b>C SABLEFISH BIOSAMPLE COLLECTION PROGRAM NOTICE</b>	<b>15</b>
<b>D TAGGED SABLEFISH PROGRAM NOTICE</b>	<b>16</b>
<b>8 References</b>	<b>17</b>

## ABSTRACT

L.C. Lacko, Temple K.L., Holt K.R., Supernault J.x., and Kronlund A.R., Wyeth M.R., and Connors, B.M. 2022. Development of methods in support of a head-only sampling program for Sablefish (*Anoplopoma fimbria*) in British Columbia. Can. Tech. Rep. Fish. Aquat. Sci. nnn: vi + 17 p.

Routine biological sampling of whole round Sablefish from commercial fishing operations in British Columbia began the early 1990's. Historically, specimens were obtained through the voluntary fishery catch sampling and tag recovery programs. We investigate the potential for obtaining sex and length information using heads, rather than the entire fish. In 2016, 438 Sablefish (240-1080 mm) were sampled at sea and six different fish head measurements were collected. Genetic samples (137) were obtained to develop methods of DNA-based sex identification. A pilot study occurred in 2017 with 360 head-only samples collected and sexed by a commercial vessel, followed by scientific sampling on shore. Regression analysis results reveal that all six cranial dimensions can be used to accurately predict length. However, interorbital distance was not only a good predictor of length, but samplers ranked it the most efficient to measure and easily repeatable. Genomic DNA were successfully processed for 130 of the 137 samples. Fisher sex determination was accurate xx% of the time. Given the results, the 2018 sampling collection program was modified so that returns of whole round Sablefish were replaced by head-only samples with knife cuts on the operculum to indicate sex.

## RÉSUMÉ

L.C. Lacko, Temple K.L., Holt K.R., Supernault J.x., and Kronlund A.R., Wyeth M.R., and Connors, B.M. 2022. Development of methods in support of a head-only sampling program for Sablefish (*Anoplopoma fimbria*) in British Columbia. Can. Tech. Rep. Fish. Aquat. Sci. nnn: vi + 17 p.

## 1 Introduction

Biological samples of British Columbia (BC) Sablefish (*Anoplopoma fimbria*) from trap and hook & line fisheries have been collected from a fishery-based voluntary catch sampling program since 1995 (Haist and Wyeth 2001). Prior to 2018, samples of whole fish were collected dockside by Fisheries and Oceans Canada (DFO) port samplers and/or contracted service providers via the dockside monitoring program (DMP). In addition, tagged fish were collected whole from commercial fisheries (trap, trawl, hook & line) at the point of landing, also by the dockside monitoring program. Biological data collected during catch sampling included fork length and sex. Otoliths were collected and archived for future ageing analyses. These data provide a fishery-dependent source of age and size composition data for the two-sex age structured operating model used as part of the BC Management Strategy Evaluation (MSE) (Cox et al. 2019, DFO 2020).

Between 2016 and 2017, DFO undertook research to evaluate the potential change to a ‘head-only’ catch sampling program for Sablefish trap and hook & line fisheries. Head-only sampling allows commercial crew to collect samples directly while at sea, thereby eliminating the need to rely on port samplers for dockside data collection as well as the need for fishing vessels to store whole fish. Instead, commercial crew J-cut the fish at-sea as per commercial practice, examine the gonads to determine sex, mark the sex with knife cuts on the operculum, and store the frozen head (and/or floy tag) for later sampling by science personal. The key motivation for switching to a head-only sampling program was to increase the number and spatial distribution of fishery samples, while simultaneously maintaining the quality of biological data.

Evaluating the potential success of a head-only sampling program for Sablefish required the development and testing of new methods. First, a method was required to estimate Sablefish fork length from head samples. Previous research on other fish species has shown that lengths can be accurately estimated from head dimensions (Serafy et al. 1996; Park et al. 2007), head and mandible lengths (Isermann and Vandergoot 2005), and head height to eye diameter ratio (Richardson et al. 2015), so we considered six different cranial measurement linked to these as a starting point for Sablefish. Second, the accuracy of methods used by commercial fishing crew to identify and record sex needed to be assessed, which required the development of DNA analysis techniques for Sablefish gender detection.

In this technical report, we describe a research investigation in 2016 that collected Sablefish samples to: 1) identify cranial measurements that are reliable predictors of fork length for Sablefish; 2) identify cranial measurements that are practical to measure; and 3) develop DNA analysis techniques for gender detection. We also describe the fishery pilot study in 2017 to: 1) assess the ease of sample collection for commercial fishers; 2) determine the accuracy of fisher sex identification using DNA analysis; followed by 3) among-sampler measurements of each head morphometric.

## 2 Methods

### 2.1 Experimental Study 2016

#### 2.1.1 Data Collection

Sablefish were randomly selected for sampling during the 2016 biennial DFO Groundfish Synoptic Bottom Trawl surveys using a length stratified sampling protocol. A total of 212 fish were sampled on the West Coast Vancouver Island survey (Williams et al. 2018) and 219 fish on the West Coast Haida Gwaii survey (Nottingham et al. 2018) (Figure 1). In addition, seven small Sablefish were collected during the 2016 salmon survey (put reference here when obtain).

For each sampled fish, fork length, round weight, sex, and maturity were recorded at sea. The heads were removed, labelled with an unique ID and frozen. On shore, six different cranial dimensions were considered, including upper jaw ( $L_{UJ}$ ), eye diameter ( $L_{ED}$ ), interorbital distance ( $L_{ID}$ ), snout length ( $L_{SL}$ ), post orbital to preoperculum distance ( $L_{PP}$ ) and post orbital head length ( $L_{PO}$ ) (Table 1, Appendix A). All measurements were taken using Mitutoyo Absolute® 500-762-20 coolant proof digimatic calipers. The post orbital head length ( $L_{PO}$ ) measurement was abandoned after testing 130 Sablefish due to low sample quality and technical issues.

At the time of sampling, each cranial dimension was evaluated by three experienced samplers on a five point rating scale in terms of two distinct metrics: 1. ease of use and 2. repeatability. The ease of use metric focused on three key attributes of the measurement: learn-ability (task understanding), efficiency (task-completion time) and degree of difficulty (task performance ease). The repeatability metric focused on ranking each measurement under repeated caliper placement, taking into consideration that a proper understanding of how to place caliper jaws on soft or hard (bone) tissues may affect the final measurement.

Once head measurements were completed, sagittal otoliths were collected for future ageing from all fish. Operculum clips (DNA) were also collected from the first 137 fish measured and forwarded to the Molecular Genetics Laboratory (MGL) for analysis and gender test development.

#### 2.1.2 Data Analysis

Using the `lm()` function in R (R Core Team 2021), fork lengths ( $L_{FL}$ ) were estimated by a simple linear regression model, using the six cranial measurements as predictor variables to see how well each could predict fork length. Model fit was evaluated in terms of statistical significance (p-value less than 0.05) and coefficients of determination ( $R^2$ ).

#### 2.1.3 Gender Test Development

In the Molecular Genetics Laboratory (MGL), gender test methods were successfully developed. X-insert and Y-insert specific primers developed by Rondeau et al. (2013) were used in a

modified format (J. Taylor, personal communication, August 4, 2022) to produce polymerase chain reaction (PCR) amplified products 155 and 170bp in size, respectively. These fluorescently labelled sex specific alleles were then size fractionated with the ABI 3730xl DNA Analyzer and scored using GeneMapper Software v5.0 (Applied Biosystems, Carlsbad, CA) and an internal lane sizing standard.

## 2.2 Fishery Pilot Study 2017

### 2.2.1 Data Collection

In 2017, a pilot study was conducted with the commercial sector returning Sablefish samples. A total of 360 heads were collected from J-cut Sablefish on a limited-entry fishery trip to the Cobb and Eickelberg seamounts (Figure 1). Each operculum was marked by commercial fishers with either one knife cut (male) or two knife cuts (female) (Appendix B).

Scientific sampling occurred on shore, with the first 99 heads of the pilot study measured by three experienced technicians for  $L_{ID}$ ,  $L_{SL}$ ,  $L_{UJ}$  and  $L_{PP}$  to determine the most repeatable measurement. The cranial dimensions of  $L_{ED}$  and  $L_{PO}$  were eliminated after the results from the 2016 experimental study.

### 2.2.2 Data Analysis

The precision of cranial measurements among readers was quantified using and Index of Average Error (IAE) (Beamish and Fournier 1981). Beamish and Fournier (1981) used IAE as a metric for comparing the precision of age determinations among different readers; however, given that three highly trained technicians were used to conduct the same measurements, we do not look at differences in IAE among technicians. Instead, we compare IAE among the four cranial dimensions considered to identify which one provided the most consistent measurement among readers. In our case, IAE was defined as:

$$IAE = \frac{1}{N} \sum_{j=1}^N \left[ \frac{1}{R} \sum_{i=1}^R \frac{|X_{ij} - X_j|}{X_j} \right]$$

where N is the number of Sablefish measured for each cranial measurement, R is the number of times each cranial measurement was taken by a technician,  $X_{ij}$  is the average cranial length for the jth Sablefish, and  $X_{ij}$  is the ith cranial measurement of the jth Sablefish.

Fin clips (95 of 99) were forwarded to the molecular genetics lab for an audit of fisher sex determination.

### 3 Results

#### 3.1 Experimental Study 2016

A total of 438 specimens comprising 222 males and 216 females were evaluated for this study from three different surveys (Table 2). The post orbital head length ( $L_{PO}$ ) measurement had a lower number of measurements compared to other methods because it was abandoned after testing 130 Sablefish due to technical issues. The smallest fork length of the collected specimens was 240 mm, the largest was 1080 mm, and the average was 573.3 mm.

Table 3 lists the statistics of the cranial dimensions ( $L_{UJ}$ ,  $L_{ED}$ ,  $L_{ID}$ ,  $L_{SL}$ ,  $L_{PP}$ ,  $L_{PO}$ ) as predictors of the response variable fork length ( $L_{FL}$ ). All cranial dimensions were highly correlated with fork length with high R-squared values and significant p-values (Figure 2 and Figure 3).

The correlation coefficient ( $r$ ) was highest for female measurements of  $L_{SL}$  (0.983) and  $L_{ID}$  (0.98) and male measurements of  $L_{UJ}$  (0.974) and  $L_{SL}$  (0.972). The R-squared values were highest for female measurements of  $L_{SL}$  (0.966) and  $L_{ID}$  (0.96) and male measurements of  $L_{UJ}$  (0.949) and  $L_{SL}$  (0.945).

In general, the samplers reported that measurement errors were more probable when the calipers were placed on soft tissues, rather than bone. The technicians scored interorbital distance ( $L_{ID}$ ) as the highest on the five point scale for ease of use and repeatable criteria, and eye diameter ( $L_{ED}$ ) and postorbital head length ( $L_{PO}$ ) were scored as the lowest (Table 4).  $L_{ID}$  (narrowest distance between the eye sockets) proved the easiest measurement as the tissue could be easily compressed with the caliper jaws to obtain bone measurements.  $L_{ED}$  (anterior-posterior diameter of eye socket) proved hard to repeat on soft tissue.  $L_{PO}$  (posterior inner edge of orbit to dorsal insertion of opercle) was difficult to measure since many opercula were damaged during head removal by J-cut, the endpoints were difficult to define and the electronic calipers malfunctioned with lengths >15 cm.

##### 3.1.1 Genetic sex-determination

Genomic DNA (130 of 137 fin clips) were successfully PCR amplified by the MGL to determine sex. The accuracy of sex detection for the experimental study by the science technicians was 92% (119/130).

#### 3.2 Fishery Pilot Study 2017

The 360 heads received from the commercial vessel were in good condition and operculum cuts worked well to indicate sex. The first 99 heads (60 from Cobb Seamount, 39 from Eikelberg seamount) were measured once by three expert science technicians for each cranial dimension of  $L_{UJ}$ ,  $L_{ID}$ ,  $L_{SL}$  and  $L_{PP}$  (Table 5). The cranial measurements that produced the lowest mean error were upper jaw ( $L_{UJ}$ ) and interorbital distance ( $L_{ID}$ ) with IAE values of 1 % and 1.1 %, respectively (Table 5). The accuracy of the commercial fisher sex detection from the DNA gender

analysis was xx% (x/95).

## 4 Discussion

### 4.1 Identification of Preferred Head Cranial Dimension

Interorbital distance ( $L_{ID}$ ) is recommended as the most suitable cranial measurement for predicting Sablefish fork length in support of a head-only sampling program. All cranial dimensions collected in the research investigation were found to be good predictors of Sablefish fork length ( $L_{FL}$ ) with resulting  $R^2$  values for male and female Sablefish >0.88. However,  $L_{ID}$  was the only method to be ranked as 5 out of 5 for both ease of use and repeatability criteria. In addition, feedback from technicians participating in both the experimental and fishery pilot studies identified  $L_{ID}$  as the preferred method because it could be performed quickly, accurately, and repeatedly. Technicians reported that measurement errors were more probable when the calipers were placed on soft tissues. Other errors were a result of the manner in which the Sablefish were handled before measurement, ie. the operculum was cut off. The high ease of use and repeatability scores for the  $L_{ID}$  method was largely due to it being based on bone measurements rather than soft tissue.

### 4.2 Implementation of Head-Only Catch Sampling in 2018

The identification of a reliable method for predicting Sablefish fork length from interorbital distance combined with both the successful development of a genetic protocol to determine sex by the Pacific Biological Station Molecular Genetics Laboratory and the confirmation that fisher sex-ID at sea was [ . . . . to be filled in with results . . . . ] resulted in a switch to head-only sampling for Sablefish starting in 2018.

Routine biological sampling procedures were modified so that commercial fishers are now only returning head samples, rather than the entire fish (Appendix C). Fishing vessels are requested to voluntarily collect a samples at each 50,000 lbs catch increment for the vessel throughout the year (e.g., 50,000 lbs, 100,000 lbs, 150,000 lbs, etc), with each sample consisting of approximately 20 to 25 Sablefish. The established sampling protocol requires sampled fish to be collected from traps or skates spread along the string being sampled. For longline gear, this is achieved by sampling the first 5 fish from each skate until the target of 20 - 25 fish is reached, while for trap gear, a select number of traps are marked for sampling ahead of time and all fish within marked traps are sampled. The tagged Sablefish program was also modified so that fishers return heads and tags only (Appendix D). These changes have improved the fishers ability to freeze, store and transport samples due to less requirement for storage space.

### 4.3 Future Monitoring and Research Requirements

We recommend that a review of the head-only sample program be undertaken in 2024-2025 to better examine how sampling rates, length composition, and spatial distribution have changed

compared to the previous sampling program. By this time, the sampling program will have been in place for at least 6 years. At least 6 years is recommended to allow for a couple years of adjustment to the new protocols, as well as to allow for some lags in at-shore sampling of heads due to the COVID-19 pandemic.

## **5 Acknowledgments**

We thank Schon Acheson and Kristina Castle for lending their skilled technical expertise for this report and the Molecular Genetics Laboratory (Mike Wetklo and Gordon Skocic) at PBS for assistance in development of the Sablefish gender tests. A special thanks to the crew of the Pacific Viking for participating in the pilot project and the Wild Canadian Sablefish Association for their continued support of science work with the Science Rewards Program.

## 6 Tables

Table 1. List of head dimensions for upper jaw length ( $L_{UJ}$ ), eye diameter ( $L_{ED}$ ), interorbital distance ( $L_{ID}$ ), snout length ( $L_{SL}$ ), post orbital to preoperculum distance ( $L_{PP}$ ) and post orbital head length ( $L_{PO}$ ) measurement descriptions and specification of caliper jaw placement. Many follow the morphological measurements described in (Shaw and McFarlane 1997). The matching images are found in Appendix A.

<b>Head dimension</b>	<b>Head description</b>	<b>Caliper jaw position</b>
$L_{UJ}$	Tip of snout to the posterior edge of the maxilla.	Outside caliper jaw measurement from forward point and centre of snout to back of maxilla.
$L_{ED}$	Anterior-posterior diameter of eye socket.	Inside caliper jaw measurement firmly stretched against eye socket at vertical midpoint of eye.
$L_{ID}$	Narrowest distance between eye sockets, measured on dorsal surface.	Outside caliper jaw measurement of the horizontal midpoint of eyes on dorsal surface.
$L_{SL}$	Tip of snout to anterior inner edge of eye socket.	Outside caliper jaw measurement from forward point and centre of snout to horizontal midpoint of anterior edge of eye socket.
$L_{PP}$	Posterior inner edge of orbit to visual insertion point of preopercle.	Outside caliper jaw measurement from back of eye socket to preopercle bone insertion point. Preopercle must be lifted to expose preopercle bone underneath.
$L_{PO}$	Posterior inner edge of orbit to dorsal insertion of opercle.	Outside caliper jaw measurement from back of eye socket to bone underneath gill cover notch at dorsal insertion of the opercle. The operculum must be held taut.

Table 2. Summary of Sablefish biological data collected during the 2016 experimental study. Tally of fork lengths ( $L_{FL}$ ), round weights (RW), upper jaw lengths ( $L_{UJ}$ ), eye diameters ( $L_{ED}$ ), interorbital distances ( $L_{ID}$ ), snout lengths ( $L_{SL}$ ), post orbital to preoperculum distances ( $L_{PP}$ ), post orbital head lengths ( $L_{PO}$ ), females (F), males (M), sagittal otoliths and operculum clips (DNA) listed by survey.

<b>Survey</b>	<b><math>L_{FL}</math></b>	<b>RW</b>	<b><math>L_{UJ}</math></b>	<b><math>L_{ED}</math></b>	<b><math>L_{ID}</math></b>	<b><math>L_{SL}</math></b>	<b><math>L_{PP}</math></b>	<b><math>L_{PO}</math></b>	<b>F</b>	<b>M</b>	<b>Otoliths</b>	<b>DNA</b>	<b>Total</b>
2016 WCHG	219	219	219	219	218	219	207	52	111	108	219	59	219
2016 WCVI	212	212	211	212	212	211	212	78	105	107	212	78	212
Salmon	7	0	7	7	7	7	7	0	0	7	0	0	7
<b>Total</b>	<b>438</b>	<b>431</b>	<b>437</b>	<b>438</b>	<b>437</b>	<b>437</b>	<b>426</b>	<b>130</b>	<b>216</b>	<b>222</b>	<b>431</b>	<b>137</b>	<b>438</b>

Table 3. Statistics associated with simple linear regressions using measurements (mm) of cranial lengths: upper jaw length ( $L_{UJ}$ ), eye diameter ( $L_{ED}$ ), interorbital distance ( $L_{ID}$ ), snout length ( $L_{SL}$ ), post orbital to preoperculum length ( $L_{PP}$ ) and post orbital head length ( $L_{PO}$ ) as predictors of the fork length ( $L_{FL}$ ) of 438 Sablefish collected from research surveys in 2016. All models were significant at  $P < 0.001$ .

Cranial Measurement	Sex	N	Slope	SE	$R^2$	r	Predictor		Response $L_{FL}$	
							Mean	SD	Mean	SD
$L_{UJ}$	female	215	7.4	0.12	0.945	0.972	59.8	16.93	586.7	129.71
	male	222	8.1	0.13	0.949	0.974	56.2	13.16	560.3	109.45
$L_{ED}$	female	216	23.9	0.5	0.914	0.956	25.9	5.19	586.7	129.41
	male	222	20.1	0.51	0.877	0.936	25.9	5.09	560.3	109.45
$L_{ID}$	female	215	11.3	0.15	0.96	0.98	41.6	11.27	586.9	129.68
	male	222	12.2	0.25	0.916	0.957	39.2	8.59	560.3	109.45
$L_{SL}$	female	215	10.9	0.14	0.966	0.983	46.6	11.7	586.7	129.71
	male	222	11.9	0.19	0.945	0.972	43.1	8.93	560.3	109.45
$L_{PP}$	female	211	13.8	0.23	0.945	0.972	32.6	9.03	583	127.89
	male	215	15.7	0.31	0.921	0.96	30.4	6.73	559.9	109.84
$L_{PO}$	female	73	6.6	0.23	0.923	0.961	62.1	22.51	574.9	154.35
	male	57	7.8	0.34	0.904	0.951	59.2	12.81	555.6	104.97

Table 4. Ease of use and repeatable five point ranking scale recorded by experienced samplers for each cranial measurement for the 2016 experimental study, where 5 = Great; 4 = Good; 3 = Moderate; 2 = Questionable; 1 = Poor.

Head Dimension	5 Point Rank		Measurement			Considerations
	Ease of use	Repeatable	Caliper limitation	Bilateral	Bone	
L <sub>UJ</sub>	3	4	x	x	x	End of the maxilla difficult to define. Caliper jaw position must be in center of snout.
L <sub>ED</sub>	3	2		x		Caliper outside jaw position on soft tissue in eye socket may result in measurement differences.
L <sub>ID</sub>	5	5			x	Tissue is compressed to obtain bone measurement. Easy to determine caliper jaw position.
L <sub>SL</sub>	4	5		x	x	Caliper jaw position must be in center of snout.
L <sub>PP</sub>	4	5		x	x	Caliper position on pre-operculum may result in measurement differences.
L <sub>PO</sub>	3	2	x	x		Operculum damage from handling was observed on several fish. Endpoints difficult to define. Electronic calipers malfunctioned with lengths >15 cm.

Table 5. Summary of Sablefish biological data collected from the 2017 pilot study to Cobb (n=60) and Eickelberg (n=39) seamounts, measured by three expert technicians using a standardized protocol for the 2017 pilot study. Index of Average Error (IAE) % values are reported for each of the cranial lengths.

Head Dimension	Sampler A	Sampler B	Sampler C	IAE
	Min - Max (mm)	Min - Max (mm)	Min - Max (mm)	
L <sub>UJ</sub>	51.65 - 91.02	52.64 - 91.53	53.82 - 92.55	1
L <sub>ID</sub>	35.43 - 61.18	35.91 - 60.81	34.94 - 60.72	1.1
L <sub>SL</sub>	39.67 - 69.27	40.38 - 70.37	41.03 - 70.32	1.2
L <sub>PP</sub>	27.69 - 49.88	26.83 - 52.51	27.25 - 52.15	2.3

## 7 Figures




Figure 1. Sample locations from the 2016 WCVI, 2016 WCHG research surveys and 2017 pilot study. The sample locations of the salmon survey are unknown.




Figure 2. Relationship between cranial lengths (UJ, ED, ID) vs fork length in millimeters. Predicted points represented by black circles, measured values colored by residual scale.





Figure 3. Relationship between cranial lengths (SL, PP, PO) vs fork length in millimeters. Predicted points represented by black circles, measured values colored by residual scale.

## APPENDIX A IMAGES OF THE SIX CRANIAL DIMENSION MEASUREMENTS.

A. Upper jaw measurement (UJ); B. Eye diameter measurement (ED); C. Interorbital distance (ID); D. Snout length (SL); E. Post orbital to preoperculum length measurement (PP); F. Post orbital head length (PO).



## APPENDIX B SEX DETERMINATION BY OPERCULUM MARKING


Instructions for sex determinations and operculum knife cuts for Sablefish males and females.

### MALES

Look for lobes that have an 'edge' to them. Due to the 'edge', they will not 'roll' easily in your fingers. The lobes are fused together at the end closest to the head.

Mark with one cut on the gill cover

**Male**



Sample photos for color and shape reference:

Juvenile: translucent-white



Developing: creamy-white



Post Spawning: brown or tan




### FEMALES

Look for tubes that are oval in cross section and will 'roll' in your fingers.

Mark with two cuts on the gill cover

**Female**



Sample photos for color and shape reference:

Juvenile: translucent pinky purple



Developing: creamy white, filled with eggs



Post Spawning: reddish-purple-brown, whitish sheen to the exterior surface



## APPENDIX C SABLEFISH BIOSAMPLE COLLECTION PROGRAM NOTICE

Instructions for the Sablefish biosample collection program.



Fisheries and Oceans  
Canada

Pêches et Océans  
Canada



## SABLEFISH BIOSAMPLE COLLECTION PROGRAM

By collecting biological samples (**HEADS ONLY**) from your Sablefish catch you play an important role in ensuring the Sablefish fishery continues to be a model for sustainable fisheries in Canada. Having an accurate understanding of the biological characteristics (e.g., size, age, sex) of Sablefish catch provides information that is critical to understanding the health of the stock, improving our ability to track changes in it (e.g., to identify when the population is growing), and reducing uncertainty in how management actions meet fishery and conservation objectives.

**SABLEFISH BIOSAMPLING KITS**  
(instructions, logbook, straps, bags) are available from  
Birgit Gettings at 778 227-0395 or [birgitg@archipelago.ca](mailto:birgitg@archipelago.ca)

**SABLEFISH BIOSAMPLE REWARDS!**  
**EACH SABLEFISH BIOSAMPLE WILL EARN 20 POINTS VALID TOWARD THE**  
**SABLEFISH REWARDS CATALOGUE**

### TO REDEEM YOUR REWARD POINTS

1. Contact: Birgit Gettings: [birgitg@archipelago.ca](mailto:birgitg@archipelago.ca) or 778-227-0395
2. **Vessel Authorized Person (AP):** Each vessel may appoint one AP to redeem points on behalf of the vessel. APs please register with Birgit to obtain the vessel's points balance. Sablefish Biosample reward points and Sablefish Tag reward points will be pooled for a combined total vessel rewards points balance. Note: points expire 3 years from the end of the year in which they were acquired.
3. **Select Rewards and Place Order:** Crews may browse the *Sablefish Rewards Catalogue* (<http://canadiansablefish.com/>) to view reward items and size charts and provide their selections to the vessel AP. When ready the APs may contact Birgit to place the order.
4. **Rewards Delivery:** If possible please select one of the AMR port offices listed in the shipping options for your rewards delivery. Orders will be fulfilled 4-5 times per year. Birgit will provide details on rewards delivery or availability for pick up.

## APPENDIX D TAGGED SABLEFISH PROGRAM NOTICE

Instructions for the tagged Sablefish collection program.




Fisheries and Oceans  
Canada

Pêches et Océans  
Canada



### TAGGED SABLEFISH PROGRAM



By returning tagged Sablefish (**HEADS and TAGS ONLY**) you play an important role in ensuring the Sablefish fishery continues to be a model for sustainable fisheries in Canada. Recoveries of tagged Sablefish provide critical information on fishery selectivity (i.e., what size Sablefish are most likely to be caught in each fishery) as well as information on movement and growth rates. This informs our understanding of the health of the stock, improves our ability to track changes in it, and reduces uncertainty in how management actions meet fishery and conservation objectives.

#### WHEN YOU CATCH A TAGGED SABLEFISH

J-cut it per commercial practice and place the HEAD ONLY, with tag tucked under gill cover, in an individual bag and freeze. Refer to the Sablefish Sex sheet for detailed guidance on determining the sex and marking of the fish.

*\*Remember to write the tag number in log book\**

At the end of your trip, bring the bag with the head and tag to your dockside validator. Vessel entries in the prize draw and reward points will be recorded and the fish head will be taken for biological sampling.

#### **TAGGED SABLEFISH REWARDS!**

**EACH TAGGED SABLEFISH HEAD/TAG RETURNED WILL BE ENTERED IN AN ANNUAL DRAW FOR FOUR CASH PRIZES OF \$2,000 EACH.**

**EACH TAGGED SABLEFISH WILL ALSO EARN 1 POINT VALID TOWARD REWARDS IN THE SABLEFISH REWARDS CATALOGUE.**

## 8 References

- Beamish, R., and Fournier, D.D.A. 1981. A method for comparing the precision of a set of age determinations. Canadian Journal of Fisheries and Aquatic Sciences 38: 982–983.
- Cox, S.P., Holt, K., and Johnson, S. 2019. Evaluating the robustness of management procedures for the Sablefish (*\*anoplopoma fimbria\**) fishery in British Columbia, Canada for 2017-18. DFO Can. Sci. Advis. Sec. Res. Doc. 2019/032. vi + 79 p.
- Haist, R.H., V., and Wyeth, M. 2001. Sablefish Stock Assessment for 2001 and Advice to Managers for 2002. DFO Can. Sci. Advis. Sec. Res. Doc. 2001/135.
- Isermann, D.A., and Vandergoot, C.S. 2005. Predicting walleye total length from head and mandible measurements. North American Journal of Fisheries Management 25(1): 316–321. Taylor & Francis.
- Nottingham, M.K., Williams, D.C., Wyeth, M.R., and Olsen, N. 2018. Summary of the west coast haida gwaii synoptic bottom trawl survey, august 25 - september 26, 2016. Can. Manuscr. Rep. Fish. Aquat. Sci. 3151: viii: 51 p.
- Park, I.S., Kim, Y.J., Choi, H.J., Oh, S.Y., Noh, C.H., and Lee, S.H. 2007. Total length estimation from head dimensions of artificially propagated brown croaker *miichthys miiuy*. Korean J. Ichthyol. 19: 128–131.
- R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Richardson, J., Shears, N., and Taylor, R. 2015. Using relative eye size to estimate the length of fish from a single camera image. Marine Ecology Progress Series 538.
- Rondeau, E.B., Messmer, A.M., Sanderson, D.S., Jantzen, S.G., Schalburg, K.R. von, Minkley, D.R., Leong, J.S., Macdonald, G.M., Davidsen, A.E., Parker, W.A., Mazzola, R.S.A., Campbell, B., and Koop, B.F. 2013. Genomics of sablefish (*anoplopoma fimbria*): Expressed genes, mitochondrial phylogeny, linkage map and identification of a putative sex gene. BMC Genomics 14(1): 452. Journal Article.
- Serafy, J.E., Schmitz, C.M., Capo, T.R., Clarke, M.E., and Ault, J.S. 1996. Total length estimation of red drum from head dimensions. The Progressive Fish-Culturist 58(4): 289–290. Taylor & Francis.
- Shaw, W., and McFarlane, G. 1997. Development of sablefish, *anoplopoma fimbria* larvae off the west coast of british columbia and transformation in the juvenile stage. NOAA Tech. Rep. NMFS 130: 3–12.
- Williams, D.C., Nottingham, M.K., Olsen, N., and Wyeth, M.R. 2018. Summary of the west coast vancouver island synoptic bottom trawl survey, may 24 - june 15, 2016. Can. Manuscr. Rep. Fish. Aquat. Sci. 3137: viii: 54 p.