
Enrollment
Allocation
Analysis
Initiated enrolment survey (N = 4,483)
Did not complete enrolment survey (n = 2,325)
Completed enrolment and randomised (N = 2,158)
Allocated to ABC (n = 522)
Allocated to ACC (n = 556)
Allocated to BCC (n = 552)
Allocated to CCC (n = 528)
Excluded: no intervention events (n = 135)
Excluded: no intervention events (n = 126)
Excluded: no intervention events (n = 143)
Excluded: no intervention events (n = 113)
Analysed: retention (RQ1; n = 387)
Analysed: retention (RQ1; n = 430)
Analysed: retention (RQ1; n = 409)
Analysed: retention (RQ1; n = 415)
Not active in both Weeks 1 & 6 (n = 251)
Not active in both Weeks 1 & 6 (n = 288)
Not active in both Weeks 1 & 6 (n = 288)
Not active in both Weeks 1 & 6 (n = 275)
Analysed: efficacy (RQ2; n = 136)
Analysed: efficacy (RQ2; n = 142)
Analysed: efficacy (RQ2; n = 121)
Analysed: efficacy (RQ2; n = 140)


ABC/ACC/BCC
transition
ABC
transition
40%
30%
20%
10%
0%
At−risk participants dropping out (%)
Week 1
Week 2
Week 3
Week 4
Week 5
Week 6
Study week
Condition
ABC
ACC
BCC
CCC


(A) Initial attempts
Estimated attempts at Week 6
(adjusted for Week 1 behaviour)
80
70
60
ABC
ACC
BCC
CCC
Condition
condition
ABC
ACC
BCC
CCC
(B) Dismissed openings
Estimated dismissal rate at Week 6, %
(adjusted for Week 1 behaviour)
27
25
23
ABC
ACC
BCC
CCC
Condition
condition
ABC
ACC
BCC
CCC


(A) Attempts
All three delays observed
No delay variation
Predicted weekly attempts
120
100
80
60
1
2
3
4
5
6
(B) Dismissed openings
Predicted dismissal rate
35%
30%
25%
20%
Delay (s)
3
6
10
(C) Active participants by delay
Active participants
1500
1000
500
0
Study week


100%
90%
80%
70%
60%
50%
40%
Predicted Pr(dropout by Week 6)
ABC
ACC
BCC
CCC
Condition
Analysis
Preregistered
(logistic regression)
Exploratory
(dose−response hazard model)


0%
−20%
−40%
−60%
Change in weekly attempts, Week 1 ... Week 6
ABC
ACC
BCC
CCC
Condition
Analysis
Preregistered (PP, efficacy)
Exploratory (ITT, effectiveness)