

STAT 429: Introduction to Time Series Analysis

Homework One

Fall 2026, University of Illinois Urbana-Champaign

Total Points = 104; Maximum Score = 100

For problems involving data analysis, you are free to use inbuilt functions from libraries in R or Python. Attach code snippets in your solutions corresponding to each problem.

1. Given data $y_1, \dots, y_n$, consider the model

$$y_i \stackrel{\text{i.i.d.}}{\sim} N(\mu, \sigma^2)$$

for two unknown parameters μ and $\sigma > 0$. We can also write this model as

$$y_i = \mu + \epsilon_i \quad \text{with } \epsilon_i \stackrel{\text{i.i.d.}}{\sim} N(0, \sigma^2).$$

Note that this is a special case of the simple linear regression model with all covariate values $x_i = 0$ (and β_0 is now denoted by μ).

- a) Write down the likelihood function, and calculate the Maximum Likelihood Estimates of μ and σ . (**1 + 2 = 3 points**)

- b) Working with the prior

$$\mu, \log \sigma \stackrel{\text{i.i.d.}}{\sim} \text{Unif}(-C, C)$$

for a large $C > 0$, write down the posterior density of μ, σ . It is okay to express the answer up to a proportionality constant. (**2 points**)

- c) Show that the posterior density of μ satisfies

$$f_{\mu|\text{data}}(\mu) \propto \frac{I\{-C < \mu < C\}}{[\sum_{i=1}^n (y_i - \mu)^2]^{n/2}},$$

where data represents $y_1, \dots, y_n$. Use $C \rightarrow \infty$ in the integration limits while calculating the necessary integral. (**3 points**)

- d) Use the posterior density of μ derived in the previous part to show that

$$\frac{\sqrt{n}(\mu - \bar{y})}{s} \mid \text{data} \sim t_{n-1}, \tag{1}$$

where

$$\bar{y} := \frac{y_1 + \dots + y_n}{n}, \quad s^2 := \frac{1}{n-1} \sum_{i=1}^n (y_i - \bar{y})^2.$$

Note that $\bar{y}$ and s are purely functions of the data $y_1, \dots, y_n$, so they become constants when conditioned on the data. (**3 points**)

- e) Using (1), argue that the posterior mean (assuming $n \geq 3$), posterior median, and posterior mode of μ are all equal to $\bar{y}$. This means that the mean $\bar{y}$ is the most natural Bayesian estimate of μ . **(1 point)**
- f) Using (1), argue that the posterior probability that μ belongs to the interval

$$\left[\bar{y} - \frac{s}{\sqrt{n}} t_{n-1, \alpha/2}, \bar{y} + \frac{s}{\sqrt{n}} t_{n-1, \alpha/2} \right] \quad (2)$$

equals $1 - \alpha$. Here $t_{n-1, \alpha/2}$ is the point for which the t_{n-1} density gives probability $\alpha/2$ to values larger than $t_{n-1, \alpha/2}$. The interval (2) is the Bayesian uncertainty interval for μ . **(1 point)**

- g) Download the dataset on the Consumer Price Index (CPI) from <https://fred.stlouisfed.org/series/CPIAUCSL>. This is the same dataset that we considered in class. Let C_t denote the monthly CPI value at time t , and define

$$y_t = \log C_{t+1} - \log C_t.$$

Use all available monthly observations and fit the model

$$y_t \stackrel{\text{i.i.d.}}{\sim} N(\mu, \sigma^2)$$

to this data. What is the interpretation of μ ? **(1 point)**

- h) Calculate the estimate and uncertainty interval (from the previous two parts) for μ . **(3 points)**
- i) Compare the estimate and interval for μ obtained above with the estimate and interval for β_1 from the model

$$\log C_t = \beta_0 + \beta_1 t + \epsilon_t, \quad \epsilon_t \stackrel{\text{i.i.d.}}{\sim} N(0, \sigma^2).$$

Which of these intervals would you prefer for the historical month-to-month inflation? Explain your reasoning. **(2 points)**

2. Download the dataset on the annual size of the Resident Population of California from <https://fred.stlouisfed.org/series/CAPOP>. This dataset gives the annual population of California, with units in thousands of persons. Drop the most recent observation and use the remaining observations as the training data. The goal is to predict the held-out final observation.

- a) Fit the model

$$y_t = \beta_0 + \beta_1 t + \epsilon_t \quad \text{with } \epsilon_t \stackrel{\text{i.i.d.}}{\sim} N(0, \sigma^2). \quad (3)$$

Provide point estimates for β_0, β_1 along with appropriate uncertainty intervals. Interpret your point estimates and explain why they make sense. **(4 points)**

- b) Along with a plot of the observed dataset, plot lines corresponding to 100 samples from the posterior distribution of (β_0, β_1) under the prior

$$\beta_0, \beta_1, \log \sigma \stackrel{\text{i.i.d.}}{\sim} \text{Unif}(-C, C)$$

for a very large C . Comment on the range of uncertainty revealed in this plot. **(4 points)**

- c) Based on model (3), provide a point estimate along with appropriate uncertainty quantification for the held-out final population observation. **(2 points)**
- d) Fit the model

$$\log y_t = \beta_0 + \beta_1 t + \epsilon_t \quad \text{with } \epsilon_t \stackrel{\text{i.i.d.}}{\sim} N(0, \sigma^2). \quad (4)$$

Provide point estimates for β_0, β_1 along with appropriate uncertainty intervals. Interpret your point estimates and explain why they make sense. **(4 points)**

- e) For model (4), under the prior

$$\beta_0, \beta_1, \log \sigma \stackrel{\text{i.i.d.}}{\sim} \text{Unif}(-C, C)$$

for a very large C , calculate the posterior probability that β_1 is larger than 0.03. **(2 points)**

- f) Based on model (4), provide a point estimate along with appropriate uncertainty quantification for the held-out final population observation. **(2 points)**
- g) Fit the model

$$\log y_t = \beta_0 + \beta_1 t + \beta_2 t^2 + \epsilon_t \quad \text{with } \epsilon_t \stackrel{\text{i.i.d.}}{\sim} N(0, \sigma^2). \quad (5)$$

Based on the fitted model, provide a point estimate along with appropriate uncertainty quantification for the held-out final population observation. **(2 points)**

- h) Which of the three models (3), (4), and (5) would you recommend for this dataset and why? Also report which of your predictions for the held-out population observation is closest to the actual observed value. **(3 points)**

3. Download the FRED dataset from <https://fred.stlouisfed.org/series/MRTSSM4453USN>, which gives monthly data on beer, wine, and liquor sales in the United States. Let y_t denote the observed data value for time t .

- a) Is the linear model

$$\log y_t = \beta_0 + \beta_1 t + \epsilon_t \quad \text{with } \epsilon_t \stackrel{\text{i.i.d.}}{\sim} N(0, \sigma^2) \quad (6)$$

appropriate for this dataset? Why or why not? **(2 points)**

- b) Let t_0 be the time point corresponding to April 2020. Fit the model

$$\log y_t = \beta_0 + \beta_1 t + \beta_2 I\{t \geq t_0\} + \beta_3 t I\{t \geq t_0\} + \epsilon_t \quad (7)$$

with

$$\epsilon_t \stackrel{\text{i.i.d.}}{\sim} N(0, \sigma^2)$$

to this dataset.

Here $I\{t \geq t_0\}$ denotes the variable which takes the value 0 for $t < t_0$ and 1 for $t \geq t_0$, and $tI\{t \geq t_0\}$ denotes the variable which takes the value 0 for $t < t_0$ and t for $t \geq t_0$.

On a single figure, plot the fitted values of models (6) and (7) along with the original data. Based on this plot, is (7) a better model for this dataset compared to (6)? Why or why not? **(4 points)**

c) In model (7), along with the prior

$$\beta_0, \beta_1, \beta_2, \beta_3, \log \sigma \stackrel{\text{i.i.d.}}{\sim} \text{Unif}(-C, C)$$

for a very large C , calculate the posterior probability that $\beta_3 < 0$. Also calculate the posterior probability that

$$\beta_2 + \beta_3 t_0 > 0.$$

(6 points)

d) For an appropriate $k \leq 4$, fit the model

$$y_t = \beta_0 + \beta_1 t + \beta_2 I\{t \geq t_0\} + \beta_3 t I\{t \geq t_0\} + \sum_{i=1}^k \left(a_i \cos\left(2\pi \frac{it}{12}\right) + b_i \sin\left(2\pi \frac{it}{12}\right) \right) + \epsilon_t. \quad (8)$$

Give reasons for your choice of k . **(4 points)**

e) Using model (8) with your chosen k , predict the sales data for the next 36 future months. Plot the predictions along with the original data and explain why your forecast is reasonable. **(4 points)**

4. Download the Google Trends time series dataset for the query *yahoo*. Use a monthly time series of worldwide search popularity beginning in January 2004.

The goal of this exercise is to fit the polynomial trend model

$$y_t = \beta_0 + \beta_1 t + \dots + \beta_k t^k + \epsilon_t \quad \text{with } \epsilon_t \stackrel{\text{i.i.d.}}{\sim} N(0, \sigma^2)$$

for an appropriate value of $k \leq 5$.

a) Visually evaluate the fit of the least squares estimate for this model to the observed data to pick an appropriate value of $k \leq 5$. Explain the reason for your choice of k . **(5 points)**

b) On a plot of the observed dataset, plot the polynomial corresponding to the least squares estimate for the model with your chosen value of k . On the same figure, plot polynomials corresponding to 300 samples from the posterior distribution of the coefficients. Comment on the range of uncertainty revealed in this plot. **(5 points)**

5. Consider the multiple regression model

$$y_i = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \dots + \beta_m x_{im} + \epsilon_i \quad \text{with } \epsilon_i \stackrel{\text{i.i.d.}}{\sim} N(0, \sigma^2). \quad (9)$$

We observe data

$$(x_{i1}, \dots, x_{im}, y_i), \quad i = 1, \dots, n.$$

In the model above, the covariates x_{ij} are treated as fixed constants.

For Bayesian analysis with the prior

$$\beta_0, \beta_1, \dots, \beta_m, \log \sigma \stackrel{\text{i.i.d.}}{\sim} \text{Unif}(-C, C),$$

we showed in class that when $C \rightarrow \infty$,

$$\beta \mid \text{data} \sim t_{n-m-1, m+1} \left(\hat{\beta}, \hat{\sigma}^2 (X^T X)^{-1} \right), \quad (10)$$

where $X, \beta, \hat{\beta}, \hat{\sigma}$ are all as defined in the lecture notes.

This exercise provides another slightly different proof of this same result. Along the way, it also discusses inference for the parameter σ . Throughout, assume that $C \rightarrow \infty$ (i.e., C is very large).

a) Prove that

$$\beta \mid \text{data}, \sigma \sim N_{m+1} \left(\hat{\beta}, \sigma^2 (X^T X)^{-1} \right). \quad (11)$$

The left-hand side above refers to the conditional density of β given the data as well as the parameter σ . **(4 points)**

b) Prove that the posterior density of σ (i.e., the density of $\sigma \mid \text{data}$) is proportional to

$$\sigma^{-n+m} \exp \left(-\frac{S(\hat{\beta})}{2\sigma^2} \right) I\{\sigma > 0\}, \quad (12)$$

where $S(\hat{\beta})$ is the Residual Sum of Squares. **(4 points)**

c) Use the results of the previous two parts along with the formula

$$f_{\beta \mid \text{data}}(\beta) = \int_0^\infty f_{\beta \mid \text{data}, \sigma}(\beta) f_{\sigma \mid \text{data}}(\sigma) d\sigma$$

to deduce (10). **(4 points)**

d) Use the result in part (b) to show that

$$\frac{S(\hat{\beta})}{\sigma^2} \mid \text{data} \sim \chi_{n-m-1}^2.$$

(3 points)

e) Use the result in the previous part to argue that the usual estimator

$$\hat{\sigma} := \sqrt{\frac{S(\hat{\beta})}{n-m-1}}$$

is a reasonable point estimate of σ . **(2 points)**

6. This problem discusses prediction. Consider the same setting as the previous problem.

a) Suppose we are interested in the parameter $a^T \beta$ for some fixed vector a . Using (11), prove that

$$a^T \beta \mid \text{data}, \sigma \sim N \left(a^T \hat{\beta}, \sigma^2 a^T (X^T X)^{-1} a \right).$$

Note that we are conditioning on the data as well as the parameter σ on the left-hand side above. **(3 points)**

b) Using (10), prove that

$$a^T \beta \mid \text{data} \sim t_{n-m-1} \left(a^T \hat{\beta}, \hat{\sigma}^2 a^T (X^T X)^{-1} a \right).$$

(3 points)

- c) Suppose we are interested in predicting the value Y_{n+1} of the response variable at a new set of covariate observations $x_{n+1,1}, \dots, x_{n+1,m}$. Assuming that

$$Y_{n+1} = \beta_0 + \beta_1 x_{n+1,1} + \dots + \beta_m x_{n+1,m} + \epsilon_{n+1}$$

for an independent error

$$\epsilon_{n+1} \sim N(0, \sigma^2),$$

prove that

$$Y_{n+1} \mid \text{data}, \sigma \sim N\left(\tilde{x}_{n+1}^T \hat{\beta}, \sigma^2 + \sigma^2 \tilde{x}_{n+1}^T (X^T X)^{-1} \tilde{x}_{n+1}\right),$$

where $\tilde{x}_{n+1}$ is the vector with entries $1, x_{n+1,1}, \dots, x_{n+1,m}$. **(4 points)**

- d) Using the result from the previous part, the formula

$$f_{Y_{n+1} \mid \text{data}}(y) = \int_0^\infty f_{Y_{n+1} \mid \text{data}, \sigma}(y) f_{\sigma \mid \text{data}}(\sigma) d\sigma,$$

and the density $\sigma \mid \text{data}$ from (12), prove that

$$Y_{n+1} \mid \text{data} \sim t_{n-m-1}\left(\tilde{x}_{n+1}^T \hat{\beta}, \hat{\sigma}^2 + \hat{\sigma}^2 \tilde{x}_{n+1}^T (X^T X)^{-1} \tilde{x}_{n+1}\right).$$

This result can be used for obtaining predictions of Y_{n+1} along with uncertainty intervals. **(5 points)**