QICT INDEPENDENT PHYSICAL-POLE EXECUTION
Mathematical specification and fail-closed handoff â 19 October 2026
Status: BLOCKED_MISSING_INPUT
Repository: github.com/sachamed/qict-code-data
Inspected reference commit: fd664503ca92d2d88007d68d6e3873f7df2beeb8
Canonical action SHA-256: c22ff80fcb94204984a65debf7e5636ef1d4e6caad89cf8fd6934dfdb8f684b7

1. FROZEN DISCRETE-TIME ACTION
The canonical JSON fixes compact gauge U(1), SU(2), SU(3), periodic BCC geometry
and a microscopic product of swaps, cell transfer, electric, magnetic and matter factors.
The primitive collision phase is pi/63. Sources follow L1,L2,L3,R1,R2,R3.
The physical contour, interacting global U-cone and independent clock are not fixed.

2. SPECTRAL OBSERVABLES
For a full physical W and isometry V, moments are C_n = V* W^n V, with C_0 = I_6.
The matrix resolvent is G(z) = V* (z-W)^(-1) V, outside unit spectrum.
Toeplitz positivity is necessary; finitely many moments do not identify a pole.
A physical mode needs spectral isolation, residue >= 0 and nonzero visible weight.
It also needs volume, buffer, regulator and quasienergy-branch stability.

3. AUTHENTIC EVIDENCE, NOT COMPLETE INPUT
The L24 N9 rank-six data are genuine PRECONTRACTION payloads, not a full U.
Nested buffer geometry does not give four certified Kato density matrices.
C10 shell geometry is not a complete CPTP/Kraus quantum map.
Synthetic control CN6 spectra are not physical six-source moments.

4. PROHIBITED INFERENCES
No physical pole, residue, mass in GeV, relic abundance or gravitational coupling
has been produced by this preflight. A read-I/O benchmark is not an apply_U benchmark.

QICT â ACCEPTANCE GATES AND COMPUTATIONAL ORDER

5. REQUIRED CHAIN

Kato/Riesz densities and certified tail -> C10 quantum maps -> boundary-to-path
-> SU(2)/SU(3)/CAR core recoupling -> rank-six source isometry -> full U and U*
-> physical CN6 -> error-certified Toeplitz -> isolated pole and positive residue
-> multi-volume and regulator checks -> independent clock -> physical mass.

6. REQUIRED NUMERICAL AND MATHEMATICAL CHECKS

Each Kato rho = F F* / Tr(F F*) must be positive and trace-one.
An independently certified strictly positive Riesz gap and tail bound are required.
The full unitary must pass matvec, adjointness, and norm-preservation tests.
Six source columns must satisfy V*V = I_6.
C_0 = I_6 and block Toeplitz >= 0 require propagated error control.
Finite-volume eigenphases alone are not thermodynamic particle evidence.

7. MULTI-VOLUME PROTOCOL

L=24,32,48,64,96,128; compatible depths n=9,13,21,29,45,61.
Compare volume laws, buffer bounds, regularization and extraction methods.
Do not use an external target mass to define the QICT scale.

8. DARK SECTOR IDENTIFIABILITY

If W=W_charged direct-sum W_dark and V=(V_charged,0), then all charged
moments V*W^nV are independent of the hidden W_dark spectrum.
Therefore a physical neutral copy-odd source V_chi must be constructed
and shown to have nonzero projected overlap with an isolated mode.

9. INDEPENDENT EXECUTION POLICY

Fix action, author-designated commit, source conventions and toolchain hashes.
Run tests and fail-closed guards before requesting HPC resources.
A physical mass can be reported only after fully independent verification.

FINAL DECISION: BLOCKED_MISSING_INPUT â no QICT dark-particle mass prediction.
This PDF is a Github-readable handoff; see the original external audit PDF
for the full two-page documentary rendering retained in the local ZIP.