TOPIC SCOPE

Yes. **You absolutely can build a useful first version without needing temporal imagery for the whole Earth.** In fact, I think this makes the project more feasible.

The key is to distinguish **“ecological condition assessment”** from **“change detection.”**

1. You don't need to start with temporal data

A single satellite observation can tell you a lot about the **current state** of a landscape.

For example:

```
Satellite image
      ↓
Deep-learning CV
      ↓
Land-cover / ecological analysis
      ↓
┌──────────────────────────────┐
│ Forest       42%             │
│ Cropland     18%             │
│ Built-up     24%             │
│ Water         6%             │
│ Bare land    10%             │
└──────────────────────────────┘
      ↓
Environmental indicators
      ↓
Ecological condition / risk
```

There are already global Sentinel-2 datasets specifically designed for this. **Sentinel2GlobalLULC**, for example, contains ~195,000 globally distributed 10 m tiles across 29 land-use/land-cover classes. (Nature)

Another dataset, **Sen-2 LULC**, contains 213,761 Sentinel-2 images/masks covering water, forests, barren land, built-up, agriculture and other classes. (ScienceDirect)

So you can train a segmentation/classification model without collecting your own imagery.

---

2. But don't call it "ecological degradation detection" from one image

This is the important scientific limitation.

From one image, you can say:

> **"This area currently exhibits characteristics associated with ecological stress/degradation."**

You generally **cannot** say:

> "The environment degraded by 23%."

That requires a baseline or historical comparison.

For example, a desert naturally has lots of bare land. A city naturally has lots of concrete. A satellite image alone cannot tell you whether those conditions are **getting worse**.

So I'd structure your system into two levels:

Level 1 — Single-image ecological assessment

What is the state of this area right now?

Level 2 — Temporal change detection

How has this state changed over time?

You can implement Level 1 for the entire project and make Level 2 an extension.

---

3. There is actually a very interesting way to do Level 1

Instead of just:

> "Classify forest / water / urban / desert."

you could create an **Ecological Condition Assessment Model**.

Input:

one Sentinel-2 scene

Output:

```
ECOLOGICAL CONDITION

Vegetation          ███████▒  78
Water presence      ████▒▒▒▒  51
Built-up pressure   ████████▒  89
Bare-soil exposure  ██▒▒▒▒▒▒  32
Vegetation stress   ██████▒▒  61

Overall ecological
condition            64 / 100

Classification:
MODERATE STRESS
```

But the score must be constructed scientifically rather than arbitrarily.

There is already research around **Remote Sensing Ecological Index (RSEI)**, which combines remotely sensed indicators such as greenness, humidity, dryness and heat into an ecological-quality measure. Researchers have also been developing deep-learning versions because traditional indices have limitations. (Taylor & Francis Online)

So you have a legitimate scientific foundation.

---


4. Here's where your project can become novel

Don't simply reproduce RSEI.

Ask:

> **Can deep-learning computer vision learn an ecological-condition representation from satellite imagery instead of relying entirely on hand-designed indices?**

For example:



That is much more interesting than ordinary image classification.

---

5. You could even make it multi-class

For example:

Input

A 10 km × 10 km satellite region.

Model outputs

```
FOREST
████████████████ 51%

AGRICULTURE
██████           22%

URBAN
████             13%

WATER
██                7%

BARE LAND
██                7%
```

Then a second model/algorithm interprets the composition:

```
Vegetation coverage:       HIGH
Human modification:        MEDIUM
Water availability:        MEDIUM
Built-up pressure:         HIGH
Potential ecological
stress:                    MODERATE
```

That is a **classification + interpretation system**, rather than just a classifier.

---

6. And you can still get temporal data later


Your concern about "we can't possibly obtain historical imagery for the whole Earth" is actually less of a problem than it sounds.

There are already global land-cover products with **annual maps going back decades**. The Copernicus global land-cover dataset, for example, provides global land-cover maps from **1992 to the present**, with yearly coverage. (Climate Data Store)

There are also annual 10 m global LULC products for **2017–2023** derived from Sentinel-2. (Open Data on AWS)

And the satellites themselves have extensive archives.

So eventually you could query:



You **don't need to download the entire Earth's history**.

The system can retrieve the relevant tiles for whatever location the user selects.

7. In fact, I'd change our project strategy slightly

Earlier we were considering:

> **Domain-generalized semantic change detection**

I would now make the research architecture:

Core research

> **Deep-learning-based ecological condition assessment from satellite imagery**

Optional temporal extension

> **Temporal change detection for identifying environmental degradation.**

That gives you a safety net.

If temporal modeling turns out to be too difficult:

You still have a complete project.

If you successfully implement it:

You have a significantly stronger project.

---

8. There is another interesting research direction: single-image + multiple spectral bands

A normal RGB image gives you:

```
R G B
```

Sentinel-2 gives you much more spectral information.

You can exploit:

```
Blue
Green
Red
Red Edge 1
Red Edge 2
Red Edge 3
NIR
SWIR
...
```

This allows models to distinguish things that visually look similar in RGB.

And this is directly relevant to ecological assessment because vegetation, soil, water and built surfaces have different spectral signatures.

A recent LULC study comparing deep-learning segmentation architectures is already using multiple land-cover categories such as built-up, roads, water, agriculture, shrubland, forest and grassland. (DOI)

---

9. The project could therefore have three levels

MVP — definitely achievable

Single satellite image → semantic segmentation/classification

```
Forest
Water
Agriculture
Urban
Bare land
```

```
Grassland
...
```

Strong project

Single image → ecological condition assessment

```
Land cover
+
Vegetation indicators
+
Human modification
+
Water
+
spectral information
        ↓
Ecological condition classification
```

Research-heavy extension

Multiple dates → environmental change detection

```
2017 ── 2020 ── 2023 ── 2026
  │       │       │       │
  └───────┴───────┴───────┘
              ↓
        CHANGE ANALYSIS
              ↓
     Forest loss
     Urban expansion
     Water change
     Vegetation degradation
```

You don't have to commit to the third one immediately.

---

10. And this actually fits your OSINT vision better

Your eventual website could show:

Current Earth

> "What does this place look like environmentally?"

Earth over time

> "How has it changed?"

Earth risk

> "What indicators suggest ecological stress?"

So:

```
             EARTH OSINT
                  │
      ┌───────────┴──────────┐
      ↓                      ↓
CURRENT CONDITION        HISTORICAL CHANGE
      │                      │
Land cover              Deforestation
Vegetation              Urbanization
Water                   Desertification
Built-up                Flooding
Human pressure          Coastal change
      │                      │
      └──────────┬───────────┘
                 ↓
```

```
            ENVIRONMENTAL
            INTELLIGENCE
```

My recommendation

For your **college project**, I would now frame the second sentence as:

> **“Application Focus: Deep Learning-Based Satellite Image Analysis for Automated Land-Cover and Ecological Condition Assessment.”**

Then the research question becomes:

> **Can deep-learning-based computer vision infer meaningful ecological condition indicators from multispectral satellite imagery, without requiring long-term temporal data?**

And **temporal change detection becomes an extension**, not a requirement.

That is safer technically, gives you huge datasets immediately, and still leaves the door open to the much larger Earth-OSINT platform you want to eventually build.

One particularly interesting finding from the literature is that even **single-date imagery has limitations**: a study comparing single-date and multi-date imagery found that medium-resolution single-date imagery was insufficient for some detailed post-deforestation land-use classification, while adding spatial detail or temporal information improved performance. (Taylor & Francis Online)

That limitation itself could become part of your research: **investigate how much ecological information can actually be extracted from a single multispectral observation, and where temporal information becomes necessary.**